
Safety Assessment of Fossil and Synthetic Waxes as Used in Cosmetics

Status: Re-Review for Panel Consideration
Release Date: August 15, 2025
Panel Meeting Date: September 8-9, 2025

History

Original Safety Assessment – published 1984

Original Re-Review – published 2005

Most Recent Action – new data being considered at this meeting

The Expert Panel for Cosmetic Ingredient Safety members are: Chair, Wilma F. Bergfeld, M.D., F.A.C.P.; Donald V. Belsito, M.D.; David E. Cohen, M.D.; Samuel M. Cohen, M.D., Ph.D.; Curtis D. Klaassen, Ph.D.; Allan E. Rettie, Ph.D.; David Ross, Ph.D.; Paul W. Snyder, D.V.M., Ph.D.; and Susan C. Tilton, Ph.D. The Cosmetic Ingredient Review (CIR) Executive Director is Bart Heldreth, Ph.D., and the Senior Director is Monice Fiume, M.B.A. This re-review document was prepared by Temima Nguyen, M.S., Scientific Analyst/Writer, CIR.



Commitment & Credibility since 1976

Memorandum

To: Expert Panel for Cosmetic Ingredient Safety Members and Liaisons
From: Temima Nguyen, M.S. Scientific Analyst/Writer, CIR
Date: August 15, 2025
Subject: Re-Review of the Safety Assessment of Fossil and Synthetic Waxes

The Expert Panel for Cosmetic Ingredient Safety (Panel) first published a review of the safety of Fossil and Synthetic Waxes in 1984 (identified as *originalreport1984_Fossil&SyntheticWaxes_092025* in the pdf). The Panel concluded these 8 ingredients are safe in cosmetics under the present practices of concentration and use. The Panel previously considered a re-review of this report and re-affirmed the 1984 conclusion, as published in 2005 (*rereview2005_Fossil&SyntheticWaxes_092025*).

Because it has been at least 15 years since the previous re-review was published, in accordance with Cosmetic Ingredient Review (CIR) Procedures, the Panel should consider whether the safety assessment of Fossil and Synthetic Waxes should be re-opened. In July 2025, an extensive search of the world's literature was performed for studies dated 2000 forward. The ingredients included for consideration for re-review, the original report journal citation, the original conclusion, a comparison of previous and new use data, and the search strategy are provided in the history and search strategy document (*history&search_Fossil&SyntheticWaxes_092025*).

Included for your review in Table 1 are updated and historical use data (*usetable_Fossil&SyntheticWaxes_092025*). According to a comparison of VCRP data from 2002 and 2023, the frequency of use of most (5/8) of the ingredients decreased. The following provides a comparison of the VCRP data from those years, with 2002 data compared to 2023 data (2002/2023): Ceresin (404/170); Emulsifying Wax NF (102/5); Microcrystalline Wax (548/1130); Montan Wax (13/3); Ozokerite (680/371); Paraffin (481/737); Synthetic Beeswax (179/164); and Synthetic Wax (215/1050). (Of note, Synthetic Wax has the highest frequency of use in RLD, with 15,166 reported uses.) In 2003, Paraffin had the highest concentration of use, at 99% in other skin care preparations; in 2022, the maximum reported concentration of use for Paraffin decreased notably to 21% in bath soaps and body washes (with the highest leave-on concentration being 18.6% in eyeliners). Overall, the ingredient with the highest reported concentration of use reported in response to the 2022 survey is Synthetic Wax, at 36% in eyebrow pencils. There are no notable changes in exposure type across this family of ingredients.

Also included is a summary of all new data that were identified as a result of the literature search. (See Tables 2 – 9; *newdata_Fossil & Synthetic Waxes_092025*.) In the new data, evidence of granuloma formulation and inflammation from Paraffin were noted in the repeated-dose toxicity studies section of the tables referenced above. However, there was no indication of new pertinent toxicity information for the other ingredients in this review.

If upon review of the updated use data and the new studies the Panel determines that a re-review is warranted, a full Draft Amended Report will be presented at an upcoming meeting.

Re-Review - Fossil & Synthetic Waxes (Ozokerite, etc.) - History and Search Strategy

Temima Nguyen – September 2025

Ingredients	Citation	Conclusion	Updated Use	Historical Use
<i>Ceresin</i> <i>Emulsifying Wax NF</i> <i>Microcrystalline Wax</i> <i>Montan Wax</i> <i>Ozokerite</i> <i>Paraffin</i> <i>Synthetic Beeswax</i> <i>Synthetic Wax</i>	JACT 3(3):43-99, 1984 IJT 24(Suppl. 1):67-74, 2005	Safe as used. Re-affirmed original conclusion.	<u>Ceresin</u> frequency of use (RLD, 2024): 5859 frequency of use (VCRP, 2023): 170 conc of use (2022): ≤ 10% <u>Emulsifying Wax NF</u> frequency of use (RLD, 2024):108 frequency of use (VCRP, 2023): 5 conc of use (2022): ≤ 1.5% <u>Microcrystalline Wax</u> frequency of use (RLD, 2024): 10,285 frequency of use (VCRP, 2023): 1130 conc of use (2022): ≤ 34.5% <u>Montan Wax</u> frequency of use (RLD, 2024): 34 frequency of use (VCRP, 2023): 3 conc of use (2022): ≤ 0.54% <u>Ozokerite</u> frequency of use (RLD, 2024): 8498 frequency of use (VCRP, 2023): 371 conc of use (2022): ≤ 23.3% <u>Paraffin</u> frequency of use (RLD, 2024): 7243 frequency of use (VCRP, 2023): 737 conc of use (2022): ≤ 21% <u>Synthetic Beeswax</u> frequency of use (RLD, 2024): 5602 frequency of use (VCRP, 2023): 164 conc of use (2022): ≤ 24.9% <u>Synthetic Wax</u> frequency of use (RLD, 2024): 15,166 frequency of use (VCRP, 2023): 1050 conc of use (2022): ≤ 36%	<u>Ceresin</u> frequency of use (2002): 404 conc of use (2003): ≤ 20% <u>Emulsifying Wax NF</u> frequency of use (2002): 102 conc of use (2003): ≤ 21% <u>Microcrystalline Wax</u> frequency of use (2002): 581 conc of use (2003): ≤ 50% <u>Montan Wax</u> frequency of use (2002): 13 conc of use (2003): ≤ 11% <u>Ozokerite</u> frequency of use (2002): 680 conc of use (2003): ≤ 22% <u>Paraffin</u> frequency of use (2002): 481 conc of use (2003): ≤ 99% <u>Synthetic Beeswax</u> frequency of use (2002): 179 conc of use (2003): ≤ 18% <u>Synthetic Wax</u> frequency of use (2002): 215 conc of use (2003): ≤ 29%

Search (from 2000 forward)

((Ozokerite) OR (12198-93-5) OR (64742-33-2)); hits/total # of useful hits: 1/16

((Ceresin) OR (8001-75-0)); hits/total # of useful hits: 1/8

((Montan wax) OR (8002-53-7)); hits/total # of useful hits: 1/7

((Paraffin) OR (8002-74-2) AND (toxicity)); hits/total # of useful hits: 13/1,173

((Microcrystalline Wax) OR (63231-60-7)); hits/total # of useful hits: 1/55

(Emulsifying Wax NF); hits/total # of useful hits: 0/1

((Synthetic Wax) OR (2658498-20-3) OR (8002-74-2) OR (9002-88-4) AND (toxicity) NOT (microplastic)); hits/total # of useful hits: 1/412

((Synthetic Beeswax) OR (71243-51-1) AND (toxicity)); hits/total # of useful hits: 0/72

Table 1. Frequency (RLD/VCRP) and concentration of use according to likely duration and exposure and by product category

	# of Uses			Max Conc of Use		# of Uses			Max Conc of Use	
	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵
Totals*	5859	170	404	2-10	0.8-20	108	5	102	1.5	0.5-21
summarized by likely duration and exposure**										
Duration of Use										
Leave-On	***	165	386	2-10	0.8-20	***	4	66	NR	0.5-12
Rinse-Off	***	5	18	NR	NR ¹	***	1	36	1.5	0.8-21
Diluted for (Bath) Use	***	NR	NR	NR	NR	***	NR	NR	NR	NR
Exposure Type										
Eye Area	***	40	121	7-10	4-12	***	NR	3	NR	NR
Incidental Ingestion	***	61	191	6.6	0.8-20	***	NR	NR	NR	NR
Incidental Inhalation-Spray	***	24 ^a ; 4 ^c	1; 21 ^a ; 6 ^c	8.2 ^a	2-13 ^a ; 5-8 ^c	***	4 ^a	4; 35 ^a ; 12 ^c	NR	1-12 ^a ; 1-10 ^c
Incidental Inhalation-Powder	***	4	6 ^c	NR	6; 5-8 ^c	***	NR	2 ^b	NR	NR
Dermal Contact	***	90	195	2-10	3-13	***	5	66	1.5	NR
Deodorant (underarm)	***	NR	NR	NR	NR	***	NR	NR	NR	NR
Hair - Non-Coloring	***	16	5	8.2	2-17	***	NR	26	NR	0.8-21
Hair-Coloring	***	NR	1	NR	NR	***	NR	7	NR	NR
Nail	***	NR	3	NR	4	***	NR	3	NR	0.5
Mucous Membrane	***	62	191	6.6	0.8-20	***	1	NR	NR	NR
Baby Products	***	1	NR	NR	NR	***	NR	2	NR	NR
as reported by product category										
Baby Products	1					2				
Baby Lotions/Oils/Powders/Creams	1	NR	NR	NR	NR	1	NR	2	NR	NR
Other Baby Products	NR	1	NR	NR	NR	1 (r.o.)	NR	NR	NR	NR
Bath Preparations	3									
Bath Oils, Tablets, and Salts	1	NR	NR	NR	NR					
Other Bath Preparations	2	NR	NR	NR	NR					
Eye Makeup Preparations (not children's)	1159									
Eyebrow Pencil	257	4	29	10	8					
Eyeliner	455	13	43	7	5-12	NR	NR	2	NR	NR
Eye Shadow	303	15	31	10	11					
Eye Lotion	2	NR	2	NR	NR					
Eye Makeup Remover	1	NR	3	NR	NR					
False Eyelashes	1	NA	NA	NR	NA					
Mascara	128	3	9	NR	4					
Eyelash and Eyebrow Adhesives, Glues, and Sealants	9	NA	NA	NR	NA					
Eyelash and Eyebrow Preparations (primers, conditioners, serums, fortifiers)	22	NA	NA	NR	NA					
Eyelash Cleansers										
Other Eye Makeup Preparations	55	5	4	10	4-11	NR	NR	1	NR	NR
Children's Eye Makeup Preparations	20									
Children's Eyeshadows	19	NA	NA	NR	NA					
Other Children's Eye Makeup	1	NA	NA	NR	NA					
Fragrance Preparations	44					4				
Cologne and Toilet Water										
Perfumes	10	NR	1	NR	NR	1	NR	NR	NR	NR
Other Fragrance Preparation	34	NR	NR	NR	NR	3	NR	4	NR	NR
Hair Preparations (non-coloring)	80					6				
Hair Conditioners	6 (l.o.); 3 (r.o.)	NR	NR	NR	NR	2 (r.o.)	NR	14	NR	0.8-2
Hair Sprays (aerosol fixatives)	6	NR	NR	NR	NR					

Table 1. Frequency (RLD/VCRP) and concentration of use according to likely duration and exposure and by product category

	<i># of Uses</i>			<i>Max Conc of Use</i>		<i># of Uses</i>			<i>Max Conc of Use</i>	
	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵
Hair Straighteners	3	NR	NR	NR	NR	1	NR	10	NR	21
Permanent Waves						NR	NR	1	NR	NR
Rinses (non-coloring)	2	NR	NR	NR	NR	NR	NR	1	NR	NR
Shampoos (non-coloring)	1	NR	NR	NR	NR					
Tonics, Dressings, and Other Hair Grooming Aids	12	11	4	8.2	2	5	NR	NR	NR	9
Wave Sets	NR	2	NR	NR	NR					
Other Hair Preparations	54 (l.o.); 2 (r.o.)	3	1	NR	17					
<i>Hair Coloring Preparations</i>	36					8				
Hair Dyes and Colors (all types requiring caution statements and patch tests)	6	NR	NR	NR	NR	7	NR	NR	NR	NR
Hair Tints	16	NR	NR	NR	NR					
Hair Rinses (coloring)	1 (r.o.)	NR	NR	NR	NR					
Hair Color Sprays (aerosol)										
Hair Lighteners with Color	3	NR	NR	NR	NR					
Hair Bleaches	1	NR	NR	NR	NR	NR	NR	1	NR	NR
Eyeshadow and Eyebrow Dyes	1	NA	NA	NR	NA					
Other Hair Coloring Preparation	5 (l.o.); 5 (r.o.)	NR	1	NR	NR	1 (r.o.)	NR	6	NR	NR
<i>Makeup Preparations (not eye; not children's)</i>	3352									
Blushers and Rouges (all types)	271	1	13	5	6					
Face Powders	52	NR	NR	NR	6					
Foundations	208 (traditional application)	NR	6	NR	4-7	NR	NR	2	NR	NR
Leg and Body Paints	NR	1	NR	NR	NR					
Lipstick and Lip Glosses	2398	61	191	6.6	0.8-20					
Makeup Bases	176 (traditional application)	NR	NR	NR	NR					
Makeup Fixatives	49	NR	NR	NR	3					
Other Makeup Preparations	249 (l.o.); 1 (r.o.)	22	15	NR	4-13					
<i>Makeup Preparations for Children (not eye)</i>	748									
Children's Blushers and Rouges (All Types)	15	NA	NA	NR	NA					
Children's Face Paints	6	NA	NA	NR	NA					
Children's Face Powders	2	NA	NA	NR	NA					
Children's Foundations										
Children's Lipsticks and Lip Glosses	718	NA	NA	NR	NA					
Other Children's Makeup	10	NA	NA		NA					
<i>Manicuring Preparations</i>	80					3				
Basecoats and Undercoats										
Cuticle Softeners	1	NR	NR	NR	NR	NR	NR	2	NR	0.5
Nail Creams and Lotions	1	NR	2	NR	NR	2	NR	1	NR	NR
Nail Extenders	9	NR	NR	NR	NR					
Nail Polish and Enamel	20	NR	1	NR	4					
Nail Polish and Enamel Removers										
Other Manicuring Preparations	53	NR	NR	NR	NR	1	NR	NR	NR	NR
<i>Oral Products</i>	2									
Dentifrices										
Other Oral Products	2	NR	NR	NR	NR					

Table 1. Frequency (RLD/VCRP) and concentration of use according to likely duration and exposure and by product category

	<i># of Uses</i>			<i>Max Conc of Use</i>		<i># of Uses</i>			<i>Max Conc of Use</i>	
	<i>RLD (2025)¹</i>	<i>VCRP (2023)²</i>	<i>VCRP (2002)³</i>	<i>% (2022)⁴</i>	<i>% (2003)⁵</i>	<i>RLD (2025)¹</i>	<i>VCRP (2023)²</i>	<i>VCRP (2002)³</i>	<i>% (2022)⁴</i>	<i>% (2003)⁵</i>
<i>Personal Cleanliness</i>	35					3				
Bath Soaps and Body Washes	5	NR	NR	NR	NR	1	NR	NR	NR	NR
Deodorants (underarm)	28	NR	NR	NR	NR					
Douches										
Feminine Deodorants										
Other Personal Cleanliness Products	1 (l.o.); 1 (r.o.)	1	NR	NR	NR	2 (r.o.)	1	NR	NR	NR
<i>Shaving Preparations</i>	3									
Aftershave Lotions	1	NR	NR	NR	NR					
Beard Softeners										
Shaving Creams (aerosol, brushless, lather)	3	NR	NR	NR	NR	NR	NR	NR	1.5	NR
Shaving Soaps (cakes, sticks, etc.)										
Other Shaving Preparation Products										
<i>Skin Care Preparations</i>	362					86				
Cleansing	29	2	13	NR	NR	7	NR	2	NR	1
Depilatories	16	NR	NR	NR	NR					
Face and Neck (excluding shaving preparations)	119 (l.o.); 16 (r.o.)	4	1	NR	5-8	5 (l.o.); 3 (r.o.)	NR	NR	NR	1
Body and Hand (excluding shaving preparations)	24 (l.o.); 2 (r.o.)	NR	5	NR	8	23 (l.o.); 3 (r.o.)	NR	12	NR	2-10
Foot Powders and Sprays										
Moisturizing	158	10	11	2 (not spray)	5	55	4	25	NR	12
Night	2	3	4	NR	5	3	NR	4	NR	1
Paste Masks (mud packs)	2	NR	1	NR	NR	2	NR	1	NR	1
Skin Fresheners	10	NR	NR	NR	NR	2	NR	NR	NR	NR
Other Skin Care Preparations	42 (l.o.); 8 (r.o.)	7	9	NR	NR	4 (l.o.); 5 (r.o.)	NR	5	NR	NR
<i>Suntan Preparations</i>	2									
Suntan Gels, Creams, and Liquids	1	NR	2	NR	13					
Indoor Tanning Preparations	1	NR	NR	NR	NR	NR	NR	5	NR	NR
Other Suntan Preparations						NR	NR	1	NR	NR
<i>Tattoo Preparations</i>	1									
Permanent Tattoo Inks	1	NA	NA	NR	NA					
Temporary Tattoo Inks	1	NA	NA	NR	NA					
Other Tattoo Preparations										
<i>Other Preparations (i.e., those preparations that do not fit another category)</i>	33	NA	NA	NR	NA					

Table 1. Frequency (RLD/VCRP) and concentration of use according to likely duration and exposure and by product category

	# of Uses			Max Conc of Use		# of Uses			Max Conc of Use	
	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵
	Microcrystalline Wax					Montan Wax				
Totals*	10,285	1130	581	0.092-34.5	0.2-50	34	3	13	0.54	0.5-11
summarized by likely duration and exposure**										
Duration of Use										
Leave-On	***	1030	564	0.092-30	0.2-50	***	3	12	0.54	0.5-11
Rinse-Off	***	100	16	0.75-34.5	0.4-10	***	NR	1	NR	NR
Diluted for (Bath) Use	***	NR	1	NR	0.5	***	NR	NR	NR	NR
Exposure Type										
Eye Area	***	235	211	2.8-24.3	0.3-19	***	2	NR	NR	0.5-9
Incidental Ingestion	***	466	229	5.9-34.5	0.5-10	***	1	7	0.54	0.5
Incidental Inhalation-Spray	***	1; 84 ^a ; 22 ^c	2; 24 ^a ; 7 ^c	0.2-1; 0.32-30 ^a	0.2-15 ^a	***	NR	NR	NR	NR
Incidental Inhalation-Powder	***	20; 22 ^c	1; 7 ^c	5.4; 0.2-2.2 ^b	NR	***	NR	NR	NR	NR
Dermal Contact	***	569	310	0.1-24.3	0.2-19	***	NR	5	NR	0.5-11
Deodorant (underarm)	***	NR	NR	7.7	NR	***	NR	NR	NR	NR
Hair - Non-Coloring	***	36	20	0.32-30	0.4-50	***	NR	NR	NR	NR
Hair-Coloring	***	NR	1	1-2	NR	***	NR	1	NR	NR
Nail	***	23	1	0.092	NR	***	NR	NR	NR	0.5
Mucous Membrane	***	494	231	5.9-34.5	0.5-10	***	1	7	0.54	0.5
Baby Products	***	NR	NR	NR	3	***	NR	NR	NR	NR
as reported by product category										
Baby Products	5									
Baby Lotions/Oils/Powders/Creams	3	NR	NR	NR	NR					
Other Baby Products	1 (l.o.); 1 (r.o.)	NR	NR	NR	3					
Bath Preparations (diluted for use)	5									
Bath Oils, Tablets, and Salts	NR	NR	1	NR	NR					
Other Bath Preparations	5	NR	NR	NR	0.5					
Eye Makeup Preparations	2299					9				
Eyebrow Pencil	403	24	15	3-24.3	6	4	NR	NR	NR	NR
Eyeliner	675	56	126	2.8-19.1	2-18	2	NR	NR	NR	2
Eye Shadow	812	90	16	3.2-14.7	2-6	NR	NR	NR	NR	9
Eye Lotion	27	3	NR	NR	0.3					
Eye Makeup Remover	1	NR	1	NR	NR					
False Eyelashes	1	NA	NA	NR	NA					
Mascara	284	36	20	4.6-11	3-6	3	2	NR	NR	3
Eyelash and Eyebrow Adhesives, Glues, and Sealants	24	NA	NA	NR	NA					
Eyelash and Eyebrow Preparations (primers, conditioners, serums, fortifiers)	44	NA	NA	NR	NA					
Eyelash Cleansers	3	NA	NA	NR	NA					
Other Eye Makeup Preparations	181	26	33	NR	8-19	NR	NR	NR	NR	0.5
Children's Eye Makeup Preparations	26									
Children's Eyeshadows	26	NA	NA	NR	NA					
Other Children's Eye Makeup	1	NA	NA	NR	NA					
Fragrance Preparations	84									
Cologne and Toilet Water										
Perfumes	19	1	1	0.2	NR					
Other Fragrance Preparation	66	NR	1	1	NR					
Hair Preparations (non-coloring)	146					2				
Hair Conditioners	4 (l.o.); 3 (r.o.)	NR	3	2	4					
Hair Sprays (aerosol fixatives)	13	NR	NR	NR	NR					
Hair Straighteners	4	NR	1	NR	0.4					

Table 1. Frequency (RLD/VCRP) and concentration of use according to likely duration and exposure and by product category

	<i># of Uses</i>			<i>Max Conc of Use</i>		<i># of Uses</i>			<i>Max Conc of Use</i>	
	<i>RLD (2025)¹</i>	<i>VCRP (2023)²</i>	<i>VCRP (2002)³</i>	<i>% (2022)⁴</i>	<i>% (2003)⁵</i>	<i>RLD (2025)¹</i>	<i>VCRP (2023)²</i>	<i>VCRP (2002)³</i>	<i>% (2022)⁴</i>	<i>% (2003)⁵</i>
Permanent Waves	4	NR	NR	NR	NR					
Rinses (non-coloring)	1	NR	NR	NR	NR					
Shampoos (non-coloring)	2 (r.o.)	NR	NR	NR	NR					
Tonics, Dressings, and Other Hair Grooming Aids	33	25	15	0.32-30	0.8					
Wave Sets										
Other Hair Preparations	82 (l.o.); 7 (r.o.)	11	1	NR	3-50	1 (l.o.); 1 (r.o.)	NR	NR	NR	NR
<i>Hair Coloring Preparations</i>	70									
Hair Dyes and Colors (all types requiring caution statements and patch tests)	27	NR	NR	1	NR					
Hair Tints	29	NR	NR	NR	NR					
Hair Rinses (coloring)	1 (r.o.)	NR	NR	2	NR					
Hair Color Sprays (aerosol)										
Hair Lighteners with Color	2	NR	NR	NR	NR					
Hair Bleaches										
Eyelash and Eyebrow Dyes	1	NA	NA	NR	NA					
Other Hair Coloring Preparation	11 (l.o.); 2 (r.o.)	NR	1	NR	NR	NR	NR	1	NR	NR
<i>Makeup Preparations (not eye; not children's)</i>	6289					22				
Blushers and Rouges (all types)	721	41	8	2.2-13	6-10					
Face Powders	246	20	1	5.4	NR					
Foundations	314 (traditional application); 7 (airbrush application)	29	35	5.6-15	9-12	NR	NR	5	NR	11
Leg and Body Paints	123 (traditional application)	2	NR	NR	NR					
Lipstick and Lip Glosses	4440	466	229	5.9-17.6	0.5-10	3	1	7	0.54	0.5
Makeup Bases	137 (traditional application); 1 (airbrush application)	8	2	0.1	NR					
Makeup Fixatives	32	1	1	NR	NR					
Other Makeup Preparations	525 (l.o.); 3 (r.o.)	87	33	16.8	3-13	19 (l.o.)	NR	NR	NR	0.5
<i>Makeup Preparations for Children (not eye)</i>	206					1				
Children's Blushers and Rouges (All Types)	10	NA	NA	NR	NA		NA	NA		NA
Children's Face Paints	7	NA	NA	NR	NA	1	NA	NA	NR	NA
Children's Face Powders	2	NA	NA	NR	NA					
Children's Foundations	1	NA	NA	NR	NA					
Children's Lipsticks and Lip Glosses	167	NA	NA	NR	NA					
Other Children's Makeup	26	NA	NA	NR	NA					
<i>Manicuring Preparations (Nail)</i>	410					1				
Basecoats and Undercoats	24	1	NR	NR	NR					
Cuticle Softeners	12	NR	NR	NR	NR					
Nail Creams and Lotions	24	1	NR	NR	NR					
Nail Extenders	231	1	NR	NR	NR					
Nail Polish and Enamel	329	15	1	0.092	NR	1	NR	NR	NR	0.5
Nail Polish and Enamel Removers	26	NR	NR	NR	NR					
Other Manicuring Preparations	42	5	NR	NR	NR					

Table 1. Frequency (RLD/VCRP) and concentration of use according to likely duration and exposure and by product category

	<i># of Uses</i>			<i>Max Conc of Use</i>		<i># of Uses</i>			<i>Max Conc of Use</i>	
	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵
<i>Oral Hygiene Products</i>	3									
Dentifrices										
Other Oral Products	3	NR	NR	34.5	10					
<i>Personal Cleanliness Products</i>	36									
Bath Soaps and Body Washes	23	22	NR	NR	NR					
Deodorants (underarm)	2	NR	NR	7.7 (not spray)	NR					
Douches										
Feminine Deodorants										
Other Personal Cleanliness Products	2 (l.o.); 9 (r.o.)	6	1	NR	NR					
<i>Shaving Preparations</i>	13									
Aftershave Lotions										
Beard Softeners	2	NR	NR	NR	NR					
Shaving Creams (aerosol, brushless, lather)	3	NR	NR	NR	NR					
Shaving Soaps (cakes, sticks, etc.)	NR	NR	NR	10	NR					
Other Shaving Preparation Products	8	NR	NR	NR	0.5					
<i>Skin Care Preparations</i>	782									
Cleansing	69	58	6	0.75-10	2					
Depilatories	50	13	1	8	4					
Face and Neck (excluding shaving preparations)	186 (l.o.); 30 (r.o.)	20	2	0.85-2.2 (not spray)	NR					
Body and Hand (excluding shaving preparations)	84 (l.o.); 2 (r.o.)	2	5	0.2-0.5 (not spray)	NR					
Foot Powders and Sprays	3	NR	NR	NR	NR					
Moisturizing	335	52	5	0.37 (not spray)	NR					
Night	4	5	4	1	2					
Paste Masks (mud packs)	22	1	2	1	NR					
Skin Fresheners	30	NR	NR	NR	NR					
Other Skin Care Preparations	87 (l.o.); 30 (r.o.)	NR	10	NR	1					
<i>Suntan Preparations</i>	9									
Suntan Gels, Creams, and Liquids	5	2	NR	NR	1-15					
Indoor Tanning Preparations	1 (traditional application)	NR	NR	NR	15					
Other Suntan Preparations	5	NR	NR	NR	0.2-0.5					
<i>Tattoo Preparations</i>	11									
Permanent Tattoo Inks										
Temporary Tattoo Inks	1	NA	NA	NR	NA					
Other Tattoo Preparations	10	NA	NA	NR	NA					
<i>Other Preparations (i.e., those preparations that do not fit another category)</i>	153	NA	NA	NR	NA					

Table 1. Frequency (RLD/VCRP) and concentration of use according to likely duration and exposure and by product category

	# of Uses			Max Conc of Use		# of Uses			Max Conc of Use	
	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵
	Ozokerite					Paraffin				
Totals*	8498	371	680	0.1-23.3	0.08-22	7243	737	481	0.01-21	0.03-99
summarized by likely duration and exposure**										
Duration of Use										
Leave-On	***	367	621	23.3	1-22	***	543	400	0.01-18.6	99
Rinse-Off	***	4	59	0.1	0.08-22	***	194	81	0.45-21	32
Diluted for (Bath) Use	***	NR	NR	0	15	***	NR	NR	NR	NR
Exposure Type										
Eye Area	***	71	83	0.98-15.2	2-22	***	227	52	0.19-18.6	1-20
Incidental Ingestion	***	138	374	5.8-12	2-21	***	118	134	9-18.2	2-10
Incidental Inhalation-Spray	***	2; 23 ^a ; 4 ^c	13; 37 ^a	2-23.3 ^a	15; 1-15 ^a ; 7 ^c	***	68 ^a ; 12 ^c	58 ^a ; 24 ^c	7; 2.2-13.5 ^a	0.3-20 ^a ; 3 ^c
Incidental Inhalation-Powder	***	13; 4 ^b	1; 1 ^b ; 12 ^c	2; 6 ^b	1; 2 ^b	***	12; 1 ^b ; 12 ^c	24 ^c	0.5-15 ^b	1 ^b ; 3 ^c
Dermal Contact	***	201	228	0.1-23.3	0.08-22	***	375	264	0.091-21	0.3-99
Deodorant (underarm)	***	59 ^a	NR	2.8	NR	***	NR	2 ^a	11-11.1	NR
Hair - Non-Coloring	***	17	6	2-6.5	NR	***	39	33	1.5-13.5	2-20
Hair-Coloring	***	NR	46	NR	0.3	***	86	13	0.45-5	NR
Nail	***	NR	1	NR	3	***	6	9	0.01-3.2	0.03-2
Mucous Membrane	***	138	375	5.8-12	0.08-21	***	126	157	9-21	2-10
Baby Products	***	NR	1	NR	2	***	1	NR	NR	1
as reported by product category										
Baby Products										
Baby Lotions/Oils/Powders/Creams	NR	NR	1	NR	2	5	1	NR	NR	1
Other Baby Products						3 (l.o.); 1 (r.o.)	NR	NR	NR	NR
Bath Preparations										
Bath Oils, Tablets, and Salts	3					4				
Other Bath Preparations	3	NR	NR	NR	15	2	NR	NR	NR	NR
Eye Makeup Preparations (not children's)										
	2084					1790				
Eyebrow Pencil	336	8	8	8	NR	171	13	4	17.9	6
Eyeliners	827	23	11	0.98-15.2	7-10	446	32	7	18.6	4-20
Eye Shadow	554	20	10	7.1	6-14	393	47	5	0.19-15.3	3-8
Eye Lotion	12	2	NR	4	NR	9	1	1	NR	1-6
Eye Makeup Remover	1	NR	3	NR	2-4	1	NR	3	NR	NR
False Eyelashes						3	NA	NA	NR	NA
Mascara	337	15	25	NR	7-9	700	113	28	9-17	6-20
Eyelash and Eyebrow Adhesives, Glues, and Sealants	17	NA	NA	NR	NA	12	NA	NA	NR	NA
Eyelash and Eyebrow Preparations (primers, conditioners, serums, fortifiers)	30	NA	NA	NR	NA	39	NA	NA	NR	NA
Eyelash Cleansers							NA	NA		NA
Other Eye Makeup Preparations	127	3	26	NR	2-22	111	21	4	6	4
Children's Eye Makeup Preparations										
Children's Eyeshadows	1					5				
Other Children's Eye Makeup	1	NA	NA	NR	NA	4	NA	NA	NR	NA
Fragrance Preparations										
	170					32				
Cologne and Toilet Water	NR	NR	1	NR	15					
Perfumes	50	1	10	NR	15	5	NR	NR	NR	NR
Other Fragrance Preparation	121	1	2	NR	15	28	NR	1	7	NR
Hair Preparations (non-coloring)										
	164					204				
Hair Conditioners	14	NR	1	NR	NR	16 (l.o.); 15 (r.o.)	5	7	1.5-2	2

Table 1. Frequency (RLD/VCRP) and concentration of use according to likely duration and exposure and by product category

	# of Uses			Max Conc of Use		# of Uses			Max Conc of Use	
	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵
Hair Sprays (aerosol fixatives)	8	NR	NR	NR	NR	11	NR	NR	NR	NR
Hair Straighteners	2	NR	NR	NR	NR	12	5	NR	NR	NR
Permanent Waves	1	NR	NR	NR	NR	7	NR	NR	NR	NR
Rinses (non-coloring)						25	NR	NR	NR	NR
Shampoos (non-coloring)						6 (r.o.)	NR	1	NR	NR
Tonics, Dressings, and Other Hair Grooming Aids	61	14	3	2-6.5	NR	48	21	23	2.2-13.5	2-20
Wave Sets	2	NR	NR	NR	NR	NR	NR	1	NR	NR
Other Hair Preparations	81 (l.o.); 5 (r.o.)	3	2	NR	NR	72 (l.o.); 14 (r.o.)	8	1	NR	2
Hair Coloring Preparations	58					885				
Hair Dyes and Colors (all types requiring caution statements and patch tests)	33	NR	46	NR	NR	787	82	NR	5	NR
Hair Tints	17	NR	NR	NR	NR	209	1	1	NR	NR
Hair Rinses (coloring)	1	NR	NR	NR	NR	1 (l.o.); 3 (r.o.)	NR	NR	NR	NR
Hair Color Sprays (aerosol)	2	NR	NR	NR	NR	NR	NR	2	NR	NR
Hair Lighteners with Color										
Hair Bleaches	1	NR	NR	NR	NR	4	1	NR	0.45	NR
Eyelash and Eyebrow Dyes	1	NA	NA	NR	NA					
Other Hair Coloring Preparation	5 (l.o.); 2 (r.o.)	NR	NR	NR	0.3	17 (l.o.); 11 (r.o.)	2	10	NR	NR
Makeup Preparations (not eye; not children's)	5333					3442				
Blushers and Rouges (all types)	307	1	10	NR	2-21	212	17	4	6	NR
Face Powders	168	13	1	2	1	75	12	NR	0.091	NR
Foundations	655	11	33	NR	4-15	404	20	33	0.5-6	2-6
	(traditional application); 3 (airbrush application)					(traditional application)				
Leg and Body Paints	68 (traditional application)	NR	NR	NR	8	103 (traditional application)	2	NR	NR	NR
Lipstick and Lip Glosses	3765	138	374	5.8-12	2-21	2330	118	134	9-18.2	2-10
Makeup Bases	116	3	11	NR	NR	162	4	37	NR	NR
	(traditional application); 1 (airbrush application)									
Makeup Fixatives	52	NR	NR	NR	NR	7	2	1	NR	NR
Other Makeup Preparations	398 (l.o.)	33	37	6.5	3-21	223 (l.o.)	29	18	0.5	2-21
Makeup Preparations for Children (not eye)	156					65				
Children's Blushers and Rouges (All Types)	3	NA	NA	NR	NA	1	NA	NA	NR	NA
Children's Face Paints	1	NA	NA	NR	NA	5	NA	NA	NR	NA
Children's Face Powders										
Children's Foundations										
Children's Lipsticks and Lip Glosses	152	NA	NA	NR	NA	45	NA	NA	NR	NA
Other Children's Makeup	1	NA	NA	NR	NA	17	NA	NA	NR	NA
Manicuring Preparations	9					49				
Basecoats and Undercoats	1	NR	NR	NR	NR	NR	1	NR	0.01	NR
Cuticle Softeners	3	NR	1	NR	NR	1	NR	1	NR	NR
Nail Creams and Lotions						1	1	3	NR	NR
Nail Extenders						1	NR	NR	NR	NR

Table 1. Frequency (RLD/VCRP) and concentration of use according to likely duration and exposure and by product category

	<i># of Uses</i>			<i>Max Conc of Use</i>		<i># of Uses</i>			<i>Max Conc of Use</i>	
	<i>RLD (2025)¹</i>	<i>VCRP (2023)²</i>	<i>VCRP (2002)³</i>	<i>% (2022)⁴</i>	<i>% (2003)⁵</i>	<i>RLD (2025)¹</i>	<i>VCRP (2023)²</i>	<i>VCRP (2002)³</i>	<i>% (2022)⁴</i>	<i>% (2003)⁵</i>
Nail Polish and Enamel	1	NR	NR	NR	3	30	2	5	NR	0.03-2
Nail Polish and Enamel Removers						3	NR	NR	NR	NR
Other Manicuring Preparations	5	NR	NR	NR	NR	14	2	NR	3.2	NR
Oral Products	2									
Dentifrices	1	NR	NR	NR	NR					
Other Oral Products	1	NR	NR	NR	NR	NR	NR	NR	11.2	NR
Personal Cleanliness	154					24				
Bath Soaps and Body Washes	1	NR	NR	NR	0.08	7	5	7	21	NR
Deodorants (underarm)	147	59	NR	2.8 (not spray)	NR	3	NR	2	11-11.1 (not spray)	NR
Douches						5 (l.o.); 9 (r.o.)	NR	NR	NR	NR
Feminine Deodorants	13 (l.o.)	NR	NR	NR	NR					
Other Personal Cleanliness Products	1 (l.o.); 4 (r.o.)	NR	1	NR	NR	NR	3	16	NR	NR
Shaving Preparations	4					13				
Aftershave Lotions	1	NR	NR	NR	NR	1	NR	NR	NR	NR
Beard Softeners						1	2	NR	NR	NR
Shaving Creams (aerosol, brushless, lather)	2	NR	NR	NR	NR	2	NR	1	NR	NR
Shaving Soaps (cakes, sticks, etc.)										
Other Shaving Preparation Products	1	NR	NR	NR	NR	9	2	NR	NR	NR
Skin Care Preparations	421					696				
Cleansing	18	2	4	0.1	1-5	51	12	26	NR	12-20
Depilatories	6	NR	NR	NR	NR	194	75	1	NR	NR
Face and Neck (excluding shaving preparations)	62 (l.o.); 15 (r.o.)	4	2	6 (not spray)	NR	99 (l.o.); 14 (r.o.)	11	4	1-8 (not spray)	NR
Body and Hand (excluding shaving preparations)	34 (l.o.)	NR	10	NR	7	121 (l.o.); 7 (r.o.)	1	19	0.5-15 (not spray)	3
Foot Powders and Sprays	13	NR	NR	NR	NR	1	NR	1	NR	NR
Moisturizing	256	9	15	0.2-3 (not spray)	1-5	208	40	24	0.8-5 (not spray)	2
Night	3	NR	7	NR	2	5	4	6	NR	0.3-8
Paste Masks (mud packs)	5	2	4	NR	NR	6	1	7	NR	32
Skin Fresheners	5	NR	NR	NR	NR	18	1	NR	NR	NR
Other Skin Care Preparations	50 (l.o.); 8 (r.o.)	6	7	NR	6	80 (l.o.); 19 (r.o.)	15	27	15	99
Suntan Preparations	15					12				
Suntan Gels, Creams, and Liquids	12	NR	9	23.3	8-15	10	2	3	NR	2-10
Indoor Tanning Preparations										
Other Suntan Preparations	3	NR	3	NR	3-12	2	NR	2	NR	20
Tattoo Preparations	9					5				
Permanent Tattoo Inks										
Temporary Tattoo Inks	2	NA	NA	NR	NA	1	NA	NA	NR	NA
Other Tattoo Preparations	7	NA	NA	NR	NA	4	NA	NA	NR	NA
Other Preparations (i.e., those preparations that do not fit another category)	62	NA	NA	NR	NA	153	NA	NA	NR	NA

Table 1. Frequency (RLD/VCRP) and concentration of use according to likely duration and exposure and by product category

	# of Uses			Max Conc of Use		# of Uses			Max Conc of Use	
	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵
	Synthetic Beeswax					Synthetic Wax				
Totals*	5602	164	179	0.0012-24.9	0.05-18	15,166	1050	215	0.05-36	0.1-29
summarized by likely duration and exposure**										
Duration of Use										
Leave-On	***	154	178	0.0012-24.9	0.05-18	***	1014	214	0.05-36	0.1-29
Rinse-Off	***	10	1	5	NR	***	35	1	0.3-11.2	5
Diluted for (Bath) Use	***	NR	NR	NR	NR	***	1	NR	NR	NR
Exposure Type										
Eye Area	***	98	23	1-24.9	0.05-18	***	234	50	0.75-36	0.8-27
Incidental Ingestion	***	23	131	2-23.3	4-7	***	536	134	2.6-16.8	3-29
Incidental Inhalation-Spray	***	3; 10 ^a ; 5 ^c	1; 5 ^a ; 3 ^c	20; 5.3 ^a	0.5-12 ^a ; 1-3 ^c	***	5; 23 ^a ; 7 ^c	3; 1 ^a	0.3; 0.05-5 ^a	1-3 ^a
Incidental Inhalation-Powder	***	1 ^b ; 5 ^c	1 ^b ; 3 ^c	0.0012-1.5 ^b	0.5 ^b ; 1-3 ^c	***	15; 7 ^c	2	5	0.5; 4 ^b ; 2-4 ^c
Dermal Contact	***	86	42	0.0012-24.9	0.06-14	***	493	70	0.1-36	0.5-27
Deodorant (underarm)	***	NR	NR	NR	NR	***	12 ^a	NR	3.4-12	NR
Hair - Non-Coloring	***	5	NR	5.3	12	***	5	1	0.05-5	NR
Hair-Coloring	***	NR	NR	NR	NR	***	NR	NR	NR	NR
Nail	***	NR	NR	NR	NR	***	NR	NR	NR	0.1
Mucous Membrane	***	23	131	2-23.3	4-7	***	548	134	2.6-16.8	3-29
Baby Products	***	1	2	NR	0.5	***	NR	NR	NR	4
as reported by product category										
Baby Products	5					4				
Baby Lotions/Oils/Powders/Creams	4	1	1	NR	0.5	2	NR	NR	NR	4
Other Baby Products	1 (l.o.)	NR	1	NR	NR	1 (l.o.); 1 (r.o.)	NR	NR	NR	NR
Bath Preparations (diluted for use)	1					4				
Bath Oils, Tablets, and Salts	1	NR	NR	NR	NR					
Other Bath Preparations						4	1	NR	NR	NR
Eye Makeup Preparations	1697					4423				
Eyebrow Pencil	144	11	6	NR	NR	682	28	2	5-36	NR
Eyeliner	470	18	9	2-24.9	0.06-14	1731	104	28	0.75-14.1	0.8-27
Eye Shadow	475	5	1	1-13.6	NR	1835	68	9	4.1-9.6	NR
Eye Lotion	16	2	NR	NR	NR	11	1	NR	NR	NR
Eye Makeup Remover	1	NR	NR	NR	NR	1	NR	NR	NR	NR
False Eyelashes	1	NA	NA	NR	NA	5	NA	NA	NR	NA
Mascara	519	50	6	8-9.3	0.05-18	226	16	10	1.5-8.7	2-11
Eyelash and Eyebrow Adhesives, Glues, and Sealants	20	NA	NA	NR	NA	19	NA	NA	NR	NA
Eyelash and Eyebrow Preparations (primers, conditioners, serums, fortifiers)	77	NA	NA	NR	NA	60	NA	NA	NR	NA
Eyelash Cleansers		NA	NA		NA		NA	NA		NA
Other Eye Makeup Preparations	86	12	1	14.7	0.06-9	256	17	1	13	NR
Children's Eye Makeup Preparations	4					2				
Children's Eyeshadows	3	NA	NA	NR	NA	1	NA	NA	NR	NA
Other Children's Eye Makeup	1	NA	NA	NR	NA	1	NA	NA	NR	NA
Fragrance Preparations	28					58				
Cologne and Toilet Water										
Perfumes	12	NR	NR	20	NR	13	1	1	0.3	NR
Other Fragrance Preparation	16	2	1	NR	NR	46	4	2		NR
Hair Preparations (non-coloring)	59					47				
Hair Conditioners	7 (l.o.); 2 (r.o.)	1	NR	NR	NR	3 (l.o.)	NR	1	0.3	NR
Hair Sprays (aerosol fixatives)	NR	1	NR	NR	NR	1	NR	NR	NR	NR
Hair Straighteners	1	NR	NR	NR	NR	1	NR	NR	NR	NR

Table 1. Frequency (RLD/VCRP) and concentration of use according to likely duration and exposure and by product category

	# of Uses			Max Conc of Use		# of Uses			Max Conc of Use	
	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵	RLD (2025) ¹	VCRP (2023) ²	VCRP (2002) ³	% (2022) ⁴	% (2003) ⁵
Other Oral Products	2	NR	NR	NR	NR	1	NR	NR	11.2	NR
Personal Cleanliness Products	15					58				
Bath Soaps and Body Washes						21	9	NR	7.9	NR
Deodorants (underarm)	8	NR	NR	NR	NR	6	12	NR	3.4-12 (not spray)	NR
Douches										
Feminine Deodorants										
Other Personal Cleanliness Products	3 (l.o.); 6 (r.o.)	NR	NR	NR	NR	31 (r.o.)	2	NR	NR	NR
Shaving Preparations	7					6				
Aftershave Lotions						NR	NR	NR	NR	1
Beard Softeners	1	NR	NR	NR	NR					
Shaving Creams (aerosol, brushless, lather)	2	NR	NR	NR	NR					
Shaving Soaps (cakes, sticks, etc.)										
Other Shaving Preparation Products	4	NR	NR	NR	NR	6	1	NR	NR	NR
Skin Care Preparations	596					558				
Cleansing	35	2	1	5	NR	118	20	NR	4-7.6	5
Depilatories	13	4	NR	NR	NR	15	2	NR	NR	NR
Face and Neck (excluding shaving preparations)	163 (l.o.); 21 (r.o.)	5	1	0.0012-0.5 (not spray)	NR	153 (l.o.); 36 (r.o.)	4	NR	1.2 (not spray)	3
Body and Hand (excluding shaving preparations)	132 (l.o.); 6 (r.o.)	NR	2	1.5 (not spray)	1-3	13 (l.o.); 5 (r.o.)	3	NR	0.1-0.3 (not spray)	2-4
Foot Powders and Sprays	1	NR	NR	NR	NR	2	NR	NR	NR	NR
Moisturizing	227	5	5	NR	0.5	165	17	1	5-5.9 (not spray)	NR
Night	36	3	NR	NR	NR	7	2	NR	NR	3
Paste Masks (mud packs)	6	3	NR	NR	NR	4	1	NR	NR	NR
Skin Fresheners	7	NR	NR	NR	NR	4	NR	NR	NR	NR
Other Skin Care Preparations	92 (l.o.); 7 (r.o.)	3	1	NR	NR	98 (l.o.); 30 (r.o.)	20	1	2-10	NR
Suntan Preparations	8					15				
Suntan Gels, Creams, and Liquids	6	NR	NR	NR	NR	10	NR	NR	1 (not spray)	NR
Indoor Tanning Preparations						1 (traditional application)	NR	NR	NR	NR
Other Suntan Preparations	2	NR	NR	NR	NR	4	NR	NR	NR	1
Tattoo Preparations	2									
Permanent Tattoo Inks										
Temporary Tattoo Inks										
Other Tattoo Preparations	2	NA	NA	NR	NA					
Other Preparations (i.e., those preparations that do not fit another category)	62	NA	NA	NR	NA	51	NA	NA	NR	NA

NR – not reported; NA – not applicable (this category was not part of the VCRP)

l.o. – leave-on; r.o. – rinse-off

*The total FOU provided for RLD refers to the ingredient count supplied by FDA, and is not a summation of the number of uses per category because each product may be categorized under multiple product categories. For data supplied via the VCRP or by the Council survey, the sum of all exposure types may not equal the sum of total uses because each ingredient may be used in cosmetics with multiple exposure types.

**Likely duration and exposure are derived from VCRP and survey data based on product category (see Use Categorization <https://www.cir-safety.org/cir-findings>)

*** In the RLD each ingredient may be reported under several product categories, making a summation of RLD misleading in comparison to VCRP data. Accordingly, RLD are presented below by product category (as supplied by FDA), but are not summarized by likely duration and exposure.

****at the time of the YEAR safety assessment, concentration of use data were not reported by the FDA, and a concentration of use survey was not conducted; YYYY data were presented in the original assessment and are reported here [if applicable]

^a It is possible these products are sprays, but it is not specified whether the reported uses are sprays.

^b It is possible these products are powders, but it is not specified whether the reported uses are powders.

^c Not specified whether a spray or a powder, but it is possible the use can be as a spray or a powder, therefore the information is captured in both categories

Table 2. Non-Cosmetic Use

Ingredient	Data	Reference	Notes										
Ozokerite Ceresin Montan Wax Paraffin* Microcrystalline Wax* Synthetic Wax*	Ozokerite, Ceresin, Montan Wax, Paraffin, Microcrystalline Wax, and Synthetic Wax are permitted for use in adhesives intended for use in packaging, transporting, or holding food.	21 CFR 175.105	Not included in original report.										
Montan Wax	Montan Wax is permitted for use as a defoaming agent in the manufacture of paper and paperboard intended for use in packaging, transporting, or holding food.	21 CFR 176.210	Not included in original report.										
Montan Wax Paraffin* Microcrystalline Wax* Synthetic Wax*	Montan Wax, Paraffin, Microcrystalline Wax, and Synthetic Wax are permitted for use in rubber articles intended for repeated use. It may be safely used in producing, manufacturing, packing, processing, preparing, treating, packaging, transporting, or holding food.	21 CFR 177.2600	Not included in original report.										
Paraffin* Microcrystalline Wax* Synthetic Wax*	Paraffin, Microcrystalline Wax, and Synthetic Wax must meet the following ultraviolet absorbance limits when subjected to the analytical procedure described in 21 CFR 172.886(b): <table border="1"> <thead> <tr> <th>Millimicrons</th> <th>Maximum</th> </tr> </thead> <tbody> <tr> <td>280 to 289</td> <td>0.15</td> </tr> <tr> <td>290 to 299</td> <td>.12</td> </tr> <tr> <td>300 to 359</td> <td>.08</td> </tr> <tr> <td>360 to 400</td> <td>.02</td> </tr> </tbody> </table> <ul style="list-style-type: none"> - may contain antioxidant permitted in food by regulations issued in accordance with section 409 of the act - may contain a total of not more than 1 weight percent of residues of polymers listed in 21 CFR 178.3710(d) - may contain 2-hydroxy-4-n-octoxybenzophenone as a stabilizer at a level not to exceed 0.01 weight percent of the petroleum wax -may contain poly(alkylacrylate) (CAS Reg. No. 27029-57-8) as a processing aid in the manufacture of petroleum wax. 	Millimicrons	Maximum	280 to 289	0.15	290 to 299	.12	300 to 359	.08	360 to 400	.02	21 CFR 178.3710	Not included in original report.
Millimicrons	Maximum												
280 to 289	0.15												
290 to 299	.12												
300 to 359	.08												
360 to 400	.02												
Paraffin* Microcrystalline Wax* Synthetic Wax*	Paraffin and Microcrystalline Wax are petrolatum waxes can be used the following: <ul style="list-style-type: none"> - in chewing gum base, as a masticatory substance up to an amount not to exceed good manufacturing practice - on cheese and raw fruits and vegetables as a protective coating up to an amount not to exceed good manufacturing practice -as a defoamer in food as based on 21 CFR 173.340. - as a component of microcapsules for spice-flavoring substances to exceed 50 percent by combined weight of the microcapsule and spice-flavoring substance mentioned in 21 CFR 172.230. 	21 CFR 172.886	Paraffin was mentioned to be used in fruit and vegetable protection in the original report, but the other uses in a chewing gum base, as a defoamer, and in microcapsules for spice-flavoring substances were not included for both Paraffin and Microcrystalline Wax. This information was not included at all for Synthetic Wax.										
Paraffin* Microcrystalline Wax* Synthetic Wax*	Paraffin and Microcrystalline Wax may be included the production of defoaming agents in articles intended for use in producing, manufacturing, packing, processing, preparing, treating, packaging, transporting, or holding food.	21 CFR 176.200	Not included in original report.										
Paraffin* Microcrystalline Wax* Synthetic Wax*	Paraffin and Microcrystalline Wax may be included in cellophane used for packaging food.	21 CFR 177.1200	Paraffin was mentioned to be used in food coating in the original report, but the specific use in cellophane was not included for both Paraffin and Microcrystalline Wax. This information was not included at all for Synthetic Wax.										
Paraffin* Microcrystalline Wax* Synthetic Wax*	Paraffin and Microcrystalline Wax may be included in surface lubricants employed in the manufacture of metallic articles that contact food.	21 CFR 178.3910	Not included in original report.										
Paraffin* Microcrystalline Wax* Synthetic Wax*	Paraffin and Microcrystalline Wax may be included not to exceed 1 pct by weight of the polymer in films prepared from basic polymers and with or without adjuvants for use during the irradiation of prepackaged foods.	21 CFR 179.45	Paraffin was mentioned to be used in waxed paper in the original report, but the specific use in films to use during the irradiation of prepackaged foods were not included for both Paraffin and Microcrystalline Wax.										

Table 2. Non-Cosmetic Use

Ingredient	Data	Reference	Notes
Paraffin* Microcrystalline Wax* Synthetic Wax*	Paraffin and Microcrystalline Wax may be used in used as components of the uncoated or coated food-contact surface of paper and paperboard intended for use in producing, manufacturing, packaging, processing, preparing, treating, packing, transporting, or holding aqueous and fatty foods.	21 CFR 176.170	In the original report, Paraffin was mentioned to be used in waxed paper and Microcrystalline Wax was mentioned to be used in laminating paper. However, the details of these papers being used in producing, manufacturing, packaging, processing, preparing, treating, packing, transporting, or holding aqueous and fatty foods were not included. This information was not included at all for Synthetic Wax.
Paraffin	Paraffin can be used as used in an antimicrobial pesticide formulation that can be used in the following; food-contact surfaces in public eating places, dairy-processing equipment, and food-processing equipment and utensils.	40 CFR 180.940	Not included in original report.
Paraffin	Paraffin may be used as substance in the preparation of an acrylate ester copolymer coating for a food-contact surface of an article intended for packaging and holding food, including heating of prepared food.	21 CFR 175.210	The use of this ingredient was mentioned to be used in food coating in the original report, but the specific use in acrylate ester copolymer coating was not included.
Paraffin	Paraffin may be used as a substance in the preparation of a resinous and polymeric coating for a food-contact surface of article intended for use in producing, manufacturing, packing, processing, preparing, treating, packaging, transporting, or holding food.	21 CFR 175.300	The use of this ingredient was mentioned to be used in food coating in the original report, but the specific use in a resinous and polymeric coating was not included.
Microcrystalline Wax*	Microcrystalline Wax may be used in used as substances in resinous and polymeric coatings may be safely used as the food-contact surface of articles intended for use in producing, manufacturing, packing, processing, preparing, treating, packaging, transporting, or holding food	21 CFR 175.320	Not included in original report.
Synthetic Wax	Synthetic Wax may be used in applications and under the same conditions where petroleum wax is permitted provided that the synthetic petroleum wax meets the definition and specifications prescribed in § 172.888 of this chapter.	21 CFR 172.888	Not included in original report.
Synthetic Wax	Synthetic Wax should conform to the following specifications: -there is not specification for congealing point except those components that have a congealing point below 50°C as mentioned in 21 CFR 175.250(b)(1) -oil content should not exceed 2.5% as determined by ASTM method D721-56T. -the absorptivity at 290 millimicrons in decahydronaphthalene at 88°C should not exceed 0.01 as determined by ASTM method E131-81a.	21 CFR 175.250	The original report mentions that Synthetic Wax can be referred to as Synthetic Paraffin but does not mention the specifications.
Prescription/OTC/Therapeutic Use Studies			
Paraffin	Paraffin can be used up to 5 percent in combination with one or more other emollient agents included in the monograph as an ophthalmic emollient.	21 CFR 349.14	The use of this ingredient was mentioned to be used in pharmaceutical ointments and salves in the original report, but the percentage was not mentioned.

CFR - Code of Federal Regulations; OTC - over-the-counter

*In the regulations, "petroleum wax" is mentioned but as Paraffin and Microcrystalline Wax are petroleum waxes, this applies to these ingredients as well. Synthetic Wax may also be used in the same applications that petroleum wax is mentioned.

Table 3. Acute toxicity studies

Test Article	Vehicle	Animals/Group	Dose/Concentration	Protocol	LD ₅₀ /LC ₅₀ /Results	Reference	Notes
ORAL							
Montan Wax	oil	rats (10/sex/dose)	12,000 mg/kg (6000 mg/kg applied in 2 doses within 2 h)	OECD TG 401 (acute oral toxicity); administered via gavage; applied 2x within 2 h & observed after 7 d.	LD ₅₀ > 12000 mg/kg bw; 1 female rat died directly after application while other rats had no symptoms.	1	No oral acute toxicity reports in the original report.
Paraffin	arachis oil	rats (5/sex/dose)	1000 and 5000 mg/kg	OECD TG 401 (acute oral toxicity); administered via gavage; weighed and observed after dosing at 20 min, 1, 24, 48, and 72 h, and 7 d.	LD ₅₀ > 5000 mg/kg bw; lack of system toxicity effects and mortality	2	This study was not included in the original report, but the oral acute toxicity studies showed that Paraffin was relatively nonlethal.
Synthetic Wax	none	Sprague-Dawley rats (5/sex/dose)	5000 mg/kg	OECD TG 420 (acute oral toxicity - fixed dose method); administered via gavage; clinical observations made at 0.5, 1, 2, and 4 h and 1x per day for 14 d; body weights recorded at 0, 7, and 14 d; organ weights and histopathology also examined	LD ₅₀ > 5000 mg/kg; no deaths or signs of systemic toxicity; body weight gain was normal; no abnormalities at necropsy	3	This study was not included in the original report, but the oral acute toxicity studies showed that Synthetic Wax did not produce toxic effects.
DERMAL							
Paraffin	olive oil	Sprague-Dawley rats (5/sex/dose)	2000 mg/kg bw	OECD TG 402 (acute dermal toxicity); dermally exposed for 24 h; observed for 14 d.	LD ₅₀ > 2000 mg/kg bw; scab formations on 4/5 male & 2/5 female rats (lesions remained throughout whole study but were not severe); lack of system toxicity effects and mortality, minimal clinical observations, standard body weight and weight gain, and necropsy results	2	This study was not included in the original report, but the dermal acute toxicity studies showed that Paraffin had to minimal to no irritation.

Table 4. Oral repeated dose toxicity studies

Test Article	Vehicle	Animals/Group	Study Duration	Dose/Concentration	Protocol	Results	Reference	Notes
Paraffin	NR	Fischer-344 and Sprague Dawley rats (9-10 animals/group)	60 d	2%	Animals were monitored for food uptake, clinical condition, and weight gain during the study. At 60 d, rats were killed to undergo hematology evaluation, serum liver enzyme analysis, MHC analysis, histopathological evaluation, and KC were examined for granuloma formation.	LOEL is 2% of dietary concentration of Paraffin but no NOAEL established; Fischer-344: microgranulomas, lymphoid cell aggregates in all, greater ALT, AST, & GGT activities in serum, elevated total white blood cell & neutrophil counts, changes in KC cells Sprague Dawley: microgranulomas in 1	2	No oral repeated dose toxicity studies in the original report.
Paraffin	none	Fischer 344 rats (20/sex/dose for waxes, 60/sex for control)	90 d	0.002, 0.02, 0.2, or 2.0% in diet (~1.5, 15, 150, 1500 mg)	OECD TG 408 (Repeated Dose 90-Day Oral Toxicity Study in Rodents); rats given test substance via diet; rats observed for mortality, food consumption, ophthalmoscopic examination, body weight changes, hematology, organ weights, histopathology, and clinical chemistry.	The NOAEL was determined to be 1.5 mg/kg/d; the rats had histiocytosis in the mesenteric lymph nodes, granuloma in the liver, and inflammation of cardiac mitral valve.	2	No oral repeated dose toxicity studies in the original report.

ALT – alanine aminotransferase, AST – aspartate aminotransferase, GGT – gamma-glutamyl transferase; KC - Kupffer cells; LOEL – lowest observed effect level; MHC – major histocompatibility complex; NOAEL - no observed adverse effect level

Table 5. Cytotoxicity studies

Test Article	Vehicle	Animals/Group	Dose/Concentration	Procedure	Results	Reference	Notes
Paraffin	Dressings, creams, and ointments	HaCaTs and primary human keratinocytes	Varied across different doses, concentrations, & mediums.	10 wound dressings and 7 creams and ointments were tested. Cytotoxicity was measured using toluidine blue staining (photo visualization) and cell count of trypan blue excluded cells after 24 h incubation period. Melolite was used as negative control.	White soft Paraffin (50% white soft & 50% liquid Paraffin) were found to be noncytotoxic.	4	No cytotoxicity reports in original report.

Table 6. Genotoxicity studies

Test Article	Vehicle	Concentration/Dose	Test System	Protocol	Results	Reference	Notes
IN VITRO							
Paraffin	DMSO	0.018, 0.037, 0.074, 0.15, 0.29, 0.59, 1.2, 1.7, 2.4, 3.4, 4.9, 7, 10 mmol/l.	mouse lymphoma L5178Y cells	OECD TG 476 (in vitro mammalian cell gene mutation test); positive controls MMS & MCA; negative control DMSO; 24 h @ 37°C under 5% carbon dioxide.	No evidence of Paraffin inducing mutant colonies over background.	2	No genotoxicity reports in original report.
Synthetic Wax	THF	0, 15, 50, 150, 500, 1500, and 5000 µg/plate	<i>Salmonella typhimurium</i> TA1535, TA1537, TA98 and TA100 <i>Escherichia coli</i> WP2 uvr A	OECD Guideline 471 (bacterial reverse mutation assay); Non-mutagenic. positive controls (4-nitroquinoline-N-oxide, N-ethyl-N-nitro-N-nitrosoguanidine, 9-aminoacridine, 2-Aminoanthracene, benzo(a)pyrene); incubated for 48 h & exposure duration was 48-72 h		3	No genotoxicity reports in original report.
Synthetic Wax	THF	Experiment I: 3, 10, 33, 100, 333, 1000, 2500; and 5000 µg/plate Experiment II: 10, 33, 100, 333, 1000, 2500, and 5000 µg/plate.	Experiment I: <i>S. typhimurium</i> strains TA1535 Experiment II: <i>coli</i> strain WP2 uvrA	OECD TG 471 (bacterial reverse mutation assay); negative controls untreated; positive controls (sodium azide, MMS, 4-nitro-o-phenylenediamine, 2-aminoanthracene); preincubation – 30 min; exposure – 72 h.	Non-mutagenic.	3	No genotoxicity reports in original report.
Synthetic Wax	THF	Pre-experiment: 0.8 µg/ml - 100 µg/ml Main experiment: 0.09; 0.27; 0.8; 1.6; 3.1; and 6.3 µg/ml	Chinese hamster lung fibroblasts (V79)	OECD TG 487 (in vitro mammalian cell micronucleus test); positive controls (ethylmethanesulfonate, 7,12-dimethylbenzanthracene); 4-h exposure before expression time at 7 d and selection time at 10 d.	Did not induce gene mutations at HRPT locus in V79 cells so non-mutagenic.	3	No genotoxicity reports in original report.
Synthetic Wax	THF	100 µg/ml	Human lymphocytes	OECD TG 487 (in vitro mammalian cell micronucleus test); positive controls (cyclophosphamide, mitomycin C, vinblastine); exposed for 3 h with and without test article and 28 h with Cytochalasin B.	Did not induce micronuclei; non-mutagenic.	3	No genotoxicity reports in original report.
Synthetic Wax	acetone	preliminary toxicity test: 9.77 to 2500 µg/ml Main experiment: 2500 µg/ml	Human lymphocytes	OECD Guidelines for Testing of Chemicals (2006) No. 487, Draft Proposal for a New Guideline 487: in vitro micronucleus test; observed 4 and 20 h.	Did not induce statistically significant increases in cell frequency with micronuclei. Synthetic Wax considered non-clastogenic & non-aneugenic to human lymphocytes in vitro.	3	No genotoxicity reports in original report.

DMSO – dimethyl sulfoxide; MCA - 3-methylcholanthrene; MMS - methyl methanesulfonate; THF - tetrahydrofuran

Table 7. Carcinogenicity studies

Test Article	Vehicle	Dose/Concentration	Test Population/System	Protocol	Results	Reference	Notes
Paraffin	NR	~5700 mg/kg	Sprague Dawley rats (~5/wax/sex); control group (140 male / 157 female)	OECD TG 451 (carcinogenicity studies); 5 Paraffin waxes were included into a feed at 10% for 2 yr. Rats were weighted & inspected every week until all the rats passed away; histological examination was completed after death.	Survival rates and growth rates – unaffected; tumors were found but incidence was similar across all groups and no other toxic effects were found. No carcinogenic potential.	2	This study was not included in the original report, but the carcinogenicity studies showed that Paraffin was not carcinogenic.

Table 7. Dermal irritation and sensitization studies

Test Article	Vehicle	Dose/Concentration	Test Population/System	Protocol	Results	Reference	Notes
IRRITATION							
IN VITRO							
Montan Wax	none	25.6 to 25.9 mg per tissue	EpiDerm™ skin model	OECD TG 431; in vitro skin irritation: reconstituted human epidermis (Rhe) test method for 60 min; incubation time was 42 h	Montan Wax was not considered irritating and it did not reduce cell viability.	1	No in vitro dermal irritation assays provided in the original report; in vivo studies resulted in no irritation or sensitization.
ANIMAL							
Paraffin	none	0.5 ml	New Zealand white rabbits (n=3)	OECD TG 404 (acute dermal irritation/corrosion); exposed to dose for 4 h under semi-occlusive conditions and observed for 96 h before irritation score was measured.	Mean erythema score – 0.86; Mean edema score - 0.67; not irritating	2	This study was not included in the original report, but the skin irritation studies in animals resulted in no irritation.
Paraffin	none	0.5 mg	New Zealand male albino Rabbits (n=3)	OECD TG 404 (acute dermal irritation/corrosion); exposed to dose for 4 h under semi-occlusive conditions and observed for at 1, 24, 48, and 72 h before evaluation.	Slight erythema at 1 h mark; considered not irritating.	2	This study was not included in the original report, but the skin irritation studies in animals resulted in no irritation.
Synthetic Wax	none	0.5 ml	New Zealand white rabbits (n=3)	OECD TG 404 (Acute Dermal Irritation / Corrosion); test material was placed via cotton gauze patch and observations were made 1, 24, 48, & 72 h afterwards for primary irritation & scoring.	Primary irritation index was 0 & Synthetic Wax was classified as a non irritant to rabbit skin; no corrosive effects.	3	This study was not included in the original report, but the skin irritation studies showed that Synthetic Wax was a nonirritant.
SENSITIZATION							
ANIMAL							
Montan Wax	methyl ethyl ketone	0, 2.5, 5.0, 10.0 %	female mice (4/sex/dose)	OECD TG 429 (local lymph node assay); applied to dorsum of each ear lobe for 3 consecutive days; positive control was hexyl cinnamic aldehyde	All treated animals survived during scheduled study period with no signs of toxicity. Montan Wax was not considered skin sensitizing.	1	No dermal sensitization assays provided in the original report.
Synthetic Wax	olive oil and liquid paraffin	Intradermal -100% used for challenge testing after sighting test of 100 %, 50 %, 25 % and 12.5 %, 6.25 % and 3.125 % Topical – 50% was used for challenge testing after sighting testing of 50 %, 25 % and 12.5 %	Dunkin Hartley female guinea pigs (n=16; 11 test animals and 5 control animals)	OECD TG 406 (Skin Sensitization); positive control was α -Hexylcinnamaldehyde; sighting tests done first intradermally and topically before challenge tests were performed	Synthetic Wax produced a 0% sensitization rate & was classified as a nonsensitizer to guinea pig skin.	3	This study was not included in the original report, but the skin sensitization studies showed that Synthetic Wax was a nonsensitizer.

Table 8. Ocular irritation studies

Test Article	Vehicle	Dose/Concentration	Test Population	Protocol	Results	Reference	Note
IN VITRO							
Montan Wax	NR	300 ml (~167.6 mg)	In vitro HET-CAM (6 eggs for test substance, 3 eggs per control)	OECD TG 405 (acute eye irritation/corrosion); negative control: physiological sodium chloride solution; positive controls: sodium dodecyl sulphate (1% solution) and sodium hydroxide (0.1-N solution); observation time: 5 min at room temperature	Positive control – severe irritation in blood vessels Negative control – no irritation Montan Wax had mean irritation score of 0.0 and was classified as not an irritant.	¹	No ocular irritation assays provided in the original report.
ANIMAL							
Paraffin Wax	NR	0.1 ml	New Zealand female white Rabbits (n=3)	OECD TG 405 (acute eye irritation/corrosion); eyes remained unwashed and animals were observed for 72 h; redness was observed 1 h after treatment but no other effects were reported.	Not irritating to rabbits.	²	This study was not included in the original report, but the ocular irritation studies showed mild to no irritation.
Synthetic Wax	NR	0.1 ml	New Zealand male white Rabbits (n=3)	OECD TG 405 (acute eye irritation/corrosion); single rabbit treated then 2 additional rabbits were treated after considered response; assessment of damage/irritation made 1, 24, 48, & 72 h after treatment.	Synthetic Wax scored a maximum group mean score of 5.3. It was classified as a minimal irritant to rabbit eyes.	³	This study was not included in the original report, but the ocular irritation studies showed that Synthetic Wax was nonirritating.

Table 9. Case reports

Ingredient	Subjects	Protocol/Study Description	Results/Case Report Summary	Reference	Notes
Ceresin	37-yr-old female	Case report on subject that had recurring cheilitis on upper and lower lip due to the usage of lipstick containing Ceresin.	A 37-yr-old woman with a history of cheilitis and personal/family history of atopy including allergic rhinitis and eczema was patch tested in 2 separate sessions. The first test consisted of the North American Contact Dermatitis Group Baseline including 47 supplementary allergens and 4 of the patient's own products. The patient only had 1 positive reaction to her own product (Clinique 'Long last shine berry freeze' lipstick) and was negative to the other 125 allergens tested. In the second test, the woman had patch testing with the individual ingredients of the lipstick formulation and only tested positive to Ceresin. The controls on day 2 and day 4 were negative.	⁵	No case studies were provided in the original report, but clinical assessments showed no to mild reaction in lipstick formulations containing Ceresin.
Paraffin	35-yr-old female	Case report on subject that was diagnosed with Paraffin-induced ELP that developed into interstitial pulmonary disease with a 25 yr follow up.	In 1979, a 35-yr-old woman with a history of working in a factory where the workers were exposed to aerosolized Paraffin. Pulmonary function tests showed a restrictive pattern, BAL showed increased cell count and a high percentage of lymphocytes. She was diagnosed with Paraffin-induced ELP and was treated with steroid therapy and discontinued Paraffin exposure. Eventually in a 25 yr follow up, her CT scan indicated extensive lung fibrosis with diffuse ground-glass, interlobular septal thickening, intralobular opacities, traction bronchiectasis, and a subpleural honeycomb pattern.	⁶	No case studies were provided in the original report.
Paraffin	29-yr-old male	Case report on subject that developed foreign body granulomas due to the subcutaneous infiltration of liquid Paraffin.	A 29-yr-old male came into a sexually-transmitted disease unit with a painless papule on his penis that had been growing for several months. However, upon examination there a 4 cm large painful tumor with irregular wound margins and covered with pus. The edges also contained pus-filled fistulae. The patient mentioned he received a penicillin injection for syphilis a month ago. Eventually, the patient disclosed that he had underwent penile enlargement using liquid paraffin at age 15. The subject had a two-stage scrotum-skin graft as the primary wound closure to cover the penis after surgical excision was recommended.	⁷	No case studies were provided in the original report, but intradermal effects were mentioned in the original report.

Table 9. Case reports

Ingredient	Subjects	Protocol/Study Description	Results/Case Report Summary	Reference	Notes
Paraffin	25-yr-old male	Case report on subject that experienced acute aspiration pneumonia due to accidentally ingesting liquid Paraffin during the act of 'fire-breathing' or 'fire eating'.	A 25-yr-old male was admitted to the emergency unit experiencing dizziness, strong headaches, and a persistent dry cough with hemoptysis. He had a fever, shortness of breath, and trouble breathing. His initial treatment included oxygen therapy via mask, IV fluids, antibiotics, corticosteroids, bronchodilators, vitamins, anticoagulant, and Ranitidine. Eventually, after 17 d of treatment, the patient showed improvement where he was transferred to a regular stationary ward. He continued the oral antibiotics, IV fluids, and vitamin therapy until he was able to leave after 4 wk. The patient was monitored for 3 mo and an x-ray was done after 5 mo which showed a regression of the pulmonary changes.	⁸	No case studies were provided in the original report.
Paraffin	83-yr-old female	Case report on subject that developed ELP due to the usage of paraffin oil as a mouthwash for an extended period of time.	An 83-yr-old female with dyspnea, hypertension, obesity, and a productive cough was admitted. In an initial evaluation, it was suspected that she had a non-specific interstitial fibrosis. A chest CT scan revealed ground-glass opacities and consolidations in the lower lobes of the lungs. When the patient confirmed that she had been using liquid Paraffin oil to manage her xerostomia, she was diagnosed with ELP. Her treatment included discontinuing the usage of liquid paraffin oil, using artificial saliva products to help with her xerostomia, avoid irritants such as caffeine and tobacco. Eventually, there was a reduction in cough and no new symptoms reported.	⁹	No case studies were provided in the original report.
Paraffin	7-yr-old. female	Case report on subject that was given liquid paraffin to treat her constipation but developed lipoid pneumonia.	A 7-yr-old female with a history of chromosomal unstable translocation and developmental delay had recurrent lower respiratory tract infections and a persistent cough for 3 yr before being admitted to the pediatric pulmonology division. A chest CT revealed that she had consolidations in the right middle lobe, right lower lobe, and left lower lobe of her lungs. Her mother revealed that she had been giving her child liquid paraffin for chronic constipation. With this information, and bronchoscopy evaluation (resulted in 70% lipid-laden macrophage), she was diagnosed with lipoid pneumonia.	¹⁰	No case studies were provided in the original report.

BAL- bronchoalveolar lavage; CT - computed tomography; ELP - exogenous liquid pneumonia; IV – intravenous

REFERENCES

1. European Chemicals Agency (ECHA). ECHA Dossier: Montan Wax (CAS No. 8002-53-7). https://chem.echa.europa.eu/100.029.373/dossier-view/aa3c62e8-414c-47ef-a436-d3c13621c10b/1ed3a570-e3f4-4307-bb7f-1b36df7241ad_1ed3a570-e3f4-4307-bb7f-1b36df7241ad. Date Accessed: July 24th, 2025.
2. European Chemicals Agency (ECHA). ECHA Dossier: Paraffin Waxes and Hydrocarbon Waxes (CAS No. 8002-74-2). https://chem.echa.europa.eu/100.029.375/dossier-view/de2d1416-d5fd-4e0b-b9f9-7447ef25bba4/fc36e56d-2bf7-4c60-9498-dcd9613b8aaf_fc36e56d-2bf7-4c60-9498-dcd9613b8aaf?searchText=paraffin%20wax. Date Accessed: July 27th, 2025.
3. European Chemicals Agency (ECHA). ECHA Dossier: Paraffin waxes (Fischer-Tropsch), isomerization (CAS No. 2658498-20-3). [https://chem.echa.europa.eu/100.350.109/dossier-view/ab59ca73-8709-47dc-b42e-1fb3070e0cdc/b861b043-680a-4887-8146-c0119be3f947_b861b043-680a-4887-8146-c0119be3f947?searchText=Paraffin%20waxes%20\(Fischer-Tropsch\)](https://chem.echa.europa.eu/100.350.109/dossier-view/ab59ca73-8709-47dc-b42e-1fb3070e0cdc/b861b043-680a-4887-8146-c0119be3f947_b861b043-680a-4887-8146-c0119be3f947?searchText=Paraffin%20waxes%20(Fischer-Tropsch)). Date Accessed: June 27th, 2025.
4. Kempf M, Kimble RM, Cuttle L. Cytotoxicity testing of burn wound dressings, ointments and creams: a method using polycarbonate cell culture inserts on a cell culture system. *Burns: Journal of the International Society for Burn Injuries*. 2011;37(6):994–1000.
5. Powell M, Moreau L. Case report: allergic contact cheilitis caused by ceresin wax. *Contact Dermatitis*. 2012;66(1):46–47.
6. Descatha A, Mompoin D, Ameille J. Occupational paraffin-induced pulmonary fibrosis: a 25-year follow-up. *Occupational Medicine*. 2006;56(7):504–506.
7. Pónyai K, Marschalkó M, Hársing J, et al. Paraffinoma. *Journal der Deutschen Dermatologischen Gesellschaft = Journal of the German Society of Dermatology*. 2010;8(9):686–688.
8. Chibishev A, Simonovska N. Acute unintentional intoxication with paraffin in a 25-year old patient - clinical case report. *Journal of Forensic and Legal Medicine*. 2014;26:1–4.
9. Siebert K, Jassem E, Porzezińska M, Jelitto M, Bernard W. Lipoid pneumonia induced by aspiration of liquid paraffin. *Annals of agricultural and environmental medicine*. 2024;31(1):144–146.
10. Ayaz E, Ademhan D, Yüksel K, et al. Complicated pneumonia due to exogenous toxic substances in children. *The Turkish Journal of Pediatrics*. 2021;63(1):149–154.

JOURNAL OF THE AMERICAN COLLEGE OF TOXICOLOGY
Volume 3, Number 3, 1984
Mary Ann Liebert, Inc., Publishers

2

Final Report on the Safety Assessment of Fossil and Synthetic Waxes

Toxicological test data on Ozokerite, Ceresin, Montan Wax, Paraffin, Microcrystalline Wax, Emulsifying Wax N.F., Synthetic Wax and Synthetic Beeswax are presented. Based on the documented animal and clinical test data, it is concluded that these waxes are safe for use as cosmetic ingredients in the present practices of concentration and use.

FOSSIL AND SYNTHETIC WAXES: INTRODUCTION

The fossil waxes used in cosmetic products are grouped in this report according to their source. The mineral waxes, which include Ozokerite, Ceresin and Montan Wax, are derived from coal and shale. Paraffin and Microcrystalline Wax are derived from petroleum.^(1,2)

This report also includes a safety review of three synthetic waxes: Emulsifying Wax N.F., Synthetic Wax, and Synthetic Beeswax, which are manufactured to meet specific use requirements.

Each of the eight waxes is reviewed separately in this report.

OZOKERITE: CHEMICAL AND PHYSICAL PROPERTIES

Structure/Composition

Ozokerite is a naturally occurring fossil wax which consists of aliphatic series of straight-chain, branched-chain, and cyclic hydrocarbons, and some oxygenated resinous bodies. It has a delicate needle or short plate microcrystalline structure.^(1,2)

Ozokerite is found near soft shale, which acts as a molecular filter and condenser. It has been suggested that the wax was produced when small hydrocarbon molecules were polymerized under pressure into large ones. Waxes from different deposits have somewhat different chemical compositions and physical properties. Most of the commercial wax is mined in Eastern Europe.⁽²⁾ Commercial products called Ozokerite may be mixed with Paraffin to reduce its cost and with Carnauba Wax, resins, or asphaltum to increase its melting point or hardness.⁽¹⁻³⁾

Properties

Crude Ozokerite is black; after refining, its color varies from yellow to white. It hardens on aging, the hardness varying according to its source and refinement. For other specific properties, see Table 1.^(1,2,4-7)

Reactivity

Ozokerite is less stable than the solid paraffins, reacting readily with fuming sulfuric acid, chlorosulfonic acid, and concentrated nitric acid. It is water and alkali resistant.⁽²⁾

Refining and Analytical Methods

The newly mined Ozokerite is first melted in boiling water and separated from the other residues. The wax is then purified with concentrated sulfuric acid and decolorized with charcoal. Any remaining color may be bleached out with more sulfuric acid.^(1,2,7)

The chemical composition of Ozokerite varies according to its source. Procedures for separating the various chemical fractions and identification have been published. The various chemical fractions are separated and usually characterized by infrared spectroscopy and gas chromatography. Normal paraffins, isoparaffins, aromatics, and naphthalenes are normal constituents.⁽⁸⁾

USE

Noncosmetic Uses

Ozokerite is used in shoe and floor polish, leather dressings, pharmaceuticals, electrical insulation, cable wax, wax figure making, candles, paper

TABLE 1. Fossil Wax Properties.

Properties	Ozokerite	Ref.	Ceresin	Ref.	Montan Wax	Ref.
Melting pt.	63°-91°C	7	53°-79°C	17	85°-88°C	18
Specific gravity	0.85-0.95	6	0.92-0.94	6	1.00	1
Acid value	nil	7	nil	17	25-30	18
Saponification value	nil	7	nil	17	62-80	18
Iodine value	7.8-9.2	1	7-10	1	14-18	18
Refractive index	1.440	1	1.4416-1.4465	1	—	—
Color	Black, yellow, green white when pure	6	White, yellow	4,6	White, brown (crude)	4,6
Soluble in	Benzene	6	Alcohol	4,6	Carbon tetrachloride	4,6
	Turpentine		Benzene		Benzene	
	Kerosene		Chloroform		Chloroform	
	Carbon disulfide		Naphtha		Dichloroethylene	
	Isopropyl ether		—		Isopropyl ether	
	Trichloroethylene		—		Naphtha	
	—		—		Trichloroethylene	
	—		—		Turpentine	
Insoluble in	Ethyl and methyl alcohol	6	Water	4,6	Water	4,6
	Water		—		—	

coating, crayons, matches, cord coating, waterproofing cloth and electrotypers plates.^(1,2,9-14)

Purpose and Use in Cosmetics

Ozokerite lends stiffness to cleansing cream preparations and lessens the brittleness of cosmetic stick formulations. The wax adds strength and thermal stability to lipstick preparations and thixotropic properties to rouges; it holds the oil phase in hair creams.⁽¹⁵⁾

Ozokerite is used in lipsticks, baby products, eye and facial makeup preparations, manicuring, skin care, suntan, fragrance, and, noncoloring hair preparations (see Table 2).^(15,16)

Ozokerite is applied to the face, hands and general body surface. It also comes in contact with the nails, hair, scalp, and lips, and periocular skin.^(15,16)

The cosmetic product formulation computer printout which is made available by the FDA is compiled through voluntary filing of such data in accordance with Title 21 part 720.4 of the Code of Federal Regulations.⁽¹⁹⁾ Ingredients are listed in prescribed concentration ranges under specific product type categories. Since certain cosmetic ingredients are supplied by the manufacturer at less than 100% concentration, the value reported by the cosmetic formulator may not necessarily reflect the actual concentration found in the finished product; the concentration in such a case would be a fraction of that reported to the FDA. The fact that data are submitted only within the framework of preset concentration ranges also provides the opportunity for overestimation of the actual concentration of an ingredient in a particular product. An entry at the lowest end of a concentration range is considered the same as one entered at the highest end of that range, thus introducing the possibility of a two- to 10-fold error in the assumed ingredient concentration.

The 1976 FDA cosmetic ingredient data list Ozokerite in 1,269 formulations. The 1979 information lists it in 1,085 formulations in concentrations up to 50%. Ozokerite may come in contact with the body from a few hours (in suntan preparations) to several days (in hair conditioners).^(16,20)

BIOLOGICAL PROPERTIES

General Effects

Microbiological Effects

Ozokerite may be metabolized by some microorganisms, and also acts as a bacteriocidal agent for others. Metabolism of this wax by some microorganisms changes its chemical composition. After microbial action there may be 12%–38% fewer hydrocarbons than in the initial product, and there can be an increase in the concentration of resin-like compounds. Such decomposition is accompanied by an accumulation of oxygen-, nitrogen-, and sulfur-rich compounds. These changes are similar to those that occur in nature.^(21,22)

Ozokerite demonstrated an antibacterial effect against *Salmonella sp.*, *Shigella flexneri*, *Staphylococcus aureus*, *Shigella sonnei*, *Corynebacterium diphtheriae* at concentrations as low as 0.62%. This effect was attributed to the presence in the Ozokerite of trace elements and organic compounds.⁽¹²⁾

ASSESSMENT: FOSSIL AND SYNTHETIC WAXES

preparations	13	—	—	1	1	7	4	—
Night skin care preparations	9	—	1	—	2	6	—	—
Wrinkle smoothers (removers)	4	—	—	1	2	1	—	—
Other skin care preparations	8	—	—	3	2	3	—	—
Suntan gels, creams, and liquids	5	—	—	2	1	2	—	—
Other suntan preparations	3	—	—	—	—	3	—	—
1976 TOTALS	1269	—	48	253	95	632	241	—
1979 TOTALS	1085	85	28	168	161	535	105	3
<i>Ceresin</i>								
Baby lotions, oils, powders, and creams	1	—	—	—	1	—	—	—
Other bath preparations	2	—	—	2	—	—	—	—
Eyeliner	3	—	—	—	—	3	—	—
Eye shadow	48	—	48	—	—	—	—	—
Eye lotion	2	—	—	—	2	—	—	—
Eye makeup remover	2	—	—	—	2	—	—	—
Mascara	32	—	—	2	21	8	1	—
Other eye makeup preparations	17	—	—	5	5	7	—	—
Colognes and toilet waters	6	—	—	—	6	—	—	—
Perfumes	8	—	—	8	—	—	—	—
Other fragrance preparations	33	—	—	33	—	—	—	—
Hair conditioners	1	—	—	—	1	—	—	—
Tonics, dressings, and other hair grooming aids	7	—	—	1	2	3	1	—
Wave sets	1	—	—	—	1	—	—	—
Other hair preparations (noncoloring)	2	—	—	—	1	1	—	—
Blushers (all types)	19	—	—	2	2	13	2	—
Face powders	4	—	—	—	4	—	—	—
Makeup foundations	7	—	—	1	3	3	—	—
Lipstick	109	—	—	1	21	80	7	—
Makeup bases	7	—	—	1	1	2	1	2
Rouges	1	—	—	1	—	—	—	—
Other makeup preparations (not eye)	19	—	—	7	1	8	3	—
Deodorants (underarm)	1	—	—	—	—	1	—	—

TABLE 2. (Continued.)

Product category	Total no. containing ingredient	No. of product formulations within each concentration range (%)						
		Unreported concentration	>25-50	>10-25	>5-10	>1-5	>0.1-1	≤0.1
Skin cleansing preparations (cold creams, lotions, liquids, and pads)	29	—	—	4	9	13	2	1
Face, body, and hand skin care preparations (excluding shaving preparations)	5	—	—	1	—	1	3	—
Hormone skin care preparations	1	—	—	—	—	1	—	—
Moisturizing skin care preparations	10	—	—	2	3	5	—	—
Night skin care preparations	12	—	—	1	4	7	—	—
Wrinkle smoothers (removers)	3	—	—	1	—	2	—	—
Other skin care preparations	9	—	—	7	2	—	—	—
Suntan gels, creams, and liquids	1	—	—	—	1	—	—	—
Other suntan preparations	1	—	—	—	1	—	—	—
1976 TOTALS	403	—	48	80	94	158	20	3
1979 TOTALS	386	—	—	77	107	180	21	1
Montan Wax	5	—	—	5	—	—	—	—
Eyebrow pencil	18	—	—	—	—	18	—	—
Eye shadow	1	—	—	—	—	—	1	—
Perfumes	323	—	—	2	—	297	24	—
Lipstick	7	—	—	—	7	—	—	—
Rouges	—	—	—	—	—	—	—	—
Face, body, and hand skin care preparations (excluding shaving preparations)	1	—	—	—	—	—	1	—
1976 TOTALS	355	—	—	7	7	333	26	—
1979 TOTALS	309	—	—	—	5	295	9	—

Data from Refs. 16, 20.

Gastric Effects

The effects of different doses of Ozokerite on the secretory activity of dog stomachs isolated by the I.P. Pavlov method have been studied. Ozokerite suspended in 150 ml tap water was injected into the pouch in 10, 20, 50, or 100 mg/kg doses, with or without a food stimulus. Ozokerite had no effect on either the secretion of gastric juice or the total and free acidity at the 10 and 20 mg/kg levels; however, 50–100 mg/kg increased these variables.⁽²³⁾

Animal Toxicology

Acute

Oral toxicity

No toxic effects were reported after gastric administration to mice of up to 200 mg/kg of a 0.2% solution of Ozokerite, or to rabbits of up to 200 mg/kg of a 2.0% solution of the wax.⁽²⁴⁾

A lipstick formulation containing 4.5% Ozokerite and another containing 19.1% were tested for acute oral toxicity in rats. The formulation with 4.5% wax was administered by gavage to 10 Sprague–Dawley rats in a 20 mg/kg dose. No deaths or toxic effects resulted.⁽²⁵⁾ The formulation containing 19.1 percent wax was given by gavage to 10 CFE rats in a dose of 25 g/kg. One animal died from intestinal obstruction, but no toxic effects were seen.⁽²⁶⁾

Formulations of 5% (two lipstick formulations), 13% (two blush cream formulations), 28% and 29% (lipstick formulations), were tested for acute oral tox-

TABLE 3. Acute Oral Toxicity: Ozokerite.

Wax conc. in test soln. (%)	Dose of test subst./kg	Solution or formulation	Species and number	Comments ^a	Ref.
0.2	50–200 mg	Solution	Mice	Ingastric administration. No toxic effect.	24
2.0	10–200 mg	Solution	Rabbits	Ingastric administration. No toxic effect.	24
4.5	20 g	Formulation-lipstick	10 Sprague–Dawley rats	Gavage. No deaths; no toxic effects.	25
5	15 g	Formulation-lipstick	5 albino rats	LD ₅₀ not determined.	27
5	15 g	Formulation-lipstick	5 albino rats	LD ₅₀ not determined.	28
13	5 g	Formulation-blush cream	5 albino rats	LD ₅₀ not determined.	29
13	5 g	Formulation-blush cream	5 albino rats	LD ₅₀ not determined.	30
19.1	25 g	Formulation-lipstick	10 CFE rats	Gavage. 1/10 died from intestinal obstruction; no toxic effects.	26
28	5 g	Formulation-lipstick	5 albino rats	LD ₅₀ not determined.	31
29	5 g	Formulation-lipstick	5 albino rats	LD ₅₀ not determined.	31

^a 21 CFR 1500.3

icity in five albino rats per sample. The animals were intubated with the compounds after an overnight fast and observed for seven days. The LD₅₀s for the formulations were not reached⁽²⁷⁻³¹⁾ (see Table 3).

Dermal irritation/toxicity

Eight samples of a 50% concentration of Ozokerite in petrolatum were tested for skin irritation on groups of nine albino rabbits. The rabbits were given three consecutive 24 h applications. The first application of each sample produced erythema in one to four animals. Erythema decreased with each subsequent application, and after the third (final) application, three samples produced no irritation. The remaining five samples caused erythema in one or two animals per group⁽³²⁻³⁹⁾ (see Table 4).

A lipstick formulation containing 4.5% Ozokerite was tested for dermal irritation on the intact and abraded skin of six New Zealand white rabbits. Three consecutive 24 h applications produced mild erythema in one rabbit's intact skin, and mild erythema in the abraded skin of three rabbits.⁽²⁵⁾ Another lipstick formulation containing 19.1% Ozokerite produced no irritation when tested under the same conditions.⁽²⁶⁾

Two lipstick formulations each containing 5% Ozokerite were tested for dermal irritation; nine albino rabbits were used per formulation. Both samples were applied full strength under a 24 h closed patch. One formulation produced a PII of 0.17 out of a possible score of 8.0, and the other produced no irritation (score = 0.0).^(27,28) Two lipstick formulations, one containing 28% and the other 29% Ozokerite, were tested for dermal irritation. Both were minimally irritating.⁽³¹⁾ Three 24 h applications of a lipstick containing 19.1% Ozokerite produced no irritation on the abraded and intact skin of six rabbits.⁽²⁶⁾

One of two blush cream formulations, each containing 13% Ozokerite and tested for dermal irritation, was minimally irritating and the other produced no irritation.^(29,30)

Ocular irritation

Eight different samples of 50% Ozokerite in petrolatum were tested according to the Draize method. Each 0.1 ml sample was instilled into one eye of each of six rabbits. Five samples produced no irritation after two days.⁽³²⁻³⁶⁾

Similar procedures were used to test eight cosmetic formulations: two lipstick formulations containing 5% Ozokerite were tested as above. One product produced no irritation. The second caused mild irritation in one rabbit on Day 1, but the irritation had cleared by Day 2.^(27,28) One of two blush cream formulations containing 13% Ozokerite produced mild eye irritation in one rabbit at four days and the other produced no irritation.^(29,30) Two lipstick formulations, one containing 28% and the other containing 29% Ozokerite, produced mild eye irritation for two days and none thereafter.⁽³¹⁾ One lipstick formulation containing 4.5% Ozokerite and another containing 19.1% were each tested for ocular irritation in six New Zealand white rabbits. When 0.1 ml volume was instilled into one eye per animal, no irritation occurred up to 72 hours^(25,26) (see Table 5).

TABLE 4. 24-hour Acute Dermal Irritation/Toxicity: Ozokerite.

Wax conc. (%)	Dose of test subst.	Solution or formulation	No. of albino rabbits	Primary irrit. score		Contact time	Comments	Ref.
					Max score (8.0)			
50	0.5 ml	Solution	9	—	—	3 applic. at 24 h intervals	Irritation in 2 rabbits at Day 1; 1 at Day 3	32
50	0.5 ml	Solution	9	—	—		Irritation in 4 rabbits at Day 1; 2 at Day 2	33
50	0.5 ml	Solution	9	—	—	3 applic. at 24 h intervals	Irritation in 4 rabbits at Day 1; 1 at Day 2; 0 at 3	34
50	0.5 ml	Solution	9	—	—		Irritation in 4 rabbits at Day 1; 1 at Day 2; 0 at 3	35
50	0.5 ml	Solution	9	—	—	3 applic. at 24 h intervals	Irritation in 2 rabbits at Day 1; 1 at Day 2 and 3	36
50	0.5 ml	Solution	9	—	—	3 applic. at 24 h intervals	Irritation in 2 rabbits at Day 1; 1 at Day 3	37
50	0.5 ml	Solution	9	—	—	3 applic. at 24 h intervals	Irritation in 4 rabbits at Day 1; 2 at Day 2; 1 at Day 3	38
50	0.5 ml	Solution	9	—	—	3 applic. at 24 h intervals	Irritation in 1 rabbit at Day 1; 3 at Day 2; 2 at Day 3	39

TABLE 4. (Continued.)

Wax conc. (%)	Dose of test subst.	Solution or formulation	No. of albino rabbits	Primary irrit. score Max score (8.0)	Contact time	Comments	Ref.
4.5	0.5 g	Formulation- lipstick	6	+1 intact site +3 abraded site	3 applic. at 24 h intervals	Open patch on abraded and intact skin. One +1 score on intact skin (mild erythema) 3 +1 on abraded skin (mild erythema)	25
5	0.5 ml	Formulation- lipstick	9	0.17	24 h	Potential for minimal irritation	27
5	0.5 ml	Formulation- lipstick	9	0.0	24 h	No irritation	28
13	0.5 ml	Formulation- blush cream	6	0.9	24 h	Potential for minimal irritation	30
13	0.5 ml	Formulation- blush cream	6	0.0	24 h	No irritation	29
19.1	0.5 g	Formulation- lipstick	6	0.0	3 applic. at 24 h intervals	Open patch on abraded and intact skin. All zero scores	26
28	0.5 ml	Formulation- lipstick	6	0.08	24 h	Potential for minimal irritation	31
29	0.5 ml	Formulation- lipstick	6	0.17	24 h	Potential for minimal irritation	31

TABLE 5. Ocular Irritation: Ozokerite.

Wax conc. in petrolatum (%)	Dose	Solution/formulation	No. of albino rabbits	Observ. time	Comments	Ref.
50	0.1 ml	Solution	6	48 h	One instillation. No irritation.	32
50	0.1 ml	Solution	6	48 h	One instillation. No irritation.	33
50	0.1 ml	Solution	6	48 h	One instillation. No irritation.	34
50	0.1 ml	Solution	6	48 h	One instillation. No irritation.	35
50	0.1 ml	Solution	6	48 h	One instillation. No irritation.	36
50	0.1 ml	Solution	6	72 h	One instillation. Minimally irritating at 48 h; no irritation at 72 h.	37
50	0.1 ml	Solution	6	72 h	One instillation. Minimally irritating at 48 h; no irritation at 72 h.	38
50	0.1 ml	Solution	6	72 h	One instillation. Minimally irritating at 48 h; no irritation at 72 h.	39
4.5	0.1 ml	Formulation-lipstick	6	72 h	One instillation. No irritation (Score = 0).	25
5.0	0.1 ml	Formulation-lipstick	6	24 h	One instillation. No irritation.	27
5.0	0.1 ml	Formulation-lipstick	6	48 h	One instillation. Mild irritation seen in one rabbit at 24 h; cleared by Day 2.	28
13	0.1 ml	Formulation-blush cream	6	7 days	One instillation. Mild irritation to Day 4.	29
13	0.1 ml	Formulation-blush cream	6	24 h	One instillation. No irritation.	30
19.1	0.1 ml	Formulation-lipstick	6	72 h	One instillation. No irritation (Score = 0).	26
28	0.1 ml	Formulation-lipstick	6	72 h	One instillation. Mild irritation to Day 2.	31
29	0.1 ml	Formulation-lipstick	6	72 h	One instillation. Mild irritation to Day 2.	31

Clinical Assessment of Safety

The results of these tests are summarized in Table 6.

Skin Irritation/Sensitization

Single 24 hour patch test

Six lots of 100% Ozokerite were each tested on 20 subjects per lot in a single insult 24 h occlusive patch test. The combined results of all tests showed one of 120 test subjects had a moderate reaction, 25 had mild to minimal reactions, and 94 produced no reaction.^(40,41)

Each of seven lipstick formulations containing Ozokerite at 5%, 28%, and 29% were applied full strength under occlusion to the upper backs or forearm of 20 subjects for 24 h. Two separate blush cream formulations containing 13% Ozokerite were similarly tested on two groups of 18 and 19 subjects. Only one volunteer in the one blush cream test showed a perceptible erythema. None of the other 176 subjects tested developed a product-induced irritation.^(42-44,46-49)

Modified Draize–Shelanski–Jordan test

A lipstick product containing 4.5% Ozokerite, applied under occlusion to 201 subjects for 24 h, caused no allergic or irritant responses.⁽²⁵⁾

TABLE 6. Clinical Data: Ozokerite.

<i>TEST</i>						
<i>Wax conc. (%)</i>	<i>Neat/formulation</i>	<i>No. of subjects</i>	<i>Contact time</i>	<i>PII</i>	<i>Comments</i>	<i>Ref.</i>
<i>24 h Patch Test</i>						
100	Neat	20	24 h	—	1/20 showed barely perceptible erythema; 1/20 had definite slight erythema.	41
100	Neat	20	24 h	0	No irritation; 1/20 mild irritation.	41
100	Neat	20	24 h	—	1/20: barely perceptible erythema; 1/20: definite erythema; 1/20: moderate erythema.	41
100	Neat	20	24 h	—	2/20: barely perceptible; 6/20: definite erythema; 1/20: moderate erythema.	40
100	Neat	20	24 h	—	6/20: definite erythema	40
100	Neat	20	24 h	—	5/20: definite erythema	40
5	Formulation-lipstick	20	24 h	0	No irritation	42
5	Formulation-lipstick	20	24 h	0	No irritation	43
5	Formulation-lipstick	20	24 h	0	No irritation	44
5	Formulation-lipstick	20	24 h	0	No irritation	45
5	Formulation-lipstick	20	24 h	0	No irritation	46
13	Formulation-blush cream	18	24 h	0.03	Barely perceptible erythema in 1 out of 18.	47
13	Formulation-blush cream	19	24 h	0	No irritation	48
28	Formulation-lipstick	20	24 h	0	No irritation	49
29	Formulation-lipstick	20	24 h	0	No irritation	49
<i>Modified Draize–Shelanski–Jordan Test</i>						
4.5	Formulation-lipstick	201	1 week	0	No allergic reactions or irritant responses were elicited by the product.	25
<i>Contact Allergy Test. Repeat</i>						
<i>Insult Patch Test</i>						
13	Formulation-blush cream	300	4 weeks	0	No allergic responses	50
<i>21-Day Cumulative Irritancy Assay Test</i>						
13	Formulation-blush cream	12	21 days	1/756	Product scored 1 out of a possible 756	51

ASSESSMENT: FOSSIL AND SYNTHETIC WAXES

55

Repeated insult patch test

None of the 300 subjects tested with a blush product containing 13% Ozokerite had an allergic response during the four weeks of the study.⁽⁵⁰⁾

21 day cumulative irritancy test

A formulation containing 13% Ozokerite was applied to the skin of 12 panelists; a sensitization score of one out of a possible 756 was recorded.⁽⁵¹⁾

Further toxicological testing on Ozokerite in a mixture is discussed in "Waxes: Plant and Insect"⁽⁵²⁾ under "Mixture Candelilla Wax (4%), Ozokerite (5%), Paraffin Wax (2.5%), and Carnauba Wax (3%) in a formulation."

CERESIN: CHEMICAL AND PHYSICAL PROPERTIES**Structure/Composition**

Ceresin Wax, a microcrystalline mixture of complex hydrocarbons, is produced by the purification of Ozokerite with sulfuric acid and filtration through bone black.⁽¹⁷⁾ The wax is available in various grades, with melting points that vary over a wide range. The name "Ceresin Wax" is also used to designate the commercial compound variety with a paraffin base.^(1,2)

See "Ozokerite" for chemical and physical properties.

BIOLOGICAL PROPERTIES**Animal Toxicology****Acute***Oral toxicity*

A lipstick formulation containing 2% Ceresin was tested for acute oral toxicity in 10 rats. A dose of 5 g/kg was administered by oral intubation and the LD₅₀ was not determined.⁽⁵³⁾

Dermal irritation

The primary skin irritation of a lipstick product containing 2% Ceresin was tested on six rabbits according to the Draize method. A 0.5 ml dose of the formulation produced minimal irritation, with a PII of 0.79 out of a maximum possible score of 8.0.⁽⁵³⁾

Five products—one cleansing cream containing 5% Ceresin, another cleansing cream with 6% Ceresin, and the other three eye creams containing 6% Ceresin—were each tested for acute skin irritancy on four New Zealand albino rabbits. A 0.5 ml undiluted sample was applied under occlusion to the clipped, intact, or abraded skin of each animal for 24 h. Sites were evaluated and scored 1, 48, and 72 h after patch removal, according to the Draize Method (maximum PII score is 8.0). The PII of the 5% sample was 1.25.⁽⁵⁴⁾ The other four samples had PIIs of 1.63, 1.00, 0.88, and 0.8.⁽⁵⁵⁻⁵⁸⁾ The irritation was minimal to mild (see Table 7).

Dermal toxicity

The acute dermal toxicity of a lipstick formulation containing 2.0% Ceresin was tested in a dose of 2.0 g/kg on 10 rabbits. The percutaneous LD₅₀ for the formulation was not determined⁽⁵³⁾ (see Table 7).

Ocular irritation

The application of an undiluted lipstick product containing 2% Ceresin caused no eye irritation in six rabbits (scores of 0.0 at 24, 48, and 72 h).⁽⁵³⁾

Three formulations, two eye creams and one cleansing cream, each containing 6% Ceresin, were tested for ocular irritation on five New Zealand rabbits each. The undiluted test material (0.1 ml) was instilled without irrigation into one eye of each rabbit and the reactions observed after 1, 2, 4, 48, and 72 h, and four and seven days. The maximum possible score for irritation was 110. One eye cream product caused mild irritation only in the conjunctivae with scores of 8, 4, 2, and 0.8 after 1, 24, 48, and 72 h, respectively. No irritation was seen thereafter.⁽⁵⁹⁾ The second eye cream product caused iridial irritation (score = 1) only at 1 h. Conjunctival irritation occurred up to 48 h with scores of 6, 2, and 0.4 after 1, 24, and 48 h.⁽⁶⁰⁾ The cleansing cream product caused similar iridial irritation after 1 h (score = 2). Conjunctival irritation was seen at 1, 24, and 48 h with scores of 8, 2, and 6.8, respectively⁽⁶¹⁾ (see Table 8).

TABLE 7. Acute Dermal Irritation/Toxicity: Ceresin.

Wax conc. (%)	Dose of test subst. (per kg)	Type of formulation	No. of albino rabbits	Primary Irrit. score Max. score	Contact time (h)	Observ. time (h)	Comments	Ref.
<i>Irritation</i>								
2	0.5 ml	Lipstick	6	0.79/8.0	24	—	Minimal irritation	53
5	0.5 ml	Cleansing cream	4	1.25/8.0	24	72	Mild irritation	54
6	0.5 ml	Cleansing cream	4	1.63/8.0	24	72	Mild irritation	55
6	0.5 ml	Eye cream	4	1.0/8.0	24	72	Mild irritation	56
6	0.5 ml	Eye cream	4	0.88/8.0	24	72	Minimal irritation	57
6	0.5 ml	Eye cream	4	0.8/8.0	24	72	Minimal irritation	58
<i>Toxicity</i>								
2.0	2 g	Lipstick	10	—	—	—	LD ₅₀ was not determined	53

TABLE 8. Ocular Irritation: Ceresin.

Wax conc. (%)	Type of formulation	Dose	No. of albino rabbits	Observ. time	Comments	Ref.
2.0	Lipstick	—	6	72 h	Scores = 0.0	53
6.0	Eye cream	0.1 ml	5	7 days	Mild conjunctival irritation to 72 h.	59
6.0	Eye cream	0.1 ml	5	7 days	Mild iridial irritation after 1 h. Mild conjunctival irritation to 48 h.	60
6.0	Cleansing cream	0.1 ml	5	7 days	Mild iridial irritation after 1 h. Mild conjunctival irritation to 48 h.	61

Clinical Assessment of Safety

One lipstick formulation containing 2% Ceresin was tested with the Schwartz–Peck prophetic patch test and the Draize–Shelanski repeated insult patch test. In the Schwartz–Peck test, “virtually” no reactions occurred in the 102 subjects under open and closed patches. In order to evaluate photosensitivity, the test sites were irradiated for one minute with a Hanovia Tannette Mark I quartz UV source (150 W) at a distance of 12 in. Irradiation occurred after the second insult patch had been read and the site was evaluated 48 h after exposure. The Draize–Shelanski test caused “virtually” no reactions in 50 subjects under open and closed patch and ultraviolet test conditions.⁽⁵³⁾

A lipstick containing 2% Ceresin was tested for human skin irritation and sensitization in a Schwartz–Peck prophetic patch test. The undiluted product was applied under occlusion to the skin of the cleansed upper back and under open patch conditions to the inner upper arm. After 48 h, the patches were removed and the sites were graded. Following a 14-day nontreatment period, second open and closed patches were applied and read 48 h later. The reading of this second insult was followed by a test for UV sensitivity. The sites where closed patches had been applied were irradiated with ultraviolet light from a Hanovia Tannette Mark I quartz lamp (150 W) at a distance of 12 in. These sites were read 48 h after irradiation. Out of the 1,078 panelists, one showed a weak, nonvesicular reaction under closed patch conditions after the first insult. No other reactions occurred after the second insult or after UV irradiation.⁽⁶²⁾

A repeated insult patch test was conducted on the same lipstick formulation containing 2% Ceresin. The undiluted product was applied under occlusion to the skin of the back for 24 h and under open conditions to the upper arm. The patches were removed, the sites graded, and the skin was allowed to recuperate for 24 h before the next in a series of 10 patches were applied. After the last induction patch, a two- to three-week rest period occurred. A challenge 48 h patch was applied to untreated sites. It was then removed and the reaction was graded. To assess UV sensitivity, sites were irradiated with an ultraviolet source after the first, fourth, seventh, and tenth induction patches, and after the 48 h challenge patch was removed. Of the 506 panelists tested in this study, one had a weak, nonvesicular reaction under the closed patch after the second induction and one had an edematous reaction under the closed patch after the sixth induction. No other reactions occurred.⁽⁶²⁾

For additional testing on “Beeswax/Ceresin Wax Composites”, see “Waxes: Plant and Insect.”⁽⁵²⁾

BEESWAX/CERESIN COMPOSITE: BIOLOGICAL PROPERTIES

Animal Toxicology

Acute

Oral toxicity

A formulation containing 6% Beeswax and 6% Ceresin was tested for acute oral toxicity in five male and five female Wistar albino rats. The animals were fasted for 16 h and given an 80.0 ml/kg formulation dose by oral intubation.

Observations were made 1 and 6 h after dosing and daily for 14 days. There were no deaths.⁽⁶³⁾

Ocular irritation

A cream formulation containing 6% Ceresin and 6% Beeswax was tested for acute eye irritation. A 0.1 ml sample was instilled into the conjunctival sac of one eye of each of nine New Zealand rabbits. The untreated eyes served as controls. Three of the treated eyes were washed with 20 ml of deionized water 30 sec after instillation. Ocular reactions were scored after 24, 48, and 72 h and four and seven days. After 24 h, four of the six rabbits with unwashed eyes showed minimal chemosis and two showed minimal conjunctival redness. No irritation occurred in the washed eyes.⁽⁶⁴⁾

Clinical Assessment of Safety

Skin Irritation/Sensitization

Prophetic patch test

A formulation containing 6% Beeswax and 6% Ceresin was tested for irritation and sensitization with the Schwartz–Peck prophetic patch test. Plastic bandages impregnated with the test product were applied to the cleansed upper backs of each of the 98 persons. Simultaneously, open patches were applied to the left volar forearm surface and the results from both applications were read 48 h later. After a rest of 14 days, there was a second application of the product with open and closed patches. To evaluate photosensitization, the test area of the back was exposed to UV radiation (Hanovia Tannette Mark I lamp) at a distance of 12 in for 1 min after the second insult had been read. This exposure area was read 48 h later. None of the subjects showed irritation or sensitization after any insult.⁽⁶⁵⁾

Repeated insult patch test

A formulation containing 6% Beeswax and 6% Ceresin was tested with the Draize–Shelanski repeated insult procedure on 49 subjects. Patches impregnated with the formulation were applied to the cleansed upper backs of each subject every Monday, Wednesday, and Friday for three and one-half weeks, for a total of 10 insults. An open patch was simultaneously applied to the left forearm, and both sets of patches were removed and the areas read 48 h later. After a 14-day rest, an eleventh open and closed patch was applied and read 48 h later. Photosensitization was evaluated with a UV light source that emitted a wavelength of 360 nm. Formulation-exposed sites were irradiated for 1 min after the final reading was taken. The light-exposed sites were read 48 h later. None of the 49 panelists showed any irritation or sensitization.⁽⁶⁵⁾

21 day cumulative irritancy test

A cream containing 6% Beeswax and 6% Ceresin was tested for 21-day cumulative irritancy on 14 panelists. Patches impregnated with the cream were applied to the back of each panelist for 21 consecutive days. The total irritation score of the entire test population was 6.4 out of a possible 630.⁽⁶⁶⁾

Contact sensitization

The contact sensitization potential of a formulation containing 6% Beeswax and 6% Ceresin was patch tested on 22 subjects. Sites on the volar forearm were pretreated with 5% sodium lauryl sulfate 24 h before the test material was applied under occlusion for alternate 48 h periods. After a 10- to 14-day rest, challenge patches were applied under occlusion to adjacent, fresh sites for 48 h. The preparation produced no irritant or allergic reactions.⁽⁶⁷⁾

In-use test

One hundred women who used a cream preparation containing 6.0% Beeswax and 6.0% Ceresin daily for two weeks experienced no irritation.⁽⁶⁸⁾

MONTAN WAX: CHEMICAL AND PHYSICAL PROPERTIES**Structure/Composition**

Montan Wax is a bituminous wax that occurs in lignites (woody coals) from central Europe, Australia, New Zealand, Russia, Great Britain, and the United States. It is extracted with a volatile solvent.^(1,2,18,69-71)

Montan Wax consists of monohydric alcohol esters, high-molecular-weight acids, and free alcohols.⁽¹⁸⁾ The crude wax also contains resinous and sulfur compounds (see Table 9).⁽¹⁾

The refined wax has a higher content of free wax acids than does the crude wax, and it is substantially free of the esters of C₂₀, C₂₂, and C₂₄ acids. The resin content varies according to geographical source.^(1,69-72)

TABLE 9. Composition of Montan Wax.

<i>Component</i>	<i>Amount Reported (%)</i>	<i>Ref.</i>
Esters of wax acids:	58-59	1
1. Octacosyl esters of C ₂₀ , C ₂₂ , C ₂₄ acids		
2. Octacosyl cerotate		
3. Octacosyl hydroxyoctacosonate		
4. Montanyl montanate		
5. Ceryl octocosonate		
Free wax acids:	17-10	
1. Carbocerotic acid		
2. Montanic acid		
3. Melissic acid		
Free primary monohydric alcohols:	3-4	
1. Tetracosonol		
2. Hexacosonol		
Secondary alcohols:	1	
1. Montanol		
Resins:	10-12	
1. C ₂₀ H ₃₀ O ₂ , and neutral resins		
2. C ₂₄ H ₃₄ O ₂ , Montan resin		
Ketones:	Less than 10	
1. Cerotones		
2. Montanones C ₂₉ H ₅₉ CO		

Properties

Crude Montan Wax is brown but becomes white after purification. The hardest of the nonvegetable waxes, Montan Wax breaks with a conchoidal fracture; in the melt it is a good solvent for basic dyes. It shows good wetting and flow in oil solutions and is fairly resistant to oxidation. For other specific properties see Table 1. ^(1,2,4,6,18)

Reactivity

Montan Wax undergoes decomposition when distilled at ordinary atmospheric pressures; in a current of steam, in vacuo, a white crystalline montanic acid is produced. ⁽⁶⁹⁾

Refining and Analytical Methods

Montan Wax is removed from lignite coal by solvent extraction. It is deresinified by solvent processing followed by chromic acid oxidation and may be further purified by concentrated and fuming sulfuric acid, chlorosulfonic acid, and alkali metal bisulfites treatment. One method of purification recommends saponification of the wax to its component alcohols and acids. The alcohol component may be decolorized by activated carbon, and the acid component by treatment with oxidizers. The results may then be reesterified. ^(1,2,69-73)

Microdistillation methods and x-ray analysis are used to determine the acid content of this wax. Thin layer, column, and gas chromatography can be used to separate and identify the constituents. Differential thermal analysis, vapor pressure osmometry, infrared spectroscopy, and nuclear magnetic resonance are also used to determine the constituents. ^(69,70,71,73-78)

Impurities

Purified Montan Wax has no known impurities, whereas crude Montan Wax may contain sulfur and resins. ⁽¹⁸⁾

USE

Noncosmetic Uses

Montan Wax is used as a substitute for Carnauba Wax as a polish, in the manufacture of rubber, in printing ink, electrical insulation, leather finishes and dressings, carbon paper, grease, phonograph records, and waterproofing. ^(1,2)

Purpose and Use in Cosmetics

Montan Wax is used in eye and facial makeup preparations, in fragrance formulations, and in skin care preparations. It stiffens eyebrow pencil, eye shadow and rouge, and is used as the solid phase of fragrance preparations. It gives form to lipsticks and is a constituent of creams for skin care products (see Table 2). ^(15,16)

The 1976 FDA voluntary submission of cosmetic data lists Montan Wax in 355 cosmetic formulations. The 1979 data listed it in 309 preparations. It is used in cosmetics which are applied near the eyes, to lips, and the general body surface. ⁽¹⁶⁾ The wax is used in concentrations varying from 0.1% in lipsticks to 25%

in some eyebrow pencils (see Table 2). It may come in contact with the body for a few hours (in fragrance preparations) to several hours daily (in makeup and skin care preparations).⁽¹⁶⁾

BIOLOGICAL PROPERTIES

Animal Toxicology

Acute

Dermal irritation

A single application (2 and 3 g/kg) of Montan Wax to the intact skin of rabbits caused no irritation.⁽⁷⁹⁾

Intraperitoneal injection

The single, minimum toxic dose of Montan Wax administered by intraperitoneal injection in mice was 7.5 g/kg.⁽⁷⁹⁾

Subchronic

Oral toxicity

Ten animals (type unspecified) that were fed Montan Wax in 1 g/kg doses for four months had no lesions; however, an increase in the weights of the adrenal glands and kidneys was observed.⁽⁷⁹⁾

Dermal toxicity/irritation

Application of 200 g/kg of Montan Wax to the backs of rabbits for 30 days produced slight, transient hyperemia, which disappeared after 8–10 days.⁽⁷⁹⁾

Inhalation toxicity

Animals showed no behavioral changes when exposed for three months to a gaseous mixture of Montan Wax (concentration of hydrocarbon 0.11 to 0.36 mg/l, benzene 0.01–0.019 mg/l).⁽⁷⁹⁾

Clinical Assessment of Safety

Repeated Insult Patch Test

Four lipstick formulations containing Montan Wax were tested for irritation and sensitization using a Modified Draize–Shelanski–Jordan procedure. The first product, containing 1.61% Montan, was applied under occlusion in eight 24 h induction patches to the upper backs of 92 women and 18 men. Following an 11-day nontreatment period, two consecutive 48 h challenge patches were applied to the back. The material caused no irritation or sensitization.⁽⁸⁰⁾

Similar studies were conducted on lipstick products containing 1.93%, 1.81%, and 2.53% Montan Wax. Subjects testing each formulation received 10 occlusive 24 h induction patches, followed by a 13-day rest. Each subject then received two 48 h challenge patches seven days apart. In 54 women testing the

lipstick containing 1.93% wax, 41 women and 11 men testing the lipstick containing 1.81% wax, and 49 women and six men using the product with 2.53% wax, no irritation or sensitization occurred.⁽⁸¹⁻⁸³⁾

PARAFFIN: INTRODUCTION

Paraffin and Microcrystalline Wax are distillation products of petroleum. The crystalline structure of Paraffin is different from that of Microcrystalline Wax: Paraffin solidifies into plates, whereas Microcrystalline forms small, poorly-defined needles. According to one theory that attempts to explain this difference, the Microcrystalline Wax is associated with the heavy, residual stocks of petroleum and cannot be separated completely from the accompanying oil; it is, therefore, unable to form any well defined crystals. Paraffin is only associated with the light distillates and can readily be separated into macrocrystalline types.^(1,2)

Paraffin is obtained from a distillate of crude petroleum by cooling, filtering, and pressing to remove the residual oil. Refined Paraffin contains about 0.5% oil.^(1,2)

Microcrystalline Wax is separated from the Paraffin distillate residue by solvent extractions and centrifugation. Its oil content varies from 2% to 12%.^(1,2)

CHEMICAL AND PHYSICAL PROPERTIES

Structure/Composition

Paraffin is a mixture of organic compounds characterized by relatively large crystals; it contains solid hydrocarbons of the methane series and a small percentage of other organic entities. The hydrocarbon content of Paraffin varies according to the source of the crude petroleum. The four generic hydrocarbon classes of petroleum are in general the paraffins (saturated, open-chain hydrocarbons); the olefins (unsaturated, open chains); the naphthalenes (saturated, cyclic hydrocarbons, possibly with multiple ring nuclei); and the aromatics (unsaturated, cyclic hydrocarbons). The constituents of Paraffin have carbon numbers that range from 18 to 36.^(1,2) The ring structures are the tri-, tetra-, and pentacyclic compounds, but these are usually removed in the refining process. Paraffin crystallizes in plates and needles and an unidentifiable type of crystal called "malcrystalline." The plate types are straight-chain hydrocarbons and comprise the basic fraction of Paraffin; the needle and malcrystalline types are assumed to be branched-chain hydrocarbons.⁽⁸⁴⁾

Properties

The properties of Paraffin are listed in Table 10.^(1,2,4-6,85)

Reactivity

Paraffin is a relatively inert material that tends to oxidize when it is melted and large surface areas are exposed to oxygen. No information was available on the reactivity of this wax.^(1,2)

TABLE 10. Properties of Petroleum Waxes.

Properties	Paraffin	Ref.	Microcrystalline	Ref.
Melting point	43°–65°C	6	63–90.5	5,85
Specific gravity	0.880–0.915	6,1	0.90–0.94 (at 15.5°C)	1,85
Acid value	0	1	0.1–0.2	1
Saponification value	0	1	0.05–< 2.0	1
Iodine value	—		1.5	1
Color	White, translucent	4,6	Pale amber to white	5
Soluble in	Benzene	1,6	Carbontetrachloride	2
	Ligroin			
	Warm alcohol			
	Chloroform			
	Turpentine			
	Carbon disulfide			
Insoluble in	Olive oil			
	Water	4,6	Water	
	Acids		—	

Analytical Methods

The constituents of Paraffin can be identified by various chromatographic methods. Infrared spectrophotometry and fluorescence of the molten wax can determine aromatics, and thermal analysis can determine specific physical properties. The percent of unsaponifiable matter and hydrocarbon content may be determined by alcohol dilution. The quality of the refined product may be tested by the determination of melting point, tensile strength, color, and oil content. ^(1,2,84,86-89)

Impurities

Refined Paraffin used in the cosmetic industry like other petroleum waxes may contain trace impurities of polycyclic aromatic hydrocarbons (PAH) which have been shown to be carcinogenic. Improved refining techniques have been developed to eliminate these impurities. ⁽⁸⁸⁾

USE

Noncosmetic Uses

Paraffin is used in candles, waterproofing, lubricating, food coating, fruit and vegetable protection, waxed paper, polishes, crayons, pharmaceutical ointments and salves, and electrical insulation. ^(1,2,19)

Purpose and Use in Cosmetics

Paraffin is used in baby products, eye and facial makeup preparations, fragrance products, coloring and noncoloring hair preparations, manicuring products, personal cleanliness preparations, shaving and skin care products, and suntan preparations. ^(4,5,15,16,20,90)

TABLE 11. Product Formulation Data.

Product category	Total no. containing ingredient	No. of Product Formulations Within Each Concentration Range (%)							
		Unreported concentration	>50	>25-50	>10-25	>5-10	>1-5	>0.1-1	≤0.1
Paraffin									
Baby shampoos	1						1		
Eye shadow	127				1	19	18		89
Eye makeup remover	3					1	2		
Mascara	3						3		
Other eye makeup preparations	20			2	1	5	8	4	
Colognes and toilet waters	3				3				
Perfumes	30			30					
Sachets	6					6			
Other fragrance preparations	15			15					
Hair conditioners	14					14			
Hair rinses (noncoloring)	3						3		
Hair shampoos (noncoloring)	2							2	
Tonics, dressings, and other hair grooming aids	29				13	6	8	2	
Wave sets	1				1				
Other hair preparations (noncoloring)	3					1	2		
Other hair coloring preparations	3			1	2				
Blushers (all types)	65				1	10	28	26	
Face powders	15							15	
Makeup foundations	76					2	51	23	
Lipstick	439				18	54	290	76	1
Makeup bases	95					13	81	1	
Rouges	29				1	1	25	2	
Makeup fixatives	3						3		
Other makeup preparations (not eye)	44			3	11	13	16	1	
Cuticle softeners	3						3		
Nail creams and lotions	2				1		1		

COSMETIC INGREDIENT REVIEW

TABLE 11. (Continued.)

Product category	Total no. containing ingredient	No. of Product Formulations Within Each Concentration Range (%)							
		Unreported concentration	>50	>25-50	>10-25	>5-10	>1-5	>0.1-1	≤0.1
Hair conditioners	2	--	--	--	--	--	2	--	--
Tonics, dressings, and other hair grooming aids	14	--	--	4	2	2	8	--	--
Other hair preparations (noncoloring)	2	--	--	1	--	--	1	--	--
Other hair coloring preparations	1	--	--	--	--	--	1	--	--
Blushers (all types)	38	--	--	9	17	12	--	--	--
Face powders	1	--	--	--	--	1	--	--	--
Makeup foundations	9	--	--	--	6	2	2	1	--
Lipstick	640	--	--	13	121	382	124	--	--
Makeup bases	6	--	--	--	2	2	1	1	--
Other makeup preparations (not eye)	17	--	1	--	3	12	1	--	--
Cuticle softeners	1	--	--	--	1	--	--	--	--
Nail creams and lotions	1	--	--	--	--	1	--	--	--
Skin cleansing preparations (cold creams, lotions, liquids, and pads)	12	--	--	--	1	6	5	--	--
Face, body, and hand skin care preparations (excluding shaving preparations)	10	--	--	3	1	5	1	--	--
Moisturizing skin care preparations	8	--	--	--	3	3	1	--	--
Night skin care preparations	7	--	--	--	1	3	2	1	--
Other skin care preparations	4	--	--	--	2	2	--	--	--
1976 TOTALS	868	--	1	9	56	189	471	141	1
1979 TOTALS	899	13	1	4	67	138	591	84	1

Data from Refs. 16, 20.

Paraffin imparts stiffness to cosmetic preparations and forms the solid portion of stick cosmetics. It holds the oil phase of lipsticks and forms emulsions. It is also used as the wax component of depilatories.^(4,5,15,90)

Cosmetics containing Paraffin may be applied to the body surface from several times per day, as in fragrance and skin care preparations, to a few times per month, as in hair coloring preparations. It may remain on the body for a few hours in colognes or toilet waters, or for several days in hair conditioners.^(16,20)

The 1976, FDA data on cosmetic ingredients list Paraffin in 1,208 uses; in 1979 it had 945 uses. Its concentration of use ranges from less than 0.1% in some facial makeup, to up to 50% in eye and facial makeups and fragrance preparations (see Table 11).^(16,20)

BIOLOGICAL PROPERTIES

Animal Toxicology

Acute

Oral toxicity

Doses of 5 g/kg of a 75% and a 100% sample of Paraffin were not lethal to six and five rats, respectively.^(91,92)

A dose of 25 ml/kg of an undiluted, melted eye shadow formulation containing 5% Paraffin was administered to four adult beagle dogs. No deaths occurred and no abnormal behavior developed during the subsequent 14 days; no abnormalities were found at necropsy.⁽⁹³⁾ The same compound was given to 10 albino Wistar rats in 60 ml/kg doses. One animal died on Day 4. The remaining nine animals were normal and survived to the end of the experiment on Day 14.⁽⁹⁴⁾

Doses of 10 ml/kg of three eye shadow products containing 8% Paraffin, and a 5 ml/kg dose of a foot care cream containing 16% Paraffin were intubated into five albino rats per product. None of the 20 animals died in seven days⁽⁹⁵⁻⁹⁸⁾ (see Table 12).

Skin irritation

Raw Paraffin, solutions, and formulations containing Paraffin were patch tested on the clipped, intact skin of nine rabbits each (only six animals were used in one test). They were applied in 0.5 ml volumes by either open or closed patches for three repeated applications. The patches were removed after 24 h, and the sites were inspected immediately and at 72 h after the application. The results were scored according to the Draize method with a maximum PII score of 8.0. The results are shown in Table 13 and may be summarized as follows.

A sample of 100% Paraffin Wax was applied full strength under a single, closed patch to the skin of nine rabbits. No irritation developed.⁽⁹⁹⁾

Three samples of 50% Paraffin in petrolatum were tested in repeated, open patch applications to six rabbits. Two samples produced erythema in four animals that lasted three days,^(100,101) and one produced erythema in one rabbit that lasted two days.⁽⁹¹⁾

Four eye shadow cream formulations containing 8% Paraffin were tested using closed and open patches. The tests were repeated three times. The closed patches produced PIIs of 3.66 and 3.33 (potential for severe irritation), 2.33 (poten-

TABLE 12. Acute Oral Toxicity: Paraffin.

Wax conc. (%)	Dose of test mixture/kg	Solution or formulation	Species and number	LD ₅₀ /kg material	LD ₅₀ /kg Wax	Comments	Ref.
75	5 g	corn oil solution	6 rats	—	—	No deaths.	91
100	5 g	Pure	5 rats	> 5 g	> 5 g	—	92
5.0	25 ml	Formulation-eye shadow	4 beagle dogs			Single dose. 14-day observation period. No deaths or abnormalities. LD ₅₀ not reached.	93
5.0	60 ml	Formulation-eye shadow	10 albino Wistar rats			Single dose. 14-day observation period. One animal died on Day 4; findings included urinary staining of abdomen; intestines filled with red fluid. No other animals affected.	94
8.0	10 g	Formulation-eye cream	5 albino rats			No deaths	95
8.0	10 g	Formulation-eye cream	5 albino rats			No deaths	96
8.0	10 g	Formulation-eye cream	5 albino rats			No deaths	97
16.0	5 g	Formulation-foot care cream	5 albino rats			No deaths	98

ASSESSMENT: FOSSIL AND SYNTHETIC WAXES

TABLE 13. Acute Dermal Irritation/Toxicity: Paraffin.

Wax conc. (%)	Dose of test subst./kg	Petrolatum solution or formulation	No. of albino rabbits	Primary irrit. score		Contact time (h)	Observ. time (h)	Comments	Ref.
				Max. score	—				
<i>Irritation</i>									
50	0.5 ml	Solution	6	—	—	24	72	Erythema in 4 animals to Day 3	100
50	0.5 ml	Solution	6	—	—	24	72	Erythema in 4 animals to Day 3	101
50	0.5 ml	Solution	6	—	—	24	72	Erythema in 1 animal to Day 2	91
100	0.5 ml	Pure	9	0.0	0.0	24	72	No irritation	99

8.0	0.5 ml	Formulation-eye shadow	9	3.66/8.0	3.66/8.0	24	72	Severe irritation	95
8.0	0.5 ml	Formulation-eye shadow	9	3.33/8.0	3.33/8.0	24	72	Severe irritation	96
8.0	0.5 ml	Formulation-eye shadow	9	2.33/8.0	2.33/8.0	24	72	Moderate irritation	97
8.0	0.5 ml	Formulation-eye shadow	9	0.17/8.0	0.17/8.0	24	72	Minimal irritation	102
15	0.5 ml	Formulation-foot cream	9	0.61/8.0	0.61/8.0	24	72	Minimal irritation	103
16	0.5 ml	Formulation-foot cream	9	0.95/8.0	0.95/8.0	24	72	Minimal irritation	98
16	0.5 ml	Formulation-foot cream	9	1.06/8.0	1.06/8.0	24	72	Mild irritation	104
16	0.5 ml	Formulation-foot cream	9	1.22/8.0	1.22/8.0	24	72	Mild irritation	105
<i>Toxicity</i>									
50	4 ml	Solution	—	—	—	24	—	No systemic effects	91

tial for moderate irritation), and 0.17 (potential for minimal irritation).^(95-97,102) The open patch tests of three of these same products produced PIs of 0.0, 0.0, and 0.17.⁽⁹⁵⁻⁹⁷⁾

A single, open insult patch using a foot cream containing 15% Paraffin produced a PI of 1.61 (potential for minimal irritation).⁽¹⁰³⁾

A single, closed insult patch of three foot cream formulations containing 16% Paraffin caused PIs of 0.95 (potential for minimal irritation), 1.06 and 1.22 (potential for mild irritation).^(98,104,105)

Dermal toxicity

The acute dermal toxicity of 50% Paraffin in petrolatum was tested in rabbits (number not given). A dose of 4 ml/kg, applied under a closed patch for 24 h, produced no systemic effects, and no abnormalities were observed at necropsy⁽⁹¹⁾ (see Table 13).

Ocular irritation

Four 50% solutions of Paraffin in petrolatum were each instilled into the eyes of six albino rabbits with no rinse. Eyes were observed for irritation for three days. Two of the samples caused mild irritation in one rabbit on Day 1;^(91,101) the other samples were not irritating.^(99,100)

A 0.1 ml volume of an eye shadow product containing 5% Paraffin was instilled into the left eyes of six rhesus monkeys. Three treated eyes were washed with 20 ml warm tap water 30 sec after instillation. Observations up to 72 h after instillation showed no signs of irritation or corneal damage.⁽¹⁰⁶⁾

Another eye shadow formulation containing 5% Paraffin tested similarly caused no irritation or damage to the treated eyes of six monkeys.⁽¹⁰⁶⁾ A 0.1 ml volume of the sample was instilled into one eye of each of nine albino rabbits; in three, the treated eyes were washed 30 sec after instillation with 20 ml of deionized water. The eyes were inspected at 24, 48, and 72 h, and after four and seven days. Minimal conjunctival redness occurred in four of six animals with unwashed eyes 48 h after instillation. Two of three animals with irrigated eyes showed minimal conjunctival redness after 48 h.⁽¹⁰⁷⁾

Four eye shadow products containing 8% Paraffin were each instilled full strength, with no rinse, into the eyes of six albino rabbits. Eyes were observed for irritation for three days. Three formulations caused mild irritation at 24 h in one rabbit, and the fourth caused mild irritation after 48 h in one rabbit.^(95-97,102)

A foot cream containing 15% Paraffin, tested as above, caused mild irritation in three of six animals at Day 1.⁽¹⁰³⁾

Two foot care formulations with 16% Paraffin were also similarly tested, one causing mild irritation in one animal at 48 h,⁽⁹⁸⁾ and the other causing irritation in two rabbits at 24 h⁽¹⁰⁴⁾ (see Table 14).

Subchronic

Dermal toxicity

An eye shadow formulation containing 8% Paraffin was applied daily, five days per week for 13 consecutive weeks, to groups of female albino rats at "exaggerated dosage levels" of at least 100 times the normal human use concentration. The compound produced neither systemic toxic effects nor abnormal cumulative dermal effects.⁽¹⁰⁸⁾

ASSESSMENT: FOSSIL AND SYNTHETIC WAXES

71

TABLE 14. Ocular Irritation: Paraffin.

Wax conc. (%)	Dose	Solution/ formulation	Species and number	Observ. time (days)	Comments	Ref.
50	0.1 ml	Solution	6 albino rabbits	3	Mild irritation in 1 rabbit at 24 h	91
50	0.1 ml	Solution	6 albino rabbits	3	Mild irritation in 1 rabbit at 24 h	101
50	0.1 ml	Solution	6 albino rabbits	3	No irritation	99
50	0.1 ml	Solution	6 albino rabbits	3	No irritation	100
5	0.1 ml	Formulation- eye shadow	6 rhesus monkeys	7	No irritation	109
5	0.1 ml	Formulation- eye shadow	6 rhesus monkeys	7	No irritation	109
5	0.1 ml	Formulation- eye shadow	9 albino rabbits	7	Minimal conjunctival redness in 4/6 with unwashed eyes and 2/3 with washed eyes to 48 h	107
8	0.1 ml	Formulation- eye shadow	6 albino rabbits	3	Mild irritation in one rabbit at Day 1	95
8	0.1 ml	Formulation- eye shadow	6 albino rabbits	3	Mild irritation in one rabbit at Day 1	96
8	0.1 ml	Formulation- eye shadow	6 albino rabbits	3	Mild irritation in one rabbit at Day 1	97
8	0.1 ml	Formulation- eye shadow	6 albino rabbits	3	Mild irritation in one rabbit at 48 h	102
15	0.1 ml	Formulation- foot cream	6 albino rabbits	3	Mild irritation in 3/6 rabbits at Day 1	103
16	0.1 ml	Formulation- foot cream	6 albino rabbits	3	Mild irritation in 1 at 48 h	98
16	0.1 ml	Formulation- foot cream	6 albino rabbits	3	Mild irritation in 2 at 24 h	104

Special Studies

Carcinogenicity

Tumorigenicity: The tumorigenic properties of Paraffin and its intermediate products were studied in mice, rats, and rabbits. Paraffin, in distillate, filtrate, oil, raw, and purified forms, was applied repeatedly to the skin of the mice and rabbits and subcutaneously to the rats. After nine months, the distillate caused hyperkeratosis, followed by the development of benign papillomas in nearly all animals. The tumorigenic effects decreased according to the sequence of Paraffin products listed above, with pure Paraffin being practically without effect. The tumorigenic effect of raw petroleum is not attributable to Paraffin, but rather to its intermediate products.⁽¹¹⁰⁾

Five petroleum waxes were tested for carcinogenicity by repeated skin application of a 15% mixture in benzene, as well as in a year-long feeding study in rats. In neither test were the waxes positive for carcinogenic activity.⁽¹¹¹⁾

Bladder Cancer: Chapman et al.⁽¹¹²⁾ surgically implanted pellets of Paraffin waxes as artificial stones into rat bladder with prepared pouches and reported that the presence of urine was necessary for tumor induction. The investigators suggested that the pellets have a possible cocarcinogenic effect. They stressed, however, that the calculi growing around the pellets were the reason for the tumor enhancing effect. Studies by Ball et al.,⁽¹¹³⁾ Bonser et al.,^(114,115) Allen et al.,⁽¹¹⁶⁾ and Podilchak⁽¹¹⁷⁾ demonstrated also that foreign objects including Paraffin and glass beads may lead, by local irritation, to tumors in the urinary bladder of mice and rats. Clayson⁽¹¹⁸⁾ concluded that "if a foreign body in the presence of urine is all that is needed for tumor formation in the rat or mouse urothelium, chemicals provoking urinary calculus formation need careful scrutiny before being accepted as directly carcinogenic" (or tumorigenic).

Jull⁽¹¹⁹⁾ also studied the effects of Paraffin implants as carriers of carcinogenic chemicals in the mouse bladder. The author concluded that even though tumors may be produced by foreign bodies alone, the effect was not unique to Paraffin. These findings then are analogous to the induction of sarcomas by subcutaneous implants of various waxes and other inert materials.⁽¹¹¹⁾

Clinical Assessment of Safety

Intradermal effects

A mixture of liquid and solid Paraffin injected into the breasts for cosmetic purposes caused foreign body granuloma and calcification.⁽¹¹⁰⁾ Similar effects occurred after injection into the penis, scrotum, bridge of nose, cheeks, and eyelids.⁽¹²⁰⁾

Skin Irritation/Sensitization

The results of these tests are summarized in Table 15.

Single 24-hour patch test

The acute irritation to human skin by Paraffin and products containing Paraffin was tested by the patch test method. A single insult of the material was applied under occlusion for 24 h to the forearm or the upper back. Readings were taken immediately after patch removal.

TABLE 15. Clinical Data: Paraffin.

<i>TEST</i>						
<i>Wax conc. (%)</i>	<i>Pure or formulation</i>	<i>No. of subjects</i>	<i>Days on test</i>	<i>PII/Max. score</i>	<i>Comments</i>	<i>Ref.</i>
<i>24-h patch test</i>						
100	Pure	20	1	—	1/20 barely perceptible erythema	121
100	Pure	20	1	—	1/20 pink uniform erythema	122
8	Formulation- unspecified	18	1	0	No irritation	125
8	Formulation- unspecified	19	1	0	No irritation	123
8	Formulation- unspecified	20	1	0	No irritation	124
15	Formulation- foot cream	19	1	0	No irritation	126
16	Formulation- foot cream	17	1	0.15	1/17 had mild pink erythema	127
16	Formulation- foot cream	18	1	0.24	2/18 pink-red erythema	128
16	Formulation- foot cream	18	1	0.75	9/18 pink to pink-red erythema	129
16	Formulation- foot cream	10	1	0.35	—	130
<i>Repeat insult patch test</i>						
15	Formulation- smoothing cream	48	—	0	No reactions	131
<i>Maximization test</i>						
5	Formulation- eye shadow	25	—	0	No irritation or contact sensitization	132
5	Formulation- eye shadow	30	—	0	No irritation or contact sensitization	133
5	Formulation- eye shadow	29	—	0	No irritation or contact sensitization	134
<i>21-day cumulative irritancy test</i>						
5	Formulation- eye shadow	10	21	18/630	Essentially nonirritating	106
<i>In-use test</i>						
5	Formulation- eye shadow	187	14	0	No irritation from use	135

Two samples of 100% Paraffin were patch tested on 20 people per sample. The first caused barely perceptible erythema in one person. All others were negative.⁽¹²¹⁾ The second caused uniform erythema in one of 20 subjects.⁽¹²²⁾

Three formulations containing 8% Paraffin produced no irritation in panels of 18, 19, and 20 people.⁽¹²³⁻¹²⁵⁾

A formulation containing 15% Paraffin produced no irritation in 19 panelists.⁽¹²⁶⁾

Four products containing 6% Paraffin were tested. One formulation produced a mild erythema over the test sites in one of 17 subjects.⁽¹²⁷⁾ A second formulation caused erythema in 2 of 18 subjects.⁽¹²⁸⁾ The third formulation caused erythema in nine (PII = 0.75).⁽¹²⁹⁾ The fourth product had an irritation score of 0.35 out of a possible maximum score of 40.⁽¹³⁰⁾

Repeated insult patch test

A repeated insult patch test on a formulation containing 15% Paraffin was conducted on 48 panelists. The undiluted formulation was applied under occlusion to the skin of the back or the arm every other day for a total of 9–15 induction results. After a rest of 10–21 days, a challenge patch was applied to an adjacent, untreated site. Reactions were scored immediately after patch removal, and after 24, 48, and 72 hours this formulation caused no irritation and no sensitization in the 48 panelists.⁽¹³¹⁾

Maximization test

The contact sensitization of a formulation containing 5% Paraffin was tested on three different panels of 25, 30, and 39 people. The material was applied under occlusion to the same site on the volar forearm of all subjects for five 48 h periods. The patch sites were pretreated for 24 h with 2.5% aqueous sodium lauryl sulfate under occlusion. A challenge patch was applied after a 14-day rest and the sites were read on removal of the patch and 24 h thereafter. No irritation and no sensitization were observed.⁽¹³²⁻¹³⁴⁾

21-day cumulative irritancy test

A 21-day cumulative irritancy test of a formulation containing 5% Paraffin was conducted on 10 people. Patches containing the material were applied daily to the same site on the backs of each panelist for four consecutive days. The patches remained in contact with the skin for 23 h and scores were read just before the next patch application. There was a score of 18 out of a maximum possible irritation score of 630 indicating that the formulation was nonirritating.⁽¹⁰⁶⁾

In-use test

An in-use test of a product containing 5% Paraffin was conducted to determine irritancy on 187 women. After two weeks of daily use there was no irritation.⁽¹³⁵⁾

MICROCRYSTALLINE WAX: CHEMICAL AND PHYSICAL PROPERTIES

Structure/Composition

Microcrystalline Wax, like Paraffin, is a distillation product of crude petroleum; however, this wax is distinctly different from Paraffin.^(1,2) The name “micro-

crystalline" refers to the small, needle-like crystalline manifestations of the hydrocarbons in the wax. These crystals consist of long-chain, saturated hydrocarbons of high molecular weight. The molecules possess from 41 to 50 carbon atoms, with respective molecular weights of 580–700 mass units.^(1,2,6,90,136)

Properties

Microcrystalline Wax is a tough, flexible substance, with a high tensile strength and melting point, and a high penetration value and refractive index. It is adhesive (tacky), nonlustrous, somewhat greasy, plastic, and tends to flow under compression. This wax is compatible with other mineral waxes and with most vegetable waxes and resins.^(1,2,4-6,15,90,136) See Table 10 for specific properties.

Reactivity

No information was available.

Refining and Analytical Methods

Microcrystalline Wax is separated from the residue of crude petroleum left in the still after distillation and removal of the oil fraction containing Paraffin. The residue also contains heavy residual lubricating oil and asphalt. The asphalt is removed by treatment with concentrated sulfuric acid. The residue may then undergo one of several dewaxing processes, which involves dilution with organic solvents, chilling, and centrifugation to remove the wax. The oil is then removed by further organic solvent dilution, heating and filtering. The oil content of refined wax may vary from 2% to 12%.^(1,2,136)

The analytical methods used to determine the composition of Microcrystalline Wax include the following. Differential thermal analysis and molecular sieving procedures determine the total amount of normal paraffins in the wax and the solid hydrocarbons can be studied by differential thermal analysis; gas and gel permeation chromatography may also be used to determine the chain length of major constituents and the overall carbon chain number.⁽¹³⁷⁻¹³⁹⁾

Impurities

The natural hydrocarbon waxes may contain traces of benzopyrenes,⁽¹⁴⁰⁾ acids or alkalies, sulfur compounds, heavy metals, and arsenic.^(1,2,6)

USES

Noncosmetic Uses

Microcrystalline Wax is used in electrical insulation, waterproofing paper, box-board, textiles, leather and wood polish, and laminating paper and cloth, in rubber compounding, pattern making, binding for pipe covering, filter for packing, as a moulding wax base, and in miscellaneous polishes.^(1,2,6)

Purpose and Use in Cosmetics

The cosmetic uses of Microcrystalline Wax include facial and eye makeup preparations, fragrance products, coloring and noncoloring hair care preparations, manicuring products and skin care preparations.^(1,2,4-6,15,90)

In cosmetics, Microcrystalline Wax imparts firmness to makeup, fragrance products, hair grooming products, lipsticks, and solid stick-form deodorants.^(15,90)

Microcrystalline Wax may come into contact with the face, hands, and general body surface, around the eyes, scalp, and hair.^(16,20)

The 1976 FDA submission of cosmetic data reports Microcrystalline Wax in 868 formulations; the 1979 formulation data list 889 uses. It is used in concentrations of less than 0.1% in some preparations, and greater than 50% in others. The wax may be in contact with the body for several hours daily as in fragrance and makeup preparations, to several continuous days, as in hair conditioners^(16,20) (see Table 11).

BIOLOGICAL PROPERTIES

Animal Toxicology

Acute

Oral toxicity

A 20% Microcrystalline Wax-corn oil suspension was given by stomach tube to five groups of five albino rats. Each group received doses of either 0.464, 1.0, 2.15, 4.64, or 10.0 g/kg. After 14 days, the rats in the 0.464, 1.00, 2.15 and 4.64 g/kg dosage groups remained normal, but those in the 10 g/kg group showed progressive excessive salivation, decreased righting and placement reflexes, bloody discharges around the nose and mouth, unkempt hair and coats, and diarrhea; in the latter part of the observation period they were emaciated. Necropsy findings of the animals of the first four dosage groups were negative for abnormalities. Rats ingesting 10.0 g/kg showed congestion of lungs and kidneys, hyperemia of the large and small intestines, and solid wax in the stomach. No animals died from administration of the four lower dosage levels, but three died in the highest dose group. The LD₅₀ was estimated to be approximately 10 g/kg.⁽¹⁴¹⁾

The acute oral toxicity of a blusher formulation containing 4.35% Microcrystalline Wax was tested on five male and five female Sprague-Dawley rats. A 25 g/kg dose, administered by gavage, produced neither toxic effects nor deaths.⁽¹⁴²⁾

Skin irritation

A single 0.5 g application of 100% Microcrystalline Wax administered to the intact and abraded skin of six albino rabbits for 24 h caused slight erythema and edema in intact and abraded sites. The PII was 0.48 out of a possible maximum score of 8.0.⁽¹⁴³⁾

Three 0.5 g applications of a blusher product containing 4.35% Microcrystalline Wax were made at 24 h intervals under open patch conditions to intact and abraded skin of six New Zealand rabbits. After the third application, the intact sites of three animals and the abraded site of one animal had mild erythema. The product was considered to be practically nonirritating to the skin of rabbits.⁽¹⁴²⁾

Ocular irritation

A 0.1 g sample of 100% Microcrystalline Wax was applied to the left eye of each of six albino rabbits and the right eye was left untreated. After 24, 48, and

72 h, five of the six animals showed no irritation. One animal showed slight conjunctival erythema and edema after 24 h.⁽¹⁴⁴⁾

A 0.1 g volume of blush formulation containing 4.35% Microcrystalline Wax was instilled in one eye of each of six New Zealand white rabbits. After 72 h, all animals were free from irritation.⁽¹⁴²⁾

A 0.1 ml volume of a lipstick containing 15% Microcrystalline Wax was instilled into one eye of each of nine albino rabbits. Three of the treated eyes were washed with 20 ml of water 30 sec after treatment. After 24, 48, and 72 h and four and seven days, both washed and unwashed eyes had an irritation score of 3.3 out of a possible 110, indicating that the compound was a nonirritant.⁽¹⁴⁵⁾

Clinical Assessment of Safety

Skin Irritation/Sensitization

The results of these tests are summarized in Table 16.

Modified Draize–Shelanski–Jordan patch test

A blusher product containing 4.35% Microcrystalline Wax, when applied under occlusion to the skin of 205 men and women, caused mild erythema in three persons. The author concluded that the irritant reactions were insignificant.⁽¹⁴²⁾

21-day cumulative irritancy test

Microcrystalline Wax, tested in a 21-day cumulative irritancy test on eight subjects, produced no irritation.⁽¹⁴⁶⁾

A lipstick formulation containing 15% (0.3 ml) Microcrystalline Wax was applied under occlusion to the backs of 10 subjects. After 23 h, the patches were removed, the sites evaluated, and the next patch was applied. After testing for 21 days, the formulation produced an irritation score of 130 out of a possible 630. This score indicates a slight potential for mild cumulative irritation under these test conditions.⁽¹⁴⁷⁾

Maximization test

A lipstick formulation containing 15% (0.3 g) Microcrystalline Wax was applied under occlusion to the volar forearm surface of 25 subjects for five consecutive 48 h periods. After a rest of 10 days, an adjacent untreated site was pretreated with 10 percent sodium lauryl sulfate for 1 h, and then a patch of the test material was applied to the challenge site for 48 h under occlusion. This material caused no contact sensitization immediately after patch removal or 24 h later.⁽¹⁴⁸⁾

Phototoxicity test

A lipstick formulation containing 15% Microcrystalline Wax was applied under occlusion to the lower backs of 10 panelists for 24 h. The patches were then removed and the sites were irradiated for 12 min with filtered light from a Xenon Arc Solar Simulator emitting in the range of 320–400 nm. An untreated site was also irradiated as a control. After 24 and 48 h, both test and control sites showed minimal reactions. This product did not produce phototoxicity.⁽¹⁴⁹⁾

A similar test of a blusher formulation containing 4.35% Microcrystalline Wax produced no reaction in the 26 men and women tested.⁽¹⁴²⁾

TABLE 16. Clinical Data: Microcrystalline Wax.

TEST							
Wax conc. (%)	Dose	Pure or formulation	No. of subjects	Days on test	PII	Comments	Ref.
<i>Modified Draize-Shelanski-Jordan patch test</i>							
4.35	—	Formulation-blusher	205	—	1 + 1 and 2 + 2 reactions	1 + 1 reaction of mild, pink, uniform erythema 2 + 2 reactions of pink to red erythema	142
<i>21-day cumulative irritancy test</i>							
100	—	Pure	8	21	0.0/4	No irritation	146
15	0.3 ml	Formulation-lipstick	10	21	130/630	Slight potential for very mild cumulative irritation	147
<i>Maximization test</i>							
15	0.3 g	Formulation-lipstick	25	22	0/5	No instances of contact sensitization	148
<i>Phototoxicity test</i>							
4.35	—	Formulation-blusher	26	—	—	No reactions. Product exhibits no indication of phototoxicity or photoallergenicity	142
15	—	Formulation-lipstick	10	3	—	Minimal reaction. No greater than the reaction to light alone, not phototoxic	149
<i>In-use test</i>							
4.35	—	Formulation-blusher	26	—	—	Safe for intended use	142


In-use test

A blusher containing 4.35% Microcrystalline Wax caused no reactions in the 26 men and women using it.⁽¹⁴²⁾


SYNTHETIC WAXES EMULSIFYING WAX N.F.: CHEMICAL AND PHYSICAL PROPERTIES

Structure/Composition

Emulsifying Wax is wax manufactured from a mixture of cetyl and stearyl alcohol and polyoxyethylene derivatives of fatty acid esters of sorbitan:




Cetyl Alcohol



Stearyl Alcohol

3.



$R', R'', R''' =$ Fatty radicals

$n_1 + n_2 + n_3 + n_4 = N =$ total moles of ethylene oxide.

The method of manufacture is confidential.⁽¹⁵⁰⁾

Properties

The properties of Emulsifying Wax N.F. are listed in Table 17.^(150,151) This is a creamy-white solid with a characteristic odor, soluble in most hydrocarbon solvents, ether, chloroform and alcohol, and insoluble in water.⁽¹⁵¹⁾

Analytical Methods

The National Formulary⁽¹⁵¹⁾ lists the methods to determine the melting range, pH, and saponification and sodium value of Emulsifying Wax N.F. Thin-layer, column, and gas-liquid chromatography are used to characterize the various synthetic waxes.⁽¹⁵²⁾

TABLE 17. Properties: Nonnaturally Occurring Waxes.

Properties	Emulsifying Wax N.F.		Synthetic Wax		Synthetic Beeswax	
		Ref.		Ref.		Ref.
Melting range	48°–52°C	151	52°–74°C	153	60°–67°C	154
Congealing point			94°–98°C	153		
pH (at 2.9% w/w dispersion)	5.5–7.0	151				
Saponification value	14.0 Max.	151				
Iodine value	3.5 Max.	151			15 Max.	154
Hydroxyl value	178–192	151			25 Max.	154
Soluble in	Ether	151			Chloroform	154
	Chloroform				Volatile silicone	
	Alcohol				Fixed oils	
	Most hydrocarbons		Cold alcohol			

USE

Noncosmetic Uses

Emulsifying Wax N.F. is used as a pharmaceutical aid, an emulsifying agent and a stiffening agent.⁽¹⁵¹⁾

TABLE 18. Product Formulation Data.

Product category	Total no. containing ingredient	No. of product formulations within each concentration range (%)							
		Unreported concentration	>50	>25–50	>10–25	>5–10	>1–5	>0.1–1	≤0.1
<i>Emulsifying Waxes</i>									
Skin care preparations	12	—	—	—	—	7	3	2	—
1979 TOTALS	12	—	—	—	—	7	3	2	—
<i>Synthetic Wax</i>									
Blusher	1	—	—	—	—	—	1	—	—
Foundation	1	—	—	—	—	—	1	—	—
Lipstick	3	—	—	—	—	2	1	—	—
1981 TOTALS	5	—	—	—	—	2	3	—	—
<i>Synthetic Beeswax</i>									
Eye shadow	119	—	—	—	—	—	—	119	—
Blushers (all types)	14	—	—	—	—	—	—	14	—
Face powders	10	—	—	—	—	—	—	10	—
Makeup foundations	1	—	—	—	—	—	1	—	—
Rouges	2	—	—	—	—	—	2	—	—
1976 TOTALS	146	—	—	—	—	—	3	143	—
1979 TOTALS	108								

Data from Refs. 16, 20, 155.

Cosmetic Uses

Emulsifying Wax N.F. is an emulsifying and stiffening agent in cosmetics. The 1976 FDA data report that it has 12 uses, and the 1979 voluntary submission of data to the FDA reports two uses in concentrations of >0.1%–1%, three uses in concentrations of >1%–5%, and seven uses in concentrations of >5%–10%⁽²⁰⁾ (see Table 18).

BIOLOGICAL PROPERTIES

Animal Toxicology

Acute

Oral toxicity

The acute oral toxicity of Emulsifying Wax N.F. was studied in rats. Forty and 50% suspensions, and formulations containing 8%–8.5% were tested; the LD₅₀ of the wax was greater than 7.9 g/kg. Methods and results are presented in Table 19.

Ocular toxicity

The ocular irritation of Emulsifying Wax N.F. was studied in rabbits according to the Draize method. For concentrations of 100%, 5.0%, and 2.5% (raw ingredient), and 8%–8.5% for Emulsifying Wax N.F. formulations, only very mild irritation was reported. See Table 20 for results and methods.

Dermal irritation

Emulsifying Wax N.F. as a raw ingredient and in formulations was tested on rabbits by the Draize method for dermal irritation. Concentrations of 50% and 100% of one sample of an Emulsifying Wax N.F. were applied with and without occlusion. The mixtures of wax and water caused no irritation to very mild irritation (PII = 0.0–1.4). A different sample of Emulsifying Wax N.F. was tested in a similar manner at concentrations from 2.5% to 25% and produced only very mild irritation (PII = 0.3, 0.4, and 0.5). Irritation was reported for the formulations containing 8.0% wax with PIIs of 3.0 and 2.8. See Table 21 for methods and results.

Clinical Assessment of Safety

Skin Irritation/Sensitization

Patch testing

Patch tests of Emulsifying Wax N.F. were conducted on one solution of the raw ingredient and on two formulations. A concentration of 10% wax in peach kernel oil produced no irritation in 98 subjects.⁽¹⁵⁶⁾ A formulation containing 8.0%, and another 8.5% wax produced no irritation in the 101 and 100 subjects, respectively^(157,158) (see Table 22).

Repeated insult patch test

An Emulsifying Wax N.F. solution, 15% in avocado oil, was applied to 52 subjects for 48 h under occlusion, for a total of 10 applications. Sites were graded

TABLE 19. Acute Oral Toxicity: Emulsifying Wax N.F.

Conc. (%)	Dose/kg of test Solution	Pure/ Solution/ Formulation	Species and No.	Route	LD ₅₀ /kg	Comments	Ref.
25	12.5 g 15.6 g 18.75 g 21.85 g 40 ml	Solution in corn oil	CFW Carworth farm mice-10	Single dose-oral intubation	15.0 g	95% confidence limits of 12.0 to 18.75 g/kg of test solution.	159
25		Solution	Albino rats-10	Single oral dose	> 10 ml	25% solution of wax. LD ₅₀ of solution is > 40 ml/kg and of wax is > 10 ml/kg. Responses included slight depression in 5/10 animals for first 6 hrs after dosing. 14-day observation, 0/10 dead.	160
40	40 ml solution = 16 ml wax	40% suspension in corn oil	CFY rats-10	Single dose-oral intubation	> 16 ml	No deaths. Reactions after dosing included slight lethargy, decreased respiratory rate, hunched posture. Animals normal within 8 days. Body weights and autopsy normal.	161
50	15.8 g	50% suspension	Sprague-Dawley rats-5	NA ^a	> 7.9 g	50% suspension of wax in oil. LD ₅₀ of suspension is > 15.8 g/kg and of wax is > 7.9 g/kg.	162
8.0	14.7 g	Formulation-skin cream	Sprague-Dawley rats-5	NA		Dose of product was 14.7 g/kg. 8% of product is wax. LD ₅₀ was not reached.	163
8.0	7.0 ml	Formulation-moisture cream	Harlan-Wistar rats-10	Single dose by gavage		Fasted rats. No deaths or signs of toxicity were observed during the two-week study. 8% of dose is wax; LD ₅₀ was not reached.	164
8.5	14.7 g	Formulation-skin cream	Sprague-Dawley rats-5	NA		8.5% of dose is wax. No deaths.	165

^aNA = Not available.

ASSESSMENT: FOSSIL AND SYNTHETIC WAXES

83

TABLE 20. Ocular Toxicity: Emulsifying Wax N.F.

Conc. (%)	Dose	Pure or formulation	No. of rabbits	Comments	Ref.
2.5	0.1 ml	Solution	9	Draize scoring. No wash in 3 animals; wash after 30 sec in 3 rabbits; wash after 4 sec in 3 rabbits. Draize score of 0.0 in all rabbits at all times.	166
5.0	0.1 ml	Solution	9	Draize scoring. No wash in 3 animals; wash after 30 sec in 3 animals; wash after 4 sec on 3 animals. Draize score of 0.0 in all rabbits at all times.	167
100	0.1 ml = 59 mg	Pure	5	16 CFR 1500.42 Method. 14-day observation. Mild redness to Day 2 in 4/5 which persisted to Day 7 in 1 of these 4 rabbits. No irritation in 1/5. Negative result for eye irritation.	168
100	0.1 ml = 25 mg	Pure	6	21 CFR 191.1; 191.12 Method. Observed for 3 days. Compound produced very mild conjunctival irritation in 2/6 rabbits which cleared by Day 2.	169
100	0.1 ml	Pure	3	Mild conjunctival irritation which cleared by Day 2.	162
8.0	0.1 ml	Formulation-moisturizing cream	3	Draize Method. No irritation in cornea or iris. Minimal conjunctival irritation. Eye normal on Day 3.	170
8.0	0.1 ml	Formulation-moisturizing cream	6	Reactions scored after 1 h and after 1, 2, 3, and 7 days. Slight conjunctival hyperemia occurred in 4/6 animals. All normal in 24-48 h.	171
8.0	0.1 ml	Formulation-moisturizing cream	6	Reactions scored after 1 h and 1, 2, 3, and 7 days. Slight conjunctivitis in 4/6 animals 1 h after treatment. All normal at 24 h.	171
8.5	0.1 ml	Formulation-moisturizing cream	3	Draize Method. No irritation.	158

TABLE 21. Dermal Irritation: Emulsifying Wax N.F.

Conc. (%)	Dose	Solution/formulation	No. of albino rabbits	Route	PII	Comments	Ref.
2.5	0.5 ml	Water solution	6	Single application occlusion	0.3	Wax was melted and dissolved in water. Observation at 24 h showed mild redness. No irritation at 72 h.	172
5.0	0.5 ml	Water solution	6	Single application; abraded and intact	0.4	Occluded patch test of wax showed mild redness at 24 h. Wax melted and dissolved in water.	173
25	0.5 ml	Water solution	6	Single application; abraded and intact	0.5	Irritation scored after 24 and 48 h. Mild redness at 24 h. Wax melted and dissolved in water.	174
50	NA ^a	Solution	6	Clipped flanks of each animal	1.4	50% solution in corn oil under occlusion.	162
100	0.5 g in 0.5 ml water	Solution	6	Clipped intact and abraded skin of flank	<0.1	0.5 g wax mixed with 0.5 ml distilled water applied to clipped, abraded and intact skin. Very slight erythema observed in intact skin of 1/6 at 24 h only. No other irritation.	175
100	0.5 g in 0.5 ml water	Solution	6	Clipped intact and abraded skin of flank	0.0	0.5 g wax applied to each intact and abraded area under occlusion for 24 h. Skin evaluated after 24 and 48 h. No irritation.	176
100	0.5 g in distilled water	Solution	6	Clipped intact and abraded skin of flank	0.0	0.5 g wax applied to intact and abraded skin for 24 h. No irritation seen at observation times of 24 and 48 h.	177
8.0	0.5 ml	Formulation- moisturizing cream	3	Clipped flanks of each animal	2.8	4 daily applications. Irritation evaluated daily for 7 days. Slight erythema at 24 h. Erythema and edema at 3 days. Mild desquamation at 7 days.	171
8.0	0.5 ml	Formulation- moisturizing cream	3	Clipped flanks of each animal	3.0	7 daily applications. Irritation evaluated daily for 7 days. Erythema at 24 h. Erythema and edema at 3 days. Desquamation at 7 days.	171

^aNA = Not available.

ASSESSMENT: FOSSIL AND SYNTHETIC WAXES

85

TABLE 22. Clinical Data: Emulsifying Wax N.F.

TEST		Pure/ solution/ formulation	No. of subjects	Days on test	Irrit. Score		Comments	Ref.
Wax conc. (%)	Dose				Max. score			
<i>Patch test</i>								
10.0	NA ^a	Solution in peach kernel oil	98	—	0	No irritation		156
8.0	NA	Formulation- moisturizing lotion	101	—	0	No irritation		157
8.5	NA	Formulation- moisturizing cream	100	—	0	No irritation		158
<i>Repeated insult patch test</i>								
15	NA	Solution	52	—	—	Neither irritation or sensitization		162
100	0.5 g	Pure	50	—	0.0/8.0	Neither irritation or sensitization		178
8.0	0.2 ml	Formulation- moisturizer	10	20	2/630	"Mild Material"		179
8.0	NA	Formulation- moisturizer	110	34	NA	No primary irritation; no sensitization.		180
8.0	NA	Formulation- moisturizer	205	6 weeks	NA	One subject developed erythema and edema. Essentially nonirritating.		181
8.0	NA	Formulation moisturizer	108	—	—	Neither irritation or sensitization		182
8.0	NA	Formulation moisturizer	152	—	—	Neither irritation or sensitization		183
8.0	NA	Formulation moisturizer	189	—	—	Neither irritation or sensitization		184

^aNA = Not Available

immediately after patch removal. After a two-week nontreatment period, a challenge application caused neither primary skin irritation nor allergic sensitization.⁽¹⁶²⁾

A 0.5 g amount of undiluted Emulsifying Wax N.F. tested as above on 50 people scored 0.0 out of a maximum possible score of 8.0.⁽¹⁷⁸⁾

A moisturizing product containing 8% Emulsifying Wax N.F. was evaluated in 110 women. Patches impregnated with the material were affixed to the alcohol-cleansed backs for 10 repeated 48 h patches. Reactions were observed immediately after patch removal. A challenge was applied 14 days after the last patch was removed. No primary irritation occurred after the removal of the 48 h patch, and no sensitization occurred after the 14-day challenge.⁽¹⁸⁰⁾

Four other repeated insult patch tests were carried out on formulations containing 8.0% Emulsifying Wax N.F. One test used 231 subjects, only 205 of whom completed the study. Patches impregnated with a 0.1 g sample of the product were attached to the upper back or flexor arm for 24 h. The procedure was repeated nine times. A tenth and final 48 h patch was applied to the test site and to a fresh site, and inspected 48 and 72 h after application. Only one of the 205

people developed erythema and edema, but this subject had developed "irritation" from cosmetics in the past. No other reaction occurred.⁽¹⁸¹⁾

Three 8.0% wax formulations were tested with the modified Draize–Shelanski procedure on 108, 152, and 189 subjects, respectively. None of the three test products was a primary irritant or a contact sensitizer⁽¹⁸²⁻¹⁸⁴⁾ (see Table 22).

Cumulative irritation test

A cumulative irritation patch test was performed on each of six formulations containing 8.0% Emulsifying Wax N.F. One test used 12 subjects, 10 of whom completed the test. A 0.2 ml volume of the moisturizing lotion was applied under occlusion to the backs for 20 consecutive days, 23 h each day. The sites were scored 1 h after patch removal and a composite total score for the 10 panelists was 2 out of a maximum possible irritation score of 630.⁽¹⁷⁹⁾

SYNTHETIC WAX: CHEMICAL AND PHYSICAL PROPERTIES

Structure/Composition

Synthetic Wax (CAS number 8002-74-2), a blend of low molecular weight homopolymers of ethylene, has a molecular weight of 500–700 (C₄₀–C₅₀).⁽¹⁵³⁾

Synthetic Wax is sometimes referred to as Fisher–Tropch hydrocarbon wax or synthetic paraffin. It is produced by the catalytic reaction of hydrogen and carbon monoxide at high pressures (300–450 psi) and temperatures (230°–250°C). Homopolymerization of ethylene also produces Synthetic Wax.⁽¹⁵³⁾

Properties

The properties of Synthetic Wax are listed in Table 17. Although this is sometimes called synthetic paraffin, it is less transparent, has a finer crystalline structure, and produces a higher gloss than paraffin. It congeals at 94°–98°C, and melts between 52°C and 74°C.^(153,185)

Analytical Methods

Thin-layer, gas–liquid, and column chromatography are used to characterize Synthetic Wax.⁽¹⁵²⁾

USE

Noncosmetic Uses

Synthetic Wax is used as a melting point extender and a hardner for soft waxes.⁽¹⁸⁵⁾

Cosmetic Uses

Synthetic Wax imparts gloss and structure to cosmetics, and hardens soft waxes.^(185,186) It is reported to be used in 5 cosmetic formulations⁽¹⁵⁵⁾ (see Table 18).

BIOLOGICAL PROPERTIES

Animal Toxicology

Acute

Oral toxicity

A single 20 ml/kg dose of a lipstick formulation containing 6.0% Synthetic Wax (1.2 ml/kg wax) given by gavage to five male and five female Sprague–Dawley rats, produced no toxic effects.⁽¹⁸⁶⁾

Ocular irritation

The eyes of six New Zealand rabbits were instilled with a single 0.1 g amount of a lip formulation containing 6.0% Synthetic Wax. No irritation was observed during the next three days.⁽¹⁸⁶⁾

Skin irritation

A dose of 0.5 g of a lip formulation containing 6.0% Synthetic Wax caused mild to moderate irritation when applied to intact and abraded skin of six New Zealand white rabbits for three 24 h intervals.⁽¹⁸⁶⁾

Clinical Assessment of Safety

Skin Irritation/Sensitization

Patch test

In a modified Draize–Shelanski–Jordan patch test, 209 men and women were exposed to a lip conditioner containing 6% Synthetic Wax, under occlusion. There were no significant reactions, and the product was found to be a nonirritant and a nonsensitizer.⁽¹⁸⁶⁾

In-use test

A controlled use test of the lip product by 25 adult volunteers produced no reactions in four weeks.⁽¹⁸⁶⁾

SYNTHETIC BEESWAX: CHEMICAL AND PHYSICAL PROPERTIES

Structure/Composition

Synthetic Beeswax is a blend of fatty esters (C₃₂–C₆₂), fatty acids (C₁₆–C₃₆), fatty alcohols (C₁₆–C₃₆), and high molecular weight hydrocarbons (C₂₁–C₃₄). Esters are the most abundant, the hydrocarbons next, the acids, and then alcohols.⁽¹⁵⁴⁾

Properties

The properties of Synthetic Beeswax are listed in Table 17.

Analytical Methods

The analytical methods used to detect and characterize natural beeswax may also be used for Synthetic Beeswax. See "Beeswax, Analytical Methods," in "Waxes: Plant and Insect."⁽⁵²⁾

Cosmetic Use

Synthetic Beeswax is used in skin and hair care products to provide lubricity, emolliency, gloss, and film formation. It is used to provide rigidity of structure, gloss and feel, produce emulsions, and mold release characteristics⁽¹⁵⁴⁾ (see Table 18).

BIOLOGICAL PROPERTIES

Animal Toxicology

Acute

Oral toxicity

Dosage levels of 5.0, 7.12, 10.14, and 14.43 g/kg Synthetic Beeswax were each given to 10 fasted male Wistar rats. Toxicity and pharmacological effects were recorded 3–4 h later and once daily for 14 days. Isolated instances of chromorhinorrhea and chromodacryorrhea were reported for all groups, and diarrhea, ptosis, bulging eyes, and sniffing in the three lower doses. At the highest dose, one animal died on Day 1, and one on Day 6. No other deaths occurred.⁽¹⁸⁷⁾

Ocular irritation

The instillation of 0.1 ml of a 3.0% Synthetic Beeswax in mineral oil into the right eye of each of three albino rabbits showed no irritation after 1, 2, 3, 4, and 7 days.⁽¹⁸⁸⁾

A 0.1 ml volume of Synthetic Beeswax was instilled into the right eye of each of six New Zealand white rabbits. Reactions were recorded at 24, 48, and 72 h after dosing. Out of a possible irritation score of 110, this compound scored 6.3, 3, and 2 on Days 1, 2, and 3, respectively. It was minimally irritating on Days 1 and 2, and practically nonirritating on Day 3.⁽¹⁸⁹⁾

Dermal irritation

A patch impregnated with 5 g of Synthetic Beeswax in 1 ml of corn oil was applied to abraded and intact skin of six New Zealand white rabbits and left in place for 24 h. After 24 and 72 h, the Draize score was 2.08/8.0.⁽¹⁹⁰⁾

In another test, a 0.5 ml volume of Synthetic Beeswax was applied under occlusion to the intact and abraded skin of three albino rabbits. After 24 and 48 h, the Draize score (PII) was 0.0/8.0.⁽¹⁹¹⁾

Skin sensitization

A 50% solution of Synthetic Beeswax in distilled water with 1% Carboxymethyl Cellulose and 0.2% Tween 80 was applied to the clipped back of guinea pigs for three consecutive days, for three weeks, and once on the fourth week.

The first application was 0.05 ml, and the remaining nine, 0.1 ml. Fourteen days after the last application, the animals were challenged at a previously untreated site. Dermal reactions were scored 24 h after each treatment by the Draize Method. On a scale of 0–4, there was 0.16 erythema and 0.05 edema. The compound was neither sensitizing nor irritating.⁽¹⁹²⁾

Clinical Assessment of Safety

Skin Irritation/Sensitization

Repeated insult patch test

A lipstick formulation containing 7.2%–9.4% Synthetic Beeswax was assayed in a repeated insult patch test on 896 subjects. Formulation-impregnated patches were applied to the upper backs of the panelists three times per week, for 48 h, for three consecutive weeks. Sites were inspected before patch replacement on the next day. After 14 days, challenge patches were applied for 48 h to sites adjacent to the original patch site. None of the subjects showed irritation after 48 and 76 h.⁽¹⁹³⁾

Photosensitivity

A lipstick containing 7.2%–9.4% Synthetic Beeswax was applied under occlusion for 24 h to 83 subjects. The patches were then removed, the sites evaluated for irritation and then irradiated for 2–3 min with a Xenon Arc Solar Simulator (150 W) with a continuous emission in the UVA-UVB range (290–400 nm). Sites were inspected 48 h later, and the entire procedure was repeated for a total of six exposures. A challenge was applied 10 days later, and there were no reactions.⁽¹⁹³⁾

SUMMARY

Ozokerite, Ceresin, and Montan Wax are mineral waxes found in deposits of shale and coal.

Ozokerite is a wax found in areas of soft shale, and consists of an aliphatic series of straight-chain, branched chain, cyclic hydrocarbons and some oxygenated resinous bodies. The refined wax melts from 61.0°–88°C, and is soluble in benzene, turpentine and kerosene, and insoluble in ethyl and methyl alcohol and water. In cosmetics, it is used in baby products, eye and facial makeup preparations, hair, skin, nail, fragrance and suntan preparations, in concentrations of <0.1%–50%.

The acute oral toxicity of Ozokerite as a raw ingredient and in formulation was studied in mice, rabbits, and rats. Doses up to 200 mg/kg of a 2.0% concentration of the raw ingredient Ozokerite and 5 g/kg formulations containing up to 29% Ozokerite were nontoxic.

In primary irritation tests, eight samples of 50% Ozokerite in petrolatum applied to groups of nine animals produced erythema and edema in one to four rabbits per group. Formulations containing 4.5%, 5%, 13%, 28%, and 29% Ozokerite have, at most, a potential for mild irritation. A product containing 19.1% Ozokerite caused no irritation.

Five of eight samples of 50% Ozokerite in petrolatum tested for ocular irrita-

tion in rabbits produced no irritation and the remaining three were minimally irritating. Formulations containing 4.5%, 5%, 13%, 19.1%, 28%, and 29% Ozokerite caused mild or no irritation.

Human clinical studies include 24 h patch tests of Ozokerite alone and in formulation. Six samples of 100% Ozokerite were tested; one sample caused no irritation, and the other five caused barely perceptible to definite erythema in a small percentage of the test subjects. Formulations containing 5%, 13%, 28%, and 29% Ozokerite caused no irritation, but one formulation with 13% caused barely perceptible erythema in one person. Repeated insult patch tests of 4.5% Ozokerite and a contact allergy patch test of 13% Ozokerite in a formulation produced no reaction. A 21-day cumulative irritancy test of 13% Ozokerite in a formulation had an irritation score of 2 out of a possible total score of 756.

Ceresin is a microcrystalline mixture of complex hydrocarbons, which is produced from, and has chemical and physical properties similar to, Ozokerite.

A formulation containing 2% Ceresin given orally in a 5 g/kg dose did not produce toxic effects in rats. Dermal irritation tests in rabbits of the same product produced a potential for minimal irritation. Five other products, one containing 5%, and the other four containing 6% Ceresin, caused only mild irritation.

A 2.0% Ceresin concentration in a formulation produced no ocular irritation in rabbits. Mild irritation was produced by formulations containing 6.0% Ceresin Wax.

In clinical studies, one formulation containing 2.0% Ceresin produced virtually no reactions in both a Schwartz–Peck prophetic patch test and a Draize–Shelanski repeated insult patch test.

A formulation containing a mixture of 6% Beeswax and 6% Ceresin was tested in both animals and humans. No deaths were produced in rats given 80.0 ml/kg orally. Minimal ocular irritation was produced by the mixture in the nonirrigated rabbit eye after 24 h, but not in irrigated eyes. Clinical studies of the mixture in formulation include the prophetic patch test under UV light. The formulation caused neither irritation nor sensitization. A repeated insult patch test of the formulation caused neither irritation nor sensitization. Ultraviolet testing produced no reaction in the prophetic patch test areas. A 21-day cumulative irritancy test caused an irritancy score of 6.4 out of a possible total score of 630. No allergic or irritant reactions were caused by the formulation in a contact sensitization test and a two-week in-use test of the mixture caused no reactions.

Montan Wax, a bituminous wax extracted from lignites with volatile solvents, consists of monohydric alcohol esters, high molecular weight acids, and free alcohols. The wax melts between 72°C and 90°C, and is insoluble in water and soluble in carbon tetrachloride, benzene, and chloroform. Montan Wax is used in polishes, finishes, greases, waterproofing agents, rubber manufacture, and as a substitute for Carnauba Wax. In cosmetic formulations, Montan Wax is used in eye and facial makeup, fragrances, and skin care preparations at concentrations of 0.1%–25%.

Montan Wax when applied in 2 and 3 g/kg doses caused no irritation to intact rabbit skin. The single minimum toxic dose of the wax administered to mice by intraperitoneal injection was 7.5 g/kg. Animals fed 1 g/kg Montan Wax daily for four months had only an increase in the weights of the adrenal glands and kidneys. It caused no dermal toxicity to rabbits at a concentration of 200 g/kg, and inhalation for three months of an atmosphere containing 0.11–0.36 mg/l of wax caused no behavioral changes.

In repeated insult patch test clinical studies, lipsticks containing 1.61%, 1.93%, 1.81%, and 2.53% Montan Wax, were nonirritating and nonsensitizing.

Paraffin is a solid mixture of hydrocarbons, principally of the methane series, derived from high boiling fractions during the destructive distillation of petroleum. It consists of saturated and unsaturated hydrocarbon chains, naphthalenes and aromatics, and has both plate and needle crystalline shape. Refined Paraffin is a white, tasteless, odorless solid and melts between 43°C and 65°C. It is insoluble in water and cold alcohol, but soluble in most organic solvents.

Paraffin is used in candlemaking, lubrication, waterproofing, food, fruit and vegetable protection, pharmaceuticals, waxed paper, crayons, and polishes. In cosmetic ingredients, Paraffin is used in baby products, eye and facial makeup, and in fragrance, hair, nail, skin, shaving, and suntan preparations in concentrations of less than 0.1% to 50%.

In doses up to 5 g/kg, ingested Paraffin caused no toxic effects in rats. In tests with dogs and rats no toxic effects were produced by products containing 5%, 8%, and 16% Paraffin.

Three samples of 50% Paraffin in petrolatum caused erythema in rabbits, but an undiluted sample caused no irritation. Formulations containing 8%, 15%, and 16% Paraffin produced minimal to severe irritation. An acute dermal test of 50% Paraffin in petrolatum was negative for systemic toxicity.

Two 50% Paraffin solutions in petrolatum caused mild eye irritation in rabbits, but another two samples produced no irritation. Formulations containing 5%, 8%, 15%, and 16% Paraffin caused mild or no irritation in rabbit eyes.

Subchronic dermal toxicity studies of Paraffin in rats were negative for toxic effects.

The carcinogenic and cocarcinogenic effect of Paraffin was studied by placing implants into the urinary bladder of rodents. It was found that the wax implants acted as foreign bodies and caused local irritation which led to urinary calculus formation, and hence to bladder tumors.

In human studies, patch tests of two samples of undiluted wax caused minimal erythema in two of 40 subjects. Formulations containing 8%, 15%, and 16% Paraffin caused mild to no erythema. A repeated insult patch test of a formulation containing 15% Paraffin was negative for toxic reactions. In maximization tests, 5% Paraffin in a formulation caused no irritation or sensitization. A 21-day cumulative irritancy test, and an in-use test of 5% Paraffin in a product, were negative for irritation.

Microcrystalline Wax is a mixture of long chain, saturated hydrocarbons of high molecular weight. It is derived from the distillation residue of crude petroleum after the Paraffin Wax fraction is removed. Microcrystalline Wax is tough, nonlustrous, and greasy, with high tensile strength and a melting point of 63°C to 90°C.

Microcrystalline Wax is used in insulation, waterproofing, polishes, lamination, and rubber compounding. As a cosmetic ingredient, it is used in eye and facial makeup, and hair, nail, skin, and fragrance preparations in concentrations of less than 0.1% to greater than 50%.

In animal toxicology studies, 20% Microcrystalline Wax at doses of 0.464, 1.0, 2.15, and 4.64 g/kg were nontoxic to rats, but three of five rats in the group given doses of 10.0 g/kg, died. A formulation containing 4.35% Microcrystalline Wax was nontoxic when fed to rats.

In a skin irritation study, 100% Microcrystalline Wax produced slight

erythema and edema in intact and abraded rabbit skin. A formulation containing 4.35% wax caused mild erythema over the contact site.

In acute eye irritation studies, 100% Microcrystalline Wax produced slight irritation. Tests on products containing 4.25% and 15% were negative for ocular irritation.

In clinical studies using a Modified Draize–Shelanski–Jordon patch test, Microcrystalline Wax caused three mild reactions out of 205 individuals. A 21-day cumulative irritancy test of 100% wax and of a formulation containing 15% wax was negative for irritation and these materials had only a slight potential for mild cumulative irritation. In a maximization test, 15% wax in a product caused no reactions. Two formulations containing 4.35% and 15% Microcrystalline Wax caused no phototoxic reactions, and an in-use test of 4.35% Microcrystalline Wax in a product was negative for irritation.

Emulsifying Wax is a mixture of cetyl and stearyl alcohol and polyoxyethylene derivatives of fatty acid esters of sorbitan. It is a creamy-white solid with a characteristic odor. It is insoluble in water, soluble in hydrocarbon solvents, and has a melting-point range of 48°–52°C.

Emulsifying Wax is used in pharmaceuticals and cosmetics as an emulsifier and stiffener. In cosmetics, its concentrations range from 0.1% to 10%.

In acute oral toxicity studies, a 25% solution and 40% and 50% suspensions of Emulsifying Wax in oil produced no deaths in rats. Two products containing 8.0% Emulsifying Wax and one with 8.5% wax were likewise nontoxic. A 25% solution of Self-Emulsifying Wax was nontoxic.

In ocular irritation studies, three 100% samples of Emulsifying Wax caused mild to no irritation. Three formulations containing 8.0% Emulsifying Wax caused minimal irritation, while one containing 8.5% wax produced no irritation. Two solutions containing 2.5% and 5.0% Self-Emulsifying Wax were nonirritating.

A 50% solution of Emulsifying Wax in corn oil produced mild cutaneous irritation in rabbits. Two 100% samples of the wax produced no irritation, while a third produced minimal erythema in one of six rabbits. Solutions of 2.5%, 5.0%, and 25% Self-Emulsifying Wax produced minimal irritation.

In clinical studies using patch tests, 10% Emulsifying Wax in peach kernel oil and formulations containing 8.0% and 8.5% wax produced no irritation.

Repeated insult patch tests of a 15% solution and a 100% concentration of Emulsifying Wax were negative for irritation and sensitization. Of six products containing 8.0% Emulsifying Wax, one produced erythema and edema in one of 205 individuals, and the other five produced no irritation or sensitization.

Synthetic Wax is a blend of low molecular weight homopolymers of ethylene produced by the Fisher–Tropsh process. It melts between 52°C and 74°C, has a high gloss, and a fine crystalline structure. Synthetic Wax is used as a melting point extender and hardener for soft waxes. In cosmetics, it is used to create gloss, impart structure, and harden soft waxes.

In acute oral toxicity tests in rats, a product containing a 6% concentration of Synthetic Wax was nontoxic. This same product caused neither ocular nor skin irritation in rabbits. In clinical studies, a patch test and an in-use test of a product containing 6.0% Synthetic Wax were negative for adverse reactions.

Synthetic Beeswax is a blend of fatty esters, fatty acids, fatty alcohols, and high molecular weight hydrocarbons. Its properties and methods of analysis mimic the natural product. Synthetic Beeswax is used in cosmetics to produce emulsions, lubricity and gloss, and provides a structure for the formulation.

Synthetic Beeswax was tested for acute oral toxicity in rats. No animals died at doses up to 10.14 g/kg. In ocular irritation studies, a sample of pure Synthetic Beeswax was practically nonirritating, and a 3.0% solution in mineral oil solution caused no irritation. Pure Synthetic Beeswax, and a sample of wax dissolved in corn oil were nonirritating to rabbit skin. Neither irritation nor sensitization occurred when 50% Synthetic Beeswax was applied to guinea pig skin.

In a human clinical patch test study, a formulation containing 7.2%–9.4% Synthetic Beeswax produced no irritation and was not phototoxic.

CONCLUSION

The Panel concludes that on the basis of the available information presented in this report, Ozokerite, Ceresin, Montan Wax, Paraffin, Microcrystalline Wax, Emulsifying Wax N.F., Synthetic Wax and Synthetic Beeswax are safe as cosmetic ingredients in present practices of concentration and use.

ACKNOWLEDGMENT

Ms. Anne Moore, Scientific Analyst and writer, prepared the literature review and technical analysis used by the Expert Panel in developing this report.

REFERENCES

1. BENNET, H. (Ed.). (1956). *Commercial Waxes, 2nd ed.* New York, NY: Chemical Publishing Co.
2. WARTH, A.H. (1947). *The Chemistry and Technology of Waxes.* New York, NY: Reinhold Publishing Co.
3. KOSS, A.S. (1920). Determination of paraffin wax in ozokerite. *Przemysl Chem.* **4**, 74–6.
4. GREENBERG, L.A. and LESTER, D. (1954). *Handbook of Cosmetic Materials.* New York, NY: Interscience Publishers.
5. JAPAN COSMETIC INDUSTRY ASSOCIATION (JCIA). (1979). Tokyo, Japan: Yakuji Nippo.
6. HAWLEY, G.G. (Ed.). (1971). *The Condensed Chemical Dictionary, 8th ed.* New York, NY: Van Nostrand Reinhold Co.
7. COSMETIC, TOILETRY AND FRAGRANCE ASSOCIATION (CTFA). (Nov. 1980). Submission of data. Cosmetic ingredient chemical description—Ozokerite.*
8. YATSENKO, E.F. and DONTSOVA, G.M. (1963). A procedure for determining the chemical composition of wax and ozokerite. *Tr. Uchr. Nauchn.-Issled. Geol.-Razved. Inst.* **5**, 371–7.
9. KLAUSKE, M. (1971). Ozokerite (mountain wax) in Soviet medicine. *Dtsch. Med. J.* **22**(13), 426–9.
10. LESKOV, A.I., SELAVRI, T.V., GLADKIKH, A.S., LAKOZA, G.N., KURTEN, M.V., and TUKHTAEV, T. (1966). A pharmacological study of ozokerite. Information on the first inter-republic symposium on the experimental study of ozokerite. *Dusanlse* 32–36, 1965. From: *Ref. Zhodt Vypush Farmakol Kimioterap Sredstva Toksikol.*
11. LAVROVA, L.P., LYASKOVSKAYA, YU. N., SHISHKEVA, N.N., DYKLOP, V.K., IVENOVA, A.A., KALENOVA, M.S., DUBROVINA, L.I., and POLETAEV, T.M. (1955). Protective coatings for sausage products. *Trudy, Vsesoyuz. Nauch.-Issledovatel: Inst. Myasnoi Prom.* **7**, 48–67.
12. KUZMINOVA, M.A., KARIMOV, V.A., MAKSHSUMOV, M.N., and KHALIKOV, T.R. (1966). The antibacterial properties of the eastern drug ozokerite. *Communication 3. Tr Taskkent Form. Inst* **4**, 736–8.
13. TEZEKBAEV, S.D. (1956). Regeneration of skin after preliminary application of a chemical irritant. *Byull. Eskptl. Biol. i. Med.* **41**(5), 66–9.

*Available on request: Administrator, Cosmetic Ingredient Review, 1110 Vermont Ave., NW, Suite 810, Washington, DC 20005.

14. KHOIDAROV, K. KH., ISMAILOVA, M.B., and MORDVINTSEVA, O.V. (1966). Some information on the pharmacological activity of a preparation of ozokerite. Information on the first inter-republic symposium on the experimental study of ozokerite. Dushanbe. 71-72, 1965 From: Ref. Zh otd Vypush Formakol Khemioterap Sredstva Foxikol **5**(54), 414.
15. SENZEL, A.J. (Ed.). (1977). *Newburger's Manual of Cosmetic Analysis, 2nd ed.* Washington, DC: Association of Official Analytical Chemists.
16. FOOD AND DRUG ADMINISTRATION (FDA). (Aug. 31, 1976). Cosmetic product formulation data. FDA computer printout. Washington, DC.
17. CTFA. (Nov. 1980). Submission of data. Cosmetic ingredient chemical description—ceresin.*
18. CTFA. (Nov. 1980). Submission of data. Cosmetic ingredient chemical description—montan wax.*
19. CODE OF FEDERAL REGULATIONS. (1979). FDA, Approved Used of Paraffin, 21 CFR 172.615, 21 CFR 175.250, 21 CFR 175.300.
20. FDA. (June 20, 1979). Cosmetic product formulation data. FDA computer printout.
21. ROZANOVA, E.P. and SHTURM, L.D. (1965). A description of the microflora found in the Shor-Sou ozokerite beds. *Mikrobiologiya* **34**(5), 888-94.
22. ROZANOVA, E.P. and SHTURM, C.D. (1966). Changes in the composition of ozokerite due to the action of microorganisms. *Mikrobiologiya* **35**(1), 138-45.
23. SANGINOV, KH.S. (1966). The effect of different doses of ozokerite on the secretory activity of the stomach. Information on the first inter-republic symposium on the experimental study of ozokerite. Dushanbe. 47-51, 1965 Ref. Zh Otd Vypush Farmakol Khimioterap Sredstv toksikol. **5**(54), 413.
24. ISMAILOVA, V.N. and MAKLAKOVA, A.I. (1966). The problem of studying the drug ozokerite. *Tr. Toshkent Farm Inst.* **4**, 256-8.
25. CTFA. (Nov. 18, 1980). Submission of data. Unpublished cosmetic safety evaluation on a lipstick formulation containing 4.5 percent ozokerite.*
26. CTFA. (Nov. 18, 1980). Submission of data. Unpublished cosmetic safety evaluation on a coverstick formulation containing 19.1 percent ozokerite.*
27. CTFA. (Aug. 22, 1978). Submission of data. Unpublished toxicology summary report: animal tests on product containing 5 percent ozokerite.*
28. CTFA. (April 20, 1979). Submission of data. Unpublished toxicology summary report: animal tests on a product containing 5 percent ozokerite.*
29. CTFA. (Sept. 23, 1974). Submission of data. Unpublished biological evaluation summary report on a product containing 13 percent ozokerite.*
30. CTFA. (Sept. 23, 1974). Submission of data. Unpublished biological evaluation summary report on product containing 13 percent ozokerite.*
31. CTFA. (March 18, 1975). Submission of data. Unpublished biological evaluation summary report on two products containing 28 percent ozokerite and 29 percent ozokerite, respectively.*
32. CTFA. (Sept. 19, 1972). Submission of data. Unpublished safety data on raw ingredient ozokerite.*
33. CTFA. (Sept. 19, 1972). Submission of data. Unpublished safety data on raw ingredient ozokerite.*
34. CTFA. (Sept. 19, 1972). Submission of data. Unpublished safety data on raw ingredient ozokerite.*
35. CTFA. (Sept. 19, 1972). Submission of data. Unpublished safety data on raw ingredient ozokerite.*
36. CTFA. (Sept. 19, 1972). Submission of data. Unpublished safety data on raw ingredient ozokerite.*
37. CTFA. (Sept. 19, 1972). Submission of data. Unpublished safety data on raw ingredient ozokerite.*
38. CTFA. (Sept. 19, 1972). Submission of data. Unpublished safety data on raw ingredient ozokerite.*
39. CTFA. (Sept. 19, 1972). Submission of data. Unpublished safety data on raw ingredient ozokerite.*
40. CTFA. (Sept. 13, 1972). Submission of data. Unpublished human single insult (24-hr) occlusion patch test on ozokerite.*
41. CTFA. (Sept. 13, 1972). Submission of data. Unpublished human single insult (24-hr) occlusion patch test on ozokerite.*
42. CTFA. (March 10, 1978). Submission of data. Clinical evaluation report: human patch test on a product containing 5 percent ozokerite.*
43. CTFA. (July 24, 1978). Submission of data. Unpublished clinical evaluation report: human patch test of a product containing 5 percent ozokerite.*
44. CTFA. (July 14, 1978). Submission of data. Unpublished clinical evaluation: human patch test of product containing 5 percent ozokerite.*
45. CTFA. (July 24, 1978). Submission of data. Unpublished clinical evaluation: human patch test of a product containing 5 percent ozokerite.*
46. CTFA. (August 15, 1978). Submission of data. Unpublished clinical evaluation report: human patch test on a product containing 5 percent ozokerite.*
47. CTFA. (May 1, 1974). Submission of data. Unpublished clinical evaluation report: human patch test on product containing 13 percent ozokerite.*

48. CTFA. (Sept. 6, 1974). Submission of data. Unpublished clinical evaluation report: human patch test on a product containing 13 percent ozokerite.*
49. CTFA. (March 13, 1975). Submission of data. Unpublished clinical evaluation report: human patch test of two products—28 percent ozokerite and 29 percent ozokerite.*
50. CTFA. (Sept. 30, 1974). Submission of data. Unpublished repeated insult patch test of a formulation containing 13 percent ozokerite.*
51. HILL TOP RESEARCH LABS (HTRL). (Aug. 13, 1976). Submission of data by CTFA. Unpublished human 21-day cumulative irritancy test on product containing 13 percent ozokerite.*
52. COSMETIC INGREDIENT REVIEW (CIR). (1982). Waxes: plant and insect.
53. CTFA. (Dec. 19, 1979). Submission of data. Summary of acute oral and dermal toxicity, primary skin and ocular irritation and human patch testing of a lipstick containing 2 percent Ceresin Wax.*
54. CTFA. (Nov. 24, 1980). Submission of data. Unpublished acute skin irritancy of product containing 5 percent ceresin.*
55. CTFA. (Nov. 24, 1980). Submission of data. Unpublished acute skin irritancy on product containing 6 percent ceresin.*
56. CTFA. (Nov. 24, 1980). Submission of data. Unpublished acute skin irritancy of product containing 6 percent ceresin.*
57. CTFA. (Nov. 24, 1980). Submission of data. Unpublished acute skin irritancy on a product containing 6 percent ceresin wax.*
58. CTFA. (Nov. 24, 1980). Submission of data. Unpublished acute skin irritancy of product containing 6 percent ceresin.*
59. CTFA. (Nov. 24, 1980). Submission of data. Unpublished acute eye irritancy of product containing 6 percent ceresin.*
60. CTFA. (Nov. 24, 1980). Submission of data. Unpublished acute eye irritancy test of a product containing 6 percent ceresin.*
61. CTFA. (Nov. 24, 1980). Submission of data. Unpublished acute eye irritancy test of product containing 6 percent ceresin.*
62. CTFA. (1973). Submission of safety data on irritation and sensitization tests on a product containing 2 percent Ceresin.*
63. BIODYNAMICS. (Sept. 4, 1975). Submission of data by CTFA. Unpublished acute oral toxicity in rats of product containing 6 percent ceresin and 6 percent beeswax.*
64. BIODYNAMICS. (August 5, 1975). Submission of data by CTFA. Unpublished rabbit eye irritation of product containing ceresin wax (6.0 percent) and beeswax (6.0 percent).*
65. RESEARCH TESTING LABS (RTL). (Jan. 12, 1976). Submission of data by CTFA. Unpublished human subject patch study on a product containing 6.0 percent ceresin wax and 6.0 percent beeswax.*
66. HTRL. (Aug. 21, 1975). Submission of data by CTFA. Unpublished study of cumulative irritant properties of a product containing 6.0 percent ceresin wax and 6.0 percent beeswax.*
67. CTFA. (Sept. 23, 1975). Submission of data. Unpublished contact sensitization test on product containing 6.0 percent ceresin wax and 6.0 percent beeswax.*
68. CTFA. (May 27, 1975). Submission of data. Unpublished usage test of product containing 6.0 percent ceresin wax and 6.0 percent beeswax.*
69. MARCUSSON, J. and MELKUS, H. (1917). Montan wax and its behavior on distillation. *Dhim. Ztg* **41**, 129–36 (1917) through *J. Soc. Chem. Ind.* **36**, 540.
70. MEYERHEIM, G. (1919). Montan wax. *Seifenfabr* **39**, 365–7, 394–6.
71. HUC, P. (1922). Montan wax and cream polishes. *Industries du cuir*. pp. 217–21.
72. FUCHS, W. and DIEBERG, R. (1956). Solution of the montanic acid problem by chromatography and infrared spectroscopy. *Fette, Seifen, Anstrichm.* **58**, 826–31.
73. TATE, K.R. and WELLS, N. (1975). Application of differential thermal analysis to peat waxes. Part 2. Montan Wax and New Zealand peat waxes. *NZJ Sci.* **18**(3), 483–91.
74. SCHMIDT, H. (1972). Thin layer chromatographic identification of genuine natural waxes. *Am. Cosmet. Perfum.* **87**(10), 35–9.
75. POPL, M. and HAVEL, Z. (1975). Chromatographic separation of montan wax and identification of individual fractions by gas chromatography and mass spectrometry. *Fette, Seifen, Anstrichm.* **77**(2) 51–4.
76. LUDWIG, J.F. (1966). Modern instrumental wax analysis. *Soap Chem. Specialties* **43**(3), 70.
77. ASANO, S. (1976). Gas-liquid chromatography of montan wax and its derivatives. *Kanzei Chuo Bunsakishoko* **16**, 49–55.
78. MARCUSSON, J. (1924). Preparation of ceresin hydrocarbons from montan wax. *Z. Angew. Chem.* **37**, 413–4.
79. STATSEK, N.K. (1963). Toxicological and hygienic characteristics of lignite wax. *Gigiena i Piziol. Truda, Proizo, Toksikal, Klinika Prof. Zabolevanii.* pp. 103–5.

80. UNIVERSITY OF CALIFORNIA AT LOS ANGELES (UCLA). (September 23, 1976). Submission of data by CTFA. Modified Draize–Shelanski–Jordan patch test of a lipstick containing 1.61 percent Montan Wax.*
81. UCLA. (December 28, 1978). Submission of data by CTFA. Modified Draize–Shelanski–Jordan patch test of a lipstick containing 1.93 percent Montan Wax.*
82. UCLA. (April 26, 1979). Submission of data by CTFA. Modified Draize–Shelanski–Jordan patch test of a lipstick containing 1.81 percent Montan Wax.*
83. UCLA. (February 29, 1980). Submission of data by CTFA. Modified Draize–Shelanski–Jordan patch test of a lipstick containing 1.81 percent Montan Wax.*
84. FAUST, H.R. (1978). The thermal analysis of waxes and petrolatums. *Thermochim. Acta.* **26**(1–3), 383–98.
85. CTFA. (Nov., 1980). Submission of data. Cosmetic ingredient chemical description—paraffin wax.*
86. MAZEE, W.M., GERSMANN, H.R., and VAN DER WIEL, A. (1966). Analysis of polycyclic aromatic hydrocarbons in petrolatum waxes and white mineral oils with appraisal of merits of different methods of analysis. *Food Cosmet. Toxicol.* **4**(1), 17–33.
87. DOBIES, R.S. (1968). Thin-layer chromatographic method for determining ultraviolet absorbers in paraffin wax. *J. Chromatogr.* **35**(3), 370–5.
88. ADLARD, E.R. and WHITMEN, B.T. (1958). Application of high temperature gas–liquid chromatography in the petroleum industry. *Gas Chromatog.* pp. 351–67.
89. SCHWARTZ, R.D., MATHEWS, R.G., and PEDRO, N.A. (1977). High-temperature gas–liquid chromatography with a dendritic salt support. *J. Chromatogr.* **142**, 103–7.
90. BALSAM, M.S. and SAGARIN, E. (Eds.). (1972). *Cosmetic: Science and Technology*. New York, NY: Wiley-Interscience.
91. CTFA. (Sept. 14, 1972). Submission of data. Unpublished safety data on raw ingredient paraffin wax.*
92. CTFA. (Oct. 30, 1975). Submission of data. Unpublished biological evaluation summary report on paraffin wax.*
93. BIODYNAMICS. (Aug. 8, 1975). Submission of data by CTFA. Unpublished acute oral toxicity in dogs of product containing 5 percent paraffin wax.*
94. BIODYNAMICS. (August 1, 1975). Submission of data by CTFA. Unpublished acute oral toxicity in rats of a product containing 5 percent paraffin wax.*
95. CTFA. (March 1, 1977). Submission of data. Unpublished biological evaluation summary report on product containing 8 percent paraffin wax.*
96. CTFA. (March 24, 1977). Submission of data. Unpublished biological evaluation summary report on product containing 8 percent paraffin.*
97. CTFA. (Aug. 1, 1977). Submission of data. Unpublished biological summary report on product containing 8 percent paraffin.*
98. CTFA. (Aug. 28, 1974). Submission of data. Unpublished biological evaluation summary report on product containing 16 percent paraffin.*
99. CTFA. (Sept. 18, 1980). Submission of data. Unpublished toxicology summary report: Paraffin animal tests.*
100. CTFA. (May 24, 1972). Submission of data. Unpublished safety report on raw ingredient paraffin wax.*
101. CTFA. (May 24, 1972). Submission of data. Unpublished safety data on raw ingredient paraffin wax.*
102. CTFA. (Oct. 5, 1977). Submission of data. Unpublished biological evaluation summary report on product containing 8 percent paraffin wax.*
103. CTFA. (April 3, 1980). Submission of data. Unpublished toxicology summary report: animal tests on product containing 15 percent paraffin.*
104. CTFA. (Feb. 8, 1980). Submission of data. Unpublished toxicology summary report: animal tests on product containing 16 percent paraffin wax.*
105. CTFA. (Dec. 27, 1979). Submission of data. Unpublished toxicology summary report: animal tests on 16 percent paraffin wax.*
106. HTRL. (Aug. 19, 1975). Submission of data by CTFA. Unpublished study of cumulative irritant properties of a series of test materials. Product containing 5 percent paraffin.*
107. BIODYNAMICS. (July 16, 1975). Submission of data by CTFA. Unpublished safety data on rabbit eye irritation of product containing 5 percent paraffin wax.*
108. CTFA. (May 30, 1978). Submission of data. Unpublished subchronic dermal toxicity safety data on cosmetic formulation containing 8 percent paraffin wax.*
109. HAZELTON LABORATORIES OF AMERICA. (Aug. 15, 1975). Submission of data by CTFA. Unpublished safety data. Eye irritation study in monkeys of product containing 5 percent paraffin wax.*
110. GETMANETS, I. YA. (1966). Blastomagenic properties of paraffin and its intermediate products. *Gigiena Truda i Prof. Zabollevaniya.* **10**(5), 58–9.
111. SHUBIK, KP., SAFFIOTTI, D., LIJINSKY, W., PIETRA, G., RAPPORT, H., TOTH, B., RAHA, C.R.,

- TOMATIS, L., FELDMAN, R., and RAMAHI, H. (Nov. 1962). Studies on the toxicity of petroleum waxes. *Tox. Appl. Pharm.* Vol 4. Supplement.
112. CHAPMAN, W.H., KIRCHHEIM, D., and McROBERTS, J.W. (1973). Effect of the urine and calculus formation on the incidence of bladder tumors in rats implanted with paraffin wax pellets. *Cancer Res.* **33**(6), 1225-9.
113. BALL, J., FIELD, W.E., ROE, F.J.E., and WALTERS, M. (1964). The carcinogenic and co-carcinogenic effects of paraffin wax pellets and glass beads in the mouse bladder. *Brit. J. Urol.* **36**(2), 225-37.
114. BONSER, G.M., BRADSHAW, L., CLAYSON, D.B., and JULL, J.W. (1956). *Brit. J. Cancer* **10**, 539.
115. BONSER, G.M., BOYLAND, E., BUSBY, E.R., CLAYSON, D.B., GROVER, P.L., and JULL, J.W. (1963). *Brit. J. Cancer* **17**, 127.
116. ALLEN, M.J., BOYLAND, E., DUKES, C.E., HORNING, E.S., and WATSON, J.D. (1957). *Brit. J. Cancer* **11**, 212.
117. PODILCHAK, M.D. (1957). *Urologia*, 6:14 (Abstracted in *Excerpta Medica*, Section XVI, 1959, 7, No. 2773).
118. CLAYSON, D.B. (1974). Guest editorial. Bladder carcinogenesis in rats and mice. Possibility of artifacts. *J. Natl. Cancer Inst.* **52**, 1685-9.
119. JULL, J.W. (1975). Carcinogenesis by chemicals implanted into the bladder: an evaluation. *Gann Monogr. Cancer Res.* **17**, 383-92.
120. BENNETT, D.H. and WADE, J.S. (1969). Rectal paraffinoma. *Proc. Roy. Soc. Med.* **62**(8), 818.
121. CTFA. (Sept. 12, 1972). Submission of data. Unpublished primary skin irritation test in humans.*
122. CTFA. (April 12, 1972). Submission of data. Unpublished primary skin irritation test of paraffin in humans.*
123. CTFA. (Jan. 20, 1977). Submission of data. Unpublished clinical evaluation report: human patch test of product containing 8 percent paraffin.*
124. CTFA. (Feb. 18, 1977). Submission of data. Unpublished clinical evaluation report: human patch test of a product containing 8 percent paraffin wax.*
125. CTFA. (Dec. 12, 1977). Submission of data. Unpublished clinical evaluation report: human patch test on product containing 8 percent paraffin.*
126. CTFA. (Feb. 13, 1981). Submission of data. Unpublished clinical evaluation report: human patch test of a product containing 15 percent paraffin wax. (Date of CTFA Transmittal letter).*
127. CTFA. (April 15, 1974). Submission of data. Unpublished clinical evaluation report: human patch test of product containing 16 percent paraffin.*
128. CTFA. (Feb. 7, 1980). Submission of data. Unpublished clinical evaluation report: human patch test on product containing 16 percent paraffin.*
129. CTFA. (Feb. 13, 1981). Submission of data. Unpublished clinical evaluation report: human patch test on product containing 16 percent paraffin. (Date of transmittal letter from CTFA).*
130. CTFA. (Feb. 13, 1981). Submission of data. Unpublished clinical evaluation report: human patch test on formulation containing 16 percent paraffin. (Date of transmittal letter from CTFA).*
131. CTFA. (Aug. 29, 1980). Submission of data. Unpublished contact allergy test of a product containing 15 percent paraffin.*
132. CTFA. (Sept. 8, 1975). Submission of data. Unpublished contact sensitization test of a product containing 5 percent paraffin.*
133. CTFA. (Sept. 28, 1975). Submission of data. Unpublished contact sensitization test of a product containing 5.0 percent paraffin wax.*
134. CTFA. (Sept. 23, 1975). Submission of data. Unpublished contact sensitization test of a product containing 5 percent paraffin wax.*
135. CTFA. (Sept. 5, 1975). Submission of data. Unpublished usage test of product containing 5 percent paraffin wax.*
136. CTFA. (Nov. 1980). Submission of data. Cosmetic ingredient chemical description—microcrystalline wax.*
137. HILLMAN, D.E. (1971). Characterization and analysis of waxes by gel permeation chromatography. *Anal. Chem.* **43**(Jul), 1007-13.
138. WASHALL, T.A. (1970). Improved molecular sieving procedure for the determination of total normal paraffins in microcrystalline wax. *J. Chromatogr. Sci.* **8**(11), 663-7.
139. ANISIMOV, I.G., MITROFANOV, M.G., SOLETSKII, O.I., FEDOROVA, Z.V., and MELIKHORA, A.V. (1959). Solid hydrocarbons studied by differential thermal analysis. *Khim. Tecknol. Topl. Masel.* **14**(6), 54-7.
140. FRITZ, W. and WOGGON, H. (1978). The content of benzo (a) pyrene in paraffins and microcrystalline waxes. *X. Gesamte Hyg.* **24**(9), 697-9.
141. CTFA. (July 16, 1969). Submission of data. Unpublished safety data on cosmetic ingredient microcrys-

- talline wax.*
142. CTFA. (Nov. 18, 1980). Submission of data. Unpublished cosmetic safety evaluation on a blusher formulation containing 4.35 percent microcrystalline wax.*
 143. CTFA. (July 16, 1969). Submission of data. Unpublished acute toxicity and irritation studies on microcrystalline wax.*
 144. CTFA. (July 16, 1969). Submission of data. Unpublished ocular safety data on cosmetic ingredient microcrystalline wax.*
 145. STILLMEADOW. (Oct. 10, 1979). Submission of data by CTFA. Unpublished safety data on cosmetic formulation (lipstick). Rabbit eye irritation.*
 146. CTFA. (Feb. 13, 1981). Submission of data. Unpublished safety data on cosmetic ingredient microcrystalline wax. (Date of transmittal letter from CTFA).*
 147. HTRL. (December 10, 1979). Submission of data. Unpublished safety data on cosmetic formulation (lipstick) containing 15 percent microcrystalline wax.*
 148. CTFA. (January 3, 1980). Submission of data. Unpublished safety data on cosmetic formulation (lipstick) containing 15 percent microcrystalline wax.*
 149. TEST KIT LABORATORIES (TKL). (Feb. 5, 1980). Submission of data by CTFA. Unpublished safety data on cosmetic formulation (lipstick) containing 15 percent microcrystalline wax.*
 150. CTFA. (Nov. 1979). Submission of data. Unpublished cosmetic ingredient chemical description of emulsifying wax.*
 151. NATIONAL FORMULARY (N.F.) (1975). Fourteenth Edition (N.F. XIV), American Pharmaceutical Association, Washington, DC.
 152. ABATE, V., BADOUX, V., HICKS, S.Z., and MESSINGER, M. (1970). Characteristics of natural and synthetic waxes using combined chromatographic techniques. *J. Soc. Cosmet. Chem.* **21**(4), 119-28.
 153. CTFA. (Nov. 1980). Submission of data. Cosmetic ingredient chemical description on synthetic wax.*
 154. WICKHEN PRODUCTS. (Feb. 13, 1981). Submission of data by CTFA. Tentative specification, typical properties, functions and uses of waxenal 820 S.B. and 821 S.B. synthetic beeswax. (Date of CTFA transmittal letter).*
 155. FDA. (Dec. 5, 1981). Cosmetic product formulation data. FDA computer printout.
 156. CTFA. (Jan. 14, 1969). Submission of data. Unpublished human testing on emulsifying wax.*
 157. CTFA. (March 3, 1978). Submission of data. Unpublished human patch test of product containing 8 percent emulsifying wax.*
 158. CTFA. (June 11, 1976). Submission of data. Unpublished acute eye irritation of product containing 8.5 percent emulsifying wax.*
 159. LEBERCO LABS. (Oct. 23, 1973). Submission of data by CTFA. Unpublished acute oral toxicity in mice of emulsifying wax.*
 160. CONSUMER PRODUCT TESTING CO. (CPTC). (Sept. 10, 1976). Submission of data by CTFA. Unpublished acute oral toxicity study of self-emulsifying wax in rats.*
 161. HUNTINGDON RESEARCH CENTER. (Sept. 9, 1977). Submission of data by CTFA. Unpublished Acute Oral Toxicity in Rats of Emulsifying Wax.*
 162. CTFA. (Sept. 17, 1976). Submission of data. Unpublished safety data on emulsifying wax.*
 163. CTFA. (Feb. 5, 1976). Submission of data. Unpublished acute oral toxicity of a product containing 8 percent emulsifying wax.*
 164. CTFA. (April 8, 1980). Submission of data. Unpublished acute oral toxicity testing of a product containing 8 percent emulsifying wax.*
 165. CTFA. (Jan. 4, 1978). Submission of data. Unpublished acute oral toxicity of a product containing 8.5 percent emulsifying wax.*
 166. CPTC. (Sept. 10, 1976). Submission of data by CTFA. Unpublished ocular irritation of self-emulsifying wax.*
 167. CPTC. (Sept. 10, 1976). Submission of data by CTFA. Unpublished ocular irritation of product containing 2.5 percent self-emulsifying wax.*
 168. HUNTINGDON RESEARCH CENTER. (April 28, 1977). Submission of data by CTFA. Unpublished acute eye irritation of emulsifying wax.*
 169. LEBERCO LABS. (July 7, 1973). Submission of data by CTFA. Unpublished acute eye irritation of emulsifying wax.*
 170. CTFA. (Jan. 20, 1976). Submission of data. Unpublished acute eye irritation of a product containing 8 percent emulsifying wax.*
 171. CTFA. (Sept. 11, 1979). Submission of data. Unpublished acute dermal and ocular irritation of product containing 8 percent emulsifying wax.*
 172. CPTC. (Sept. 10, 1976). Submission of data by CTFA. Unpublished primary dermal irritation of self-emulsifying wax.*

ASSESSMENT: FOSSIL AND SYNTHETIC WAXES

99

173. CPTC. (Sept. 10, 1976). Submission of data by CTFA. Unpublished primary dermal irritation of self-emulsifying wax.*
174. CPTC. (Sept. 10, 1976). Submission of data by CTFA. Unpublished primary dermal irritation test of self-emulsifying wax.*
175. HUNTINGDON RESEARCH CENTER. (April 28, 1977). Submission of data by CTFA. Unpublished acute skin irritation of emulsifying wax.*
176. LEBERCO LABS. (April 2, 1973). Submission of data by CTFA. Unpublished acute skin irritation of emulsifying wax.*
177. LEBERCO LABS. (July 7, 1973). Submission of data by CTFA. Unpublished primary skin irritation study of emulsifying wax.*
178. BIO-TOXICOLOGY LABORATORIES (BTL). (Sept. 12, 1973). Submission of data by CTFA. Unpublished human repeated insult patch test of emulsifying wax.*
179. HTRL. (Sept. 28, 1979). Submission of data by CTFA. Unpublished human study of cumulative irritant properties of a product containing 8 percent emulsifying wax.*
180. CTFA. (Feb. 13, 1981). Submission of data. Unpublished human repeated insult patch test of product containing 8 percent emulsifying wax. (Date of CTFA Transmittal letter).*
181. CTFA. (Jan. 9, 1980). Submission of data. Unpublished human repeated insult patch test of product containing 8 percent emulsifying wax.*
182. CTFA. (Sept. 17, 1979). Submission of data. Unpublished human repeated insult patch test of product containing 8 percent emulsifying wax.*
183. LEO WINTER ASSOCIATES. (Feb. 13, 1981). Submission of data by CTFA. Unpublished repeated insult patch testing and in-use testing of 8 percent emulsifying wax information. (Date of CTFA transmittal letter).*
184. LEO WINTER ASSOCIATES. (Feb. 13, 1981). Submission of data by CTFA. Unpublished human repeated insult patch test on product containing 8 percent emulsifying wax. (Date of CTFA transmittal letter).*
185. HADERT, H. (Sept. 1971). *Wax: History, Type and Applications*. American Ink Maker.
186. CTFA. (Nov. 18, 1980). Submission of data. Unpublished cosmetic safety evaluation on formulation containing 6.0 percent synthetic wax.*
187. MB RESEARCH LABS (MBRL). (April 9, 1979). Submission of data by CTFA. Unpublished acute oral toxicity test in rats of synthetic beeswax.*
188. LEBERCO LABS. (July 13, 1976). Submission of data by CTFA. Unpublished acute ocular toxicity of synthetic beeswax.*
189. MBRL. (April 5, 1979). Submission of data by CTFA. Unpublished acute eye irritation of synthetic beeswax.*
190. MBRL. (March 23, 1979). Submission of data by CTFA. Unpublished primary dermal irritation in rabbits of synthetic beeswax.*
191. LEBERCO LABS. (July 2, 1976). Submission of data by CTFA. Unpublished primary skin irritation of synthetic beeswax.*
192. MBRL. (May 2, 1979). Submission of data by CTFA. Unpublished skin sensitization on guinea pigs of synthetic beeswax.*
193. CTFA. (Oct. 30, 1980). Submission of data. Unpublished human repeated insult patch test, photopatch and use test of product containing 10 percent white beeswax and 7.2–9.4 percent synthetic beeswax.*

Pepe, R. C., J. A. Wenninger, G. N. McEwen, Jr. eds. 2002. *International cosmetic ingredient dictionary and handbook*, 9th ed., vols. 1-4. Washington, DC: CTFA.

Suzuki, M., K. Asaba, H. Komatsu, and M. Mochizuka. 1978. Autoradiographic study on percutaneous absorption of several oils useful for cosmetics. *J. Soc. Cosmet. Chem.* 29:265-282.

Tucker, S. B., S. A. Flannigan, M. Dunbar, and R. B. Drotman. 1986. Development of an objective comedogenicity assay. *Arch. Dermatol.* 122:660-665.

OZOKERITE, CERESIN, MONTAN WAX, PARAFFIN, MICROCRYSTALLINE WAX, EMULSIFYING WAX N.F., SYNTHETIC WAX, AND SYNTHETIC BEESWAX

A safety assessment of these fossil and synthetic waxes was published in 1984 with the conclusion that these ingredients are "safe as cosmetic ingredients in the practices and concentrations of use" (Elder 1984). New studies listed at the end of this section, with updated information regarding types and concentrations of use, were considered by the CIR Expert Panel. The Panel determined to not reopen this safety assessment.

Ozokerite was used in 1269 cosmetic products in 1976, with the largest use occurring in moisturizing products at concentrations of >1% to 50%. In 2002, Ozokerite was used in 680 cosmetic products (FDA 2002), at a maximum use concentration of 22% in other eye makeup preparations (CTFA 2003).

Ceresin was used in 403 cosmetic products in 1976, with the largest use occurring in makeup bases; rouges; other makeup preparations; cleansing; face, body, and hand skin care; moisturizing; night, wrinkle smoothers; and other skin care preparations at concentrations of ≤25%. In 2002, Ceresin was used in 404 cosmetic products (FDA 2002), at a maximum use concentration of 20% in other makeup preparations and suntan gels, creams, and liquids (CTFA 2003).

Montan Wax was used in 355 cosmetic products in 1976, with the largest use occurring in eyebrow pencil and lipsticks at concentrations of >0.1% to 25%. In 2002, Montan Wax was

used in 13 cosmetic products (FDA 2002), at a maximum use concentration of 11% in foundations (CTFA 2003).

Paraffin was used in 1238 cosmetic products in 1976, with the largest use occurring in other eye makeup preparations; perfumes; other fragrance preparations; other hair coloring preparations; other makeup preparations; face, body, and hand skin care; moisturizers; and other skin care preparations at concentrations of ≤50%. In 2002, Paraffin was used in 481 cosmetic products (FDA 2002), at a maximum use concentration of 99% in other skin care preparations (CTFA 2003).

Microcrystalline Wax was used in 868 cosmetic products in 1976, with the largest use occurring in other fragrance preparations at concentrations of >25% to 50%. In 2002, Microcrystalline Wax was used in 581 cosmetic products (FDA 2002), at a maximum use concentration of 50% in other hair preparations (CTFA 2003).

Emulsifying Wax N.F. was used in 12 cosmetic products in 1976, with the only use occurring in other skin care preparations at concentrations of >0.1% to 10%. In 2002, Emulsifying Wax N.F. was used in 102 cosmetic products (FDA 2002), at a maximum use concentration of 21% in hair straighteners (CTFA 2003). This ingredient is no longer listed as a cosmetic ingredient in the *International Cosmetic Ingredient Dictionary and Handbook* (Pepe et al. 2002).

Synthetic Beeswax was used in 146 cosmetic products in 1976, with the largest use occurring in foundations and rouges at concentrations of >1% to 5%. In 2002, Synthetic Beeswax was used in 179 cosmetic products (FDA 2002), at a maximum use concentration of 18% in mascara (CTFA 2003).

Synthetic Wax was used in five cosmetic products in 1976, with the largest use occurring in lipstick at concentrations of >1 to 10%. In 2002, Synthetic Wax was used in 215 cosmetic products (FDA 2002), at a maximum use concentration of 29% in lipstick (CTFA 2003).

Table 17 presents the historical and current cosmetic product uses and use concentrations of these fossil and synthetic waxes.

TABLE 17
Historical and current cosmetic product uses and concentrations for fossil and synthetic waxes

Product type	1976 uses (Elder 1984)	2002 uses (FDA 2002)	1976 use concentrations (Elder 1984) (%)	2003 use concentrations (CTFA 2003) (%)
<i>Ozokerite</i>				
Baby lotions, oils, powders, and creams	3	1	>1-5	2
Bath preparations (other)	—	—	—	15
Eyebrow pencil	—	8	—	—
Eyeliner	—	11	—	7-10
Eye shadow	152	10	>0.1-25	6-14
Eye makeup remover	2	3	>10-25	2-4
Mascara	71	25	>1-25	7-9
Eye makeup (other)	17	26	>0.1-25	2-22
Colognes and toilet waters	—	1	—	15

(Continued on next page)

TABLE 17

Historical and current cosmetic product uses and concentrations for fossil and synthetic waxes (*Continued*)

Product type	1976 uses (Elder 1984)	2002 uses (FDA 2002)	1976 use concentrations	2003 use concentrations
			(Elder 1984) (%)	(CTFA 2003) (%)
Perfumes	41	10	>25–50	15
Sachets	3	—	>1–5	—
Fragrance preparations (other)	7	2	>10–25	15
Hair Conditioners	1	1	>5–10	—
Hair tonics, dressings, etc.	—	3	—	—
Hair preparations (other noncoloring)	1	2	>0.1–1	—
Hair dyes and colors	—	46	—	—
Hair-coloring preparations (other)	—	—	—	0.3
Blushers (all types)	65	10	>0.1–25	2–21
Face powders	1	1	>5–10	1
Foundations	45	33	>0.1–25	4–15
Leg and body paint	1	—	>1–5	8
Lipsticks	559	374	>0.1–25	2–21
Makeup bases	149	11	>0.1–25	—
Rouges	35	2	>1–25	—
Makeup fixatives	1	—	>0.1–1	—
Makeup (other)	36	37	>1–25	3–21
Cuticle softeners	1	1	>5–10	—
Nail polish and enamel	—	—	—	3
Bath soaps and detergents	—	—	—	0.08
Personal cleanliness products (other)	—	1	—	—
Skin-cleansing creams, lotions, liquids, and pads	17	4	>0.1–10	1–5
Face and neck skin care preparations	18*	2	>0.1–50*	—
Body and hand skin care preparations	—	10	—	7
Hormone preparations**	1	—	>1–5	—
Moisturizers	17	15	>0.1–25	1–5
Night skin care preparations	9	7	>1–50	2
Paste masks/mud packs	—	4	—	—
Skin care preparations (other)	8	7	>1–25	6
Suntan gels, creams, and liquids	5	9	>1–25	8–15
Suntan preparations (other)	3	3	>1–5	3–12
Total uses/ranges for Ozokerite	1269	680	>0.1–50	1–22
<i>Ceresin</i>				
Baby lotions, oils, powders, and creams	1	—	>5–10	—
Bath preparations (other)	2	—	>10–25	—
Eyebrow pencil	—	29	—	8
Eyeliner	3	43	>1–5	5–12
Eye shadow	48	31	>25–50	11
Eye lotion	2	2	>5–10	—
Eye makeup remover	2	3	>5–10	—
Mascara	32	9	>0.1–25	4
Eye makeup (other)	17	4	>1–25	4–11
Colognes and toilet waters	6	—	>5–10	—
Perfumes	8	1	>10–25	—
Fragrance preparations (other)	33	—	>10–25	—
Hair conditioners	1	—	>5–10	—

(Continued on next page)

TABLE 17
 Historical and current cosmetic product uses and concentrations for fossil and synthetic waxes (*Continued*)

Product type	1976 uses (Elder 1984)	2002 uses (FDA 2002)	1976 use concentrations	2003 use concentrations
			(Elder 1984) (%)	(CTFA 2003) (%)
Hair tonics, dressings, etc.	7	4	>0.1–25	2
Wave sets	1	—	>5–10	—
Hair preparations (other noncoloring)	2	1	>1–10	17
Hair-coloring preparations (other)	—	1	—	—
Blushers (all types)	19	11	>0.1–25	6
Face powders	4	—	>5–10	6
Foundations	7	6	>1–25	4–7
Lipsticks	109	191	>0.1–25	0.8–20
Makeup bases	7	2	≤0.1–25	—
Rouges	1	2	>10–25	—
Makeup fixatives	—	—	—	3
Makeup (other)	19	15	>0.1–25	4–13
Nail creams and lotions	—	2	—	—
Nail polish and enamel	—	1	—	4
Underarm deodorant	1	—	>1–5	—
Skin-cleansing creams, lotions, liquids, and pads	29	13	≤0.1–25	—
Face and neck skin care preparations	5*	1	>0.1–25*	5–8
Body and hand skin care preparations	—	5	—	8
Hormone preparations**	1	—	>1–5	—
Moisturizers	13	11	>1–25	5
Night skin care preparations	12	4	>1–25	5
Paste masks/mud packs	—	1	—	—
Skin care preparations (other)	9	9	>5–25	—
Suntan gels, creams, and liquids	—	2	—	13
Other suntan preparations	1	—	>5–10	—
Total uses/ranges for Ceresin	403	404	≤0.1–25	0.8–20
<i>Montan Wax</i>				
Eyebrow pencil	5	—	>10–25	—
Eyeliner	—	—	—	2
Eye Shadow	18	—	>1–5	9
Mascara	—	—	—	3
Eye makeup (other)	—	—	—	0.5
Perfumes	1	—	>0.1–1	—
Hair-coloring preparations (other)	—	1	—	—
Foundations	—	5	—	11
Lipsticks	323	7	>0.1 — 25	0.5
Makeup (other)	—	—	—	0.5
Nail polish and enamel	—	—	—	0.5
Rouges	7	—	>5–10	—
Face and neck skin care preparations	1*	—	>0.1–1*	—
Body and hand skin care preparations	—	—	—	—
Total uses/ranges for Montan Wax	355	13	>0.1–25	0.5–11
<i>Paraffin</i>				
Baby shampoos	1	—	>1–5	—
Baby lotions, oils, powders, and creams	—	—	—	1
Eyebrow pencil	—	4	—	6

(Continued on next page)

TABLE 17
 Historical and current cosmetic product uses and concentrations for fossil and synthetic waxes (*Continued*)

Product type	1976 uses (Elder 1984)	2002 uses (FDA 2002)	1976 use concentrations (Elder 1984) (%)	2003 use concentrations (CTFA 2003) (%)
Eyeliner	—	7	—	4–20
Eye shadow	127	5	>0.1–25	3–8
Eye lotion	—	1	—	1–6
Eye makeup remover	3	3	>1–10	—
Mascara	3	28	>1–5	6–20
Other eye makeup preparations	20	4	>0.1–50	4
Colognes and toilet waters	3	—	>10–25	—
Perfumes	30	—	>25–50	—
Sachets	6	—	>5–10	—
Other fragrance preparation	15	1	>25–50	—
Hair conditioners	14	7	>5–10	2
Hair rinses (noncoloring)	3	—	>1–5	—
Shampoos (noncoloring)	2	1	>0.1–1	—
Tonics, dressings, and other hair-grooming aids	29	23	>0.1–25	2–20
Wave sets	1	1	>10–25	—
Other hair preparations	3	1	>1–10	2
Hair tints	—	1	—	—
Hair colors sprays (aerosol)	—	2	—	—
Other hair-coloring preparations	3	10	>10–50	—
Blushers (all types)	65	4	>0.1–25	—
Face powders	15	—	>0.1–1	—
Foundations	76	33	>0.1–10	2–6
Lipstick	439	134	≤0.1–25	2–10
Makeup bases	95	37	>0.1–10	—
Rouges	29	—	>0.1–25	—
Makeup fixatives	3	1	>1–5	—
Other makeup preparations	44	18	>0.1–50	2–21
Cuticle softeners	3	1	>1–5	—
Nail creams and lotions	2	3	>1–25	—
Nail polish and enamel	—	5	—	0.03–2
Other manicuring preparations	1	—	>1–5	0.03
Bath soaps and detergents	3	7	≤0.1	—
Deodorants (underarm)	1	2	>5–10	—
Other personal cleanliness products	3	16	>1–10	—
Aftershave lotion	3	—	>0.1–1	—
Shaving cream	1	1	>1–5	1
Cleansing	69	26	>0.1–25	12–20
Depilatories	1	1	>5–10	—
Face and neck (excluding shave)	31*	4	≤0.1–50*	—
Body and hand (excluding shave)	—	19	—	3
Hormone preparations**	2	—	≤0.1–1	—
Foot powders and sprays	—	1	—	—
Moisturizers	28	24	>1.0–50	2
Night skin care preparations	11	6	>1–10	0.3–8
Paste masks/mud packs	4	7	>0.1–5	32
Skin lighteners**	4	—	>1–5	—

(Continued on next page)

TABLE 17
 Historical and current cosmetic product uses and concentrations for fossil and synthetic waxes (*Continued*)

Product type	1976 uses (Elder 1984)	2002 uses (FDA 2002)	1976 use concentrations	2003 use concentrations
			(Elder 1984) (%)	(CTFA 2003) (%)
Skin fresheners	1	—	>10–25	—
Skin care preparations (other)	22	27	>1–50	99
Suntan gels, creams, and liquids	13	3	>0.1–10	2–10
Indoor tanning preparations	—	—	—	2
Suntan preparations (other)	6	2	>1–25	20
Total uses/ranges for Paraffin	1238	481	≤0.1–50	0.03–99
<i>Microcrystalline Wax</i>				
Baby products (other)	—	—	—	3
Bath oils, tablets, and salts	—	1	—	—
Bath preparations (other)	—	—	—	0.5
Eyebrow pencil	2	15	>0.1–10	6
Eyeliner	—	126	—	2–18
Eye shadow	37	16	>0.1–25	2–6
Eye lotion	—	—	—	0.3
Eye makeup remover	2	1	>1–5	—
Mascara	5	20	>1–25	3–6
Eye makeup (other)	23	33	>0.1–25	8–19
Perfumes	17	1	>10–25	—
Fragrance preparations (other)	9	1	>25–50	—
Hair conditioners	2	3	>1–5	4
Hair straighteners	—	1	—	0.4
Hair tonics, dressings, etc.	14	15	>1–25	0.8
Hair preparations (other noncoloring)	2	1	>1–10	3–50
Hair-coloring preparations (other)	1	1	>1–5	—
Blushers (all types)	38	8	>1–25	6–10
Face powders	1	1	>1–5	—
Foundations	9	35	>0.1–10	9–12
Lipstick	640	229	>0.1–25	0.5–10
Makeup bases	6	2	≤0.1–10	—
Rouges	—	—	—	7
Makeup fixatives	—	1	—	—
Makeup (other)	17	33	>0.1–10	3–13
Cuticle softeners	1	—	>5–10	—
Oral hygiene products (other)	—	—	—	10
Shaving preparation products (other)	—	—	—	0.5
Nail creams and lotions	1	—	>1–5	—
Nail polish and enamel	—	1	—	—
Personal cleanliness products (other)	—	1	—	—
Skin-cleansing creams, lotions, liquids, and pads	12	6	>0.1–10	2
Depilatories	—	1	—	4
Face and neck skin care preparations	10*	2	>0.1–25*	—
Body and hand skin care preparations	—	5	—	—
Moisturizers	8	5	>0.1–25	—
Night skin care preparations	7	4	>0.1–25	2
Paste masks/mud packs	—	2	—	—
Skin care preparations	4	10	>5–25	1

(Continued on next page)

TABLE 17

Historical and current cosmetic product uses and concentrations for fossil and synthetic waxes (*Continued*)

Product type	1976 uses	2002 uses	1976 use concentrations	2003 use concentrations
	(Elder 1984)	(FDA 2002)	(Elder 1984) (%)	(CTFA 2003) (%)
Suntan gels, creams, and liquids	—	—	—	1–15
Indoor tanning products	—	—	—	15
Suntan preparations (other)	—	—	—	0.2–0.5
Total uses/ranges for Montan Wax	868	581	≤0.1–50	0.2–50
<i>Emulsifying Wax N.F.</i>				
Baby lotions, oils, powders, and creams	—	2	—	—
Eyeliners	—	2	—	—
Eye makeup (other)	—	1	—	—
Fragrance preparation (other)	—	4	—	—
Hair conditioners	—	14	—	0.8–2
Hair straighteners	—	10	—	21
Hair tonics, dressings, etc.	—	—	—	9
Rinses (noncoloring)	—	1	—	—
Permanent waves	—	1	—	—
Hair bleaches	—	1	—	—
Hair coloring preparations (other)	—	6	—	—
Foundations	—	2	—	—
Cuticle softeners	—	2	—	0.5
Nail creams and lotions	—	1	—	—
Skin-cleansing creams, lotions, liquids, and pads	—	2	—	1
Face and neck skin care preparations	—*	—	—*	1
Body and hand skin care preparations	—	12	—	2–10
Moisturizers	—	25	—	12
Night skin care preparations	—	4	—	1
Paste masks/mud packs	—	1	—	1
Skin care preparations (other)	12	5	>0.1–10	—
Indoor tanning preparations	—	5	—	—
Suntan preparations (other)	—	1	—	—
Total uses/ranges for Emulsifying Wax N.F.	12	102	>0.1–10	0.5–21
<i>Synthetic Beeswax</i>				
Baby lotions, oils, powders, and creams	—	1	—	0.5
Baby products (other)	—	1	—	—
Eyebrow pencils	—	6	—	—
Eyeliners	—	9	—	0.06–14
Eye shadow	119	1	>0.1–1	—
Mascara	—	6	—	0.05–18
Eye makeup (other)	—	1	—	0.06–9
Fragrance preparations (other)	—	1	—	—
Hair tonics, dressings, etc.	—	—	—	12
Blushers	14	—	>0.1–1	—
Face powders	10	—	>0.1–1	—
Foundations	1	2	>1–5	—
Leg and body paints	—	—	—	13
Lipstick	—	131	—	4–7
Makeup bases	—	1	—	—
Rouges	2	—	>1–5	—

(Continued on next page)

TABLE 17
Historical and current cosmetic product uses and concentrations for fossil and synthetic waxes (*Continued*)

Product type	1976 uses (Elder 1984)	2002 uses (FDA 2002)	1976 use concentrations	2003 use concentrations
			(Elder 1984) (%)	(CTFA 2003) (%)
Makeup (other)	—	9	—	1
Skin-cleansing creams, lotions, liquids, and pads	—	1	—	—
Face and neck skin care preparations	—*	1	—*	—
Body and hand skin care preparations	—	2	—	1–3
Moisturizers	—	5	—	0.5
Skin care preparations (other)	—	1	—	—
Total uses/ranges for Synthetic Beeswax	146	179	>0.1–5	0.05–18
<i>Synthetic Wax</i>				
Baby lotions, oils, powders, and creams	—	—	—	4
Eyebrow pencils	—	2	—	—
Eyeliners	—	28	—	0.8–27
Eye shadow	—	9	—	—
Mascara	—	10	—	2–11
Eye makeup (other)	—	1	—	—
Perfumes	—	1	—	—
Fragrance preparation (other)	—	2	—	—
Hair conditioners	—	1	—	—
Blushers (all types)	1	10	>1–5	—
Face powders	—	2	—	0.5
Foundations	1	6	>1–5	9
Lipsticks	3	134	>1–10	3–29
Makeup bases	—	2	—	—
Rouges	—	1	—	2
Makeup (other)	—	4	—	—
Cuticle softeners	—	—	—	0.1
Aftershave lotions	—	—	—	1
Skin-cleansing creams, lotions, liquids, and pads	—	—	—	5
Face and neck skin care preparations	—*	—	—*	3
Body and hand skin care preparations	—	—	—	2–4
Moisturizers	—	1	—	—
Night skin care preparations	—	—	—	3
Personal cleanliness products (other)	—	1	—	—
Suntan preparations (other)	—	—	—	1
Total uses/ranges for Synthetic Wax	5	215	>1–10	0.1–29

*This category was combined when the original safety assessment was performed and is now two separate categories.

**No longer considered a cosmetic product category.

REFERENCES

Benedetto, G. D., M. Pierangeli, A. Scalise, and A. Bertani. 2002. Paraffin oil injection in the body: An obsolete and destructive procedure. *Ann. Plastic Surg.* 49:391–396.

Cosmetic, Toiletry, and Fragrance Association (CTFA). 2003. Concentrations of use—mineral waxes. Unpublished data submitted by CTFA on January 21, 2003. 6 pages.¹⁸

Elder, R. L. 1984. Final report on the safety assessment of fossil and synthetic waxes. *J. Am. Col. Toxicol.* 3:43–99.

¹⁸Available from the Director, Cosmetic Ingredient Review, 1101 17th Street NW, Suite 310, Washington, DC 20036, USA.

Food and Drug Administration (FDA). 2002. Frequency of use of cosmetic ingredients. *FDA database*. Washington, DC: FDA.

Heimbach, J. T., A. R. Bodor, J. S. Douglas, et al. 2002. Dietary exposures to mineral hydrocarbons from food-use applications in the United States. *Food Chem. Toxicol.* 40:555–571.

Hintschich, C. R., C. K. Beyer-Machule, and F. H. Stefani. 1995. Paraffinoma of the periorbit—a challenge for the oculoplastic surgeon. *Ophthalm. Plastic Reconstr. Surg.* 11:39–44.

Joint FAO/WHO Expert Committee on Food Additives (JECFA). 1993. Microcrystalline wax and paraffin wax. *WHO Food Addit. Ser.* 30:253–260.

JECFA. 1996. Mineral oils (food grade), paraffin waxes and microcrystalline waxes. *WHO Food Addit. Ser.* 35:193–206.

Pepe, R. C., J. A. Wenninger, and G. N. McEwen, Jr. 2002. *International cosmetic ingredient dictionary and handbook*. 9th ed. Washington, DC: CTFA.

Smith, J. H., A. K. Mallett, A. J. Preston, et al. 1996. Ninety-day feeding study in Fischer-344 rats of highly refined petroleum-derived food-grade white oils and waxes. *Toxicol. Pathol.* 24:214–230.

Vigneswaran, W. T., and S. S. Ramasastry. 1996. Paraffin plompage of the chest revisited. *Ann. Thorac. Surg.* 62:1837–1839.

Witschel, H., and K. Geiger. 1994. Paraffin induced sclerosing lipogranuloma of eyelids and anterior orbit following endonasal sinus surgery. *Br. J. Ophthalmol.* 78:61–65.

~~PEG-2, 3, 4, 5, 6, 6-32, 7, 8, 9, 10, 12, 14, 15, 18, 20, 23, 25, 30, 32, 35, 36, 40, 45, 50, 55, 75, 90, 100, 120, AND 150 STEARATES~~

A safety assessment of PEG-2, 6, 8, 12, 20, 32, 40, 50, 100, and 150 Stearates was published in 1983 with the conclusion “safe as cosmetic ingredients in the present practices of

concentration and use” (Elder 1983). Studies available since that safety assessment was completed, along with updated information regarding use concentrations, were considered by the CIR Expert Panel. The Panel determined not to reopen this safety assessment.

In 1979, PEG Stearates were used in 374 cosmetic products, typically at concentrations ranging from >0.1% to 10%. Currently, there are uses reported in 1459 products, typically at concentrations <4%. Table 18 presents the available use information.

REFERENCES

Barany, E., M. Lindberg, and M. Loden. 2000. Unexpected skin barrier influence from nonionic emulsifiers. *Int. J. Pharm.* 195:189–195.

Bryan, A. J., R. Kaur, G. Robinson, N. W. Thomas, and C. G. Wilson. 1980. Histological and physiological studies on the intestine of the rat exposed to solutions of Myrj 52 and PEG 2000. *Int. J. Pharm.* 7:145–156.

TABLE 18
Historical and current cosmetic product uses and concentrations for PEG Stearates

Product type	1976 uses (Elder 1983)	2002 uses (FDA 2002)	1976 use concentrations (Elder 1983) (%)	2002 use concentrations (CTFA 2002) (%)
<i>PEG-2 Stearate</i>				
Bath preparations (other)	1	—	>0.1–1	—
Eye makeup (other)	—	—	—	2
Sachets	—	1	—	—
Fragrance preparations	11	—	>0.1–1	—
Shampoos (noncoloring)	—	2	—	—
Hair tonics, dressings, etc.	—	1	—	—
Hair preparations (other noncoloring)	4	—	>0.1–10	—
Hair coloring preparations (other)	1	—	>0.1–1	—
Foundations	—	4	—	0.2–1
Makeup bases	—	1	—	—
Makeup (other)	16	—	>0.1–10	2
Manicuring preparations (other)	1	—	>1–5	—
Personal cleanliness products (other)	1	—	>0.1–1	—
Aftershave lotion	—	3	—	0.4
Shaving preparations (other)	6	—	>0.1–1	—
Skin-cleansing creams, lotions, liquids, and pads	—	6	—	2
Face and neck skin care preparations	—	1	—	—
Body and hand skin care preparations	—*	7	—*	—
Moisturizers	—	13	—	—
Night skin care preparations	—	9	—	—
Paste masks/mud packs	—	4	—	—
Skin care preparations (other)	20	8	>0.1–10	—
Suntan gels, creams, and liquids	1	2	>1–5	2
Indoor tanning preparations	—	1	—	1
Suntan preparations (other)	—	1	—	—
Total uses/ranges for PEG-2 Stearate	62	64	>0.1–10	0.2–2
Body and hand skin care preparations	—	5	—	—

(Continued on next page)