
Safety Assessment of Mink Oil as Used in Cosmetics

Status: Re-Review for Panel Consideration
Release Date: September 1, 2022
Panel Meeting Date: September 26-27, 2022

The Expert Panel for Cosmetic Ingredient Safety members are: Chair, Wilma F. Bergfeld, M.D., F.A.C.P.; Donald V. Belsito, M.D.; David E. Cohen, M.D.; Curtis D. Klaassen, Ph.D.; Daniel C. Liebler, Ph.D.; Allan E. Rettie, Ph.D.; David Ross, Ph.D.; Thomas J. Slaga, Ph.D.; Paul W. Snyder, D.V.M., Ph.D.; and Susan C. Tilton, Ph.D. The Cosmetic Ingredient Review (CIR) Executive Director is Bart Heldreth, Ph.D. This report was prepared by Regina Tucker, M.S., Scientific Analyst/Writer, CIR.



Commitment & Credibility since 1976

Memorandum

To: Expert Panel for Cosmetic Ingredient Safety Members and Liaisons
From: Regina Tucker, M.S. Scientific Analyst/Writer, CIR
Date: September 1, 2022
Subject: Mink Oil

The Expert Panel for Cosmetic Ingredient Safety (Panel) first published a review on the safety of Mink Oil in 1998 (identified as *originalreport1_MinkOil_092022* in the pdf), with an insufficient data conclusion. Subsequently, the Panel's data needs were met, and in 2005, a final amended report was published with the conclusion that Mink Oil is safe as a cosmetic ingredient in the practices of use and concentration described in the safety assessment (*originalreport2_MinkOil_092022*).

Because it has been at least 15 years since the previous report was published, in accord with Cosmetic Ingredient Review (CIR) Procedures, the Panel should consider whether the safety assessment of Mink Oil should be re-opened. An exhaustive search of the world's literature was performed for studies dated 2000 forward. No relevant published data were found. An historical overview, comparison of original and new use data, and the search strategy used are enclosed herein (*newdata_MinkOil_092022*).

Also included for your review is a table of current and historical use data (*usetable_MinkOil_092022*). Since the final amended report was issued there is one new use category reported in 2022 (hair-coloring). The frequency of use has decreased, from 100 reported uses in 2001 to 32 uses in 2002, and the maximum concentration of use for Mink Oil has gone from 3% in 2001 to 0.1% in 2022.

If, upon review of the updated use data the Panel determines that a re-review is warranted, a draft amended report will be presented at an upcoming meeting.

Re-Review - Mink Oil - History and New Data

(Regina Tucker – September 2022 meeting)

Ingredient (1)	Citation	Conclusion	Use - New Data	Use -Historical Data	Notes
Mink Oil	IJT 17(S4): 71 – 81, 1998 IJT 24 (suppl. 3):57-64, 2005	insufficient data safe as used	<u>Mink Oil</u> frequency of use (2022): 32 uses conc of use (2022): 0.1%	<u>Mink Oil</u> frequency of use (2001): 100 uses conc of use (2001): 0.001-3 %	frequency of use decreased; concentration of use decreased; one new use category Hair-Coloring

NOTABLE NEW DATA			
Publication	Study Type	Results – Brief Overview	Different from Existing Data?
<i>no new published data</i>			

Search (from 2000 on)

PubMed

((("mink oil") OR (8023-74-3[EC/RN Number])) AND (("2000"[Date - Publication]: "3000"[Date - Publication])) – 1 hit; none useful

Current and historical frequency and concentration of use according to duration and exposure of Mink Oil

	# of Uses		Max Conc of Use (%)	
	2022 ¹	2001 ²	2022 ³	2001 ²
Totals*	32	100	0.1	0.001-3
<i>Duration of Use</i>				
<i>Leave-On</i>	22	63	0.1	0.001-2
<i>Rinse-Off</i>	5	36	NR	3
<i>Diluted for (Bath) Use</i>	5	1	NR	0.1
<i>Exposure Type</i>				
Eye Area	NR	2	NR	0.1-1
Incidental Ingestion	11	12	NR	0.001
Incidental Inhalation-Spray	5 ^a ;3 ^c	3;23 ^a ;12 ^c	NR	1;0.2-2 ^a ;0.1-0.5 ^c
Incidental Inhalation-Powder	1;3 ^c	1;12 ^c	NR	0.1;1 ^b ;0.1-0.5 ^c
Dermal Contact	19	48	0.1	0.1-2
Deodorant (underarm)	NR	NR	NR	NR
Hair - Non-Coloring	1	39	NR	3
Hair-Coloring	1	NR	NR	NR
Nail	NR	NR	NR	0.001-0.1
Mucous Membrane	16	13	NR	NR
Baby Products	NR	NR	NR	NR

*Because each ingredient may be used in cosmetics with multiple exposure types, the sum of all exposure types may not equal the sum of total uses.

^a It is possible these products are sprays, but it is not specified whether the reported uses are sprays.

^b It is possible these products are powders, but it is not specified whether the reported uses are powders.

^c Not specified whether a spray or a powder, but it is possible the use can be as a spray or a powder, therefore the information is captured in both categories

NR – no reported use

References

1. U.S. Food and Drug Administration Center for Food Safety & Applied Nutrition (CFSAN). 2022. Voluntary Cosmetic Registration Program - Frequency of use of Cosmetic Ingredients. College Park, MD. Obtained under the Freedom of Information Act from CFSAN; requested as "Frequency of Use Data" January 4, 2022; received January 11, 2022
2. Andersen FA. Final amended report on the safety assessment of Mink Oil. *Int J Toxicol*. 2005;24 Suppl 3:57-64.
3. . Personal Care Products Council 2020 Concentration of Use by FDA Product Category: Mink Oil. (Unpublished data submitted by the Personal Care Products Council on October 13, 2020.)

FINAL REPORT ON THE SAFETY ASSESSMENT OF MINK OIL¹

Mink Oil is obtained from the subdermal fatty tissues of the mink. The oil is a mixture of the natural glycerides of 14–20 carbon chain fatty acids. Mink Oil is used in cosmetic formulations as a hair conditioning agent, an occlusive skin conditioning agent, and as a surfactant. Current data on concentration of use is limited. A Mink Oil cream was noncomedogenic in rabbits following repeated (5 days a week for 2 weeks) applications, although minimal hyperemia and hyperplasia were noted at the treated site toward the end of the treatment. No sensitization was observed in animals treated with Mink Oil under occlusive patches, but sensitization was seen in animals treated with intradermal injections of Mink Oil in a maximization procedure. Mink Oil was not an ocular irritant in animals. Clinical studies done under occlusive patches found no irritation. In another clinical test, a rinse-off product containing 2.8% Mink Oil was applied to the skin for five consecutive days without rinsing. Some erythema was noticed, but in a repeat of the test, no erythema was seen. Mink Oil is reported to provide a minimal sun protection factor (SPF), but did not appear to increase the SPF when added to a solution containing recognized sunscreens. Studies using human volunteers indicate that Mink Oil applied to the skin remains on the skin surface. Additional safety test data are needed, including a 28-day dermal toxicity study, ultraviolet (UV) absorption data (and if the ingredient absorbs in the UVA or UVB region, then photosensitization data are also needed), and method of manufacture (extraction). Independent of these data needs, it is concluded that the total polychlorinated biphenyl (PCB)/pesticide contamination should not exceed 40 ppm with not more than 10 ppm for any specific residue. Because there are additional data needed in order to complete the safety assessment, the overall conclusion is that the available data are insufficient to support the safety of Mink Oil for use in cosmetic products.

CHEMISTRY

Definition

Mink Oil is obtained from the subdermal fatty tissues of the mink (Wenninger and McEwen 1997). The oil is a mixture of the natural glycerides of 14–20 carbon chain fatty acids; approximately 75% of the

Received 20 January 1998; accepted 30 April 1998.

¹Reviewed by the Cosmetic Ingredient Review Expert Panel. Bindu Nair, Scientific Analyst and Writer, prepared this report.

Address correspondence and reprint requests to Bindu Nair, Cosmetic Ingredient Review, 1101 17th Street, NW, Suite 310, Washington, DC 20036, USA.

composition is 16–18 carbon chain fatty acid glycerides (Nikitakis and McEwen 1990).

Physical and Chemical Properties

Mink Oil is a pale yellow liquid with a mild characteristic odor. It is soluble in chloroform, carbon tetrachloride, ether, benzene, acetone, and isopropanol and is insoluble in water. The refractive index at 25°C is between 1.4665 to 1.4675. The freezing point is 5°C maximum; the saponification value is 190 to 220; the iodine value is 80 to 90 (Nikitakis and McEwen 1990). It has a specific gravity of 0.91437, a refractive index at 40°C of 1.4623 (SGS Control Services Inc. 1994). One source reports that because of its iodine value, Mink Oil “exhibits very low oxidation and degradation” (Emulan Inc., unknown date). Table 1 lists the acid composition found in Mink Oil Wax and crude Mink Oil.

Impurities

Analysis of one sample of Mink Oil found no detectable levels of the following PCBs, Arochlor 1016, 1221, 1232, 1242, 1248, 1254, and 1260.

Table 1. Fatty acid composition of two mink oils

Acid	Carbon no.: unsat	Mink wax oil	Mink crude oil
		(% of total)	
Lauric	12	0.1	0.1
Myristic	14	4.0	3.5
Myristoleic	14:1	0.7	0.9
Pentadecanoic	15	0.2	0.1
Palmitic	16	28.0	17.2
Palmitic	16:1	13.3	17.0
Heptadecanoic	17	0.5	0.4
Heptadecanoic	17:1	0.6	0.5
Stearic	18	4.7	2.5
Oleic	18:1	35.3	40.9
Linoleic	18:2	10.6	15.0
Linolenic	18:3	0.4	0.6
Eicosanoic	20	0.1	—
Eicolenic	20:1	0.7	0.6
Total unsaturated		61.6	75.5
Total saturated		37.6	23.8
Total		99.2	99.3

Unsat, unsaturated.

Source. Emulan, date unknown.

The assay had a minimum detection limit of 0.5 $\mu\text{g/l}$ (New Jersey Laboratories/A.A. Labs Division 1995).

USE

Cosmetic

As shown in Table 2, as of January 1995, Mink Oil was reported to be contained in 139 formulations (FDA 1995). It is used in formulations as a hair conditioning agent, an occlusive skin conditioning agent, and a surfactant (Wenninger and McEwen 1997).

Concentrations of use are no longer reported to the Food and Drug Administration (FDA) (FDA 1992). Data from 1984 (FDA 1984) indicated

Table 2. Frequency of use of Mink Oil

Product category	No. formulations in category	No. containing mink oil
Bath oils, tablets, and salts	146	5
Other bath preparations	204	7
Mascara	211	2
Perfumes	260	1
Powders (excluding aftershave talcum)	290	1
Other fragrance preparations	158	1
Hair conditioner	693	10
Hair spray (aerosol fixatives)	348	5
Hair straighteners	59	2
Permanent waves	423	4
Shampoos (noncoloring)	916	5
Tonics, dressings, and other hair grooming aids	624	18
Other hair preparations	382	5
Face powders	305	1
Foundations	333	1
Lipstick	997	12
Shaving cream	152	3
Cleansing	771	12
Face and neck	261	2
Body and hand	987	11
Moisturizing	873	16
Night	220	4
Paste masks (mud packs)	276	2
Other skin care preparations	782	8
Suntan gels, creams, liquids	196	1
1995 totals		139

Source. FDA, 1995.

that mink oil was used in a variety of products at concentrations less than 25%. Data submitted directly to Cosmetic Ingredient Review (CIR) show Mink Oil to be used at 2% in self-tanning products and at 0.2% in hair products (CTFA 1995).

GENERAL BIOLOGY

Absorption

An Attenuated Total Reflectance–InfraRed Spectroscopy study was conducted to determine the penetration of a commercial preparation of Mink Oil into human stratum corneum. Five healthy Caucasian females were instructed to refrain from applying any topical agents to their forearms 24 hours prior to the study. A site on one forearm of each panelist was treated with 50 μ l of the test substance. The same site on the other forearm was not treated and served as control. Both sites were scanned prior to product application. One hour after application, the treated site was again scanned. This was followed by scans of 5, 10, and 15 stripped layers. The same scans were conducted on the untreated sites. Scanning consisted of placing the subject's forearm directly onto a germanium plate and conducting an infrared (IR) spectra for wavelengths between 800 to 4000 cm^{-1} . The germanium plate served as an internal reflection crystal. Based on the IR spectra of the test substance, wavelengths of 1746, 1466, and 1163 cm^{-1} were chosen for analysis. Presence of Mink Oil in each of the four scans (surface, 5, 10, and 15 layers all done at 1 hour post-treatment) was defined as a greater absorbance value at the treated site versus the control. For each of the four scans, there were 15 absorbance pairs (a pair consisting of an absorbance value of the control and treated sites). Mink oil was detected at the surface scan of all panelists. It was detected within 5 layers in 4 of 5 panelists; in 10 layers in 2 of 5 panelists, and in 15 layers in 2 of 5 panelists. It was concluded that after 1 hour Mink Oil remains significantly on the skin surface (Xienta Institute for Skin Research 1988).

Comedogenicity

A Mink Oil cream was considered to be noncomedogenic following repetitive applications (ten applications in two consecutive weeks) to the external ear canal of six albino rabbits. Observations noted include "minimal to mild" hyperemia and "very minimal" hyperkeratosis in the treated areas of all animals towards the end of the treatment period (Wells Laboratories, Inc. 1990b).

ANIMAL TOXICOLOGY

Acute Oral Toxicity

A commercial preparation of Mink Oil had an oral LD₅₀ of >64.0 cc/kg in albino rats (Bio-Toxicology Laboratories, Inc. 1974). In another study, a Mink Oil cream had an oral LD₅₀ of >5.0 ml/kg (Wells Laboratories, Inc. 1990e).

Dermal Irritation

Patches containing 0.5 ml of a commercial preparation of refined Mink Oil were applied to intact and scarified areas of albino rabbits. The treatment areas had earlier been clipped free of hair. Animals were immobilized for a 24-hour period; the sites were evaluated at the end of the treatment period and 24 hours afterwards. Scores were made according to the Draize scale. No erythema or edema was noted (Leberco Laboratories 1972a).

Patches containing 0.5 ml of refined Mink Oil (neat) were applied to intact and abraded skin of six rabbits. No irritation was noted (Primary Irritation Index score 0.0) at observations made 24 and 72 hours after application (Wells Laboratories, Inc. 1979b). Similar results were reported by Wells Laboratories, Inc. (1990d) when a cream containing Mink Oil was tested on three New Zealand white rabbits using the same protocol.

Dermal Sensitization

Over a period of 20 days, eight male guinea pigs received ten subcutaneous injections of a 0.1% suspension of refined Mink Oil; the injections were administered to the clipped back and sides. The test material was suspended in physiological saline and was administered in 0.1-ml doses except for the first injection which contained 0.05 ml. Two weeks after the final induction injection, a test injection of 0.05-ml was administered. Reaction areas were measured 24 hours after each injection. Reactions were noted throughout the induction period. However, as the value for the challenge was "less than for the average of the ten original readings (after induction), it can be said that (Mink Oil) did not produce any sensitization in the guinea pig" (Leberco Laboratories 1972b).

Female Dunkin-Hartley guinea pigs were treated with Mink Oil using the Magnusson-Kligman Maximization Procedure. The procedure involves the following phases: induction, dose range, booster, and challenge. During induction, sites on the upper back were injected intradermally with 50% Freund's complete adjuvant; 5% Mink Oil in propylene glycol; and 5% Mink Oil in 50% Freund's complete adjuvant. Control

animals were injected with complete adjuvant, propylene glycol, and 1:1 propylene glycol in complete adjuvant. During the dose range phase, ten animals were tested with 24-hour occlusive patches containing 25, 50, and 100% Mink Oil. Sites were evaluated for irritation at 24, 48, and 72 hours after patch application. Two of the animals reacted to the 25% concentration with a response graded as "barely perceptible"; by 72 hours the condition cleared. These two animals also reacted to the higher concentrations of Mink Oil; the reactions cleared by the observation at 72 hours. Another four had "barely perceptible" or "mild" reactions to the 50% concentration. These reactions also cleared by 72 hours and three of these four animals did not react to the 100% concentration. Three animals reacted for the first time to the 100% concentration with a "barely perceptible" response (two also had desquamation); these conditions cleared by 72 hours. During the booster phase (one week after induction injections), undiluted Mink Oil was topically applied. As this concentration did not produce irritation in the dose range phase, 5% sodium lauryl sulfate (SLS) was applied to the induction site 24 hours before the booster. Control animals were pretreated with SLS before receiving a booster of undiluted petrolatum; positive controls were treated with 25% phenylacetaldehyde without SLS pretreatment. During the booster phase, all test and control animals were wrapped occlusively for 48 hours. Two weeks following the booster, test animals were challenged with a 24-hour occlusive patch of 25% and 50% Mink Oil in petrolatum. Reactions were scored at 48 and 72 hours after patch application. There were no reactions to 25% Mink Oil noted in the ten animals at the 48 hour reading; one animal had a "barely perceptible" reaction at 72 hours. Four animals had a "barely perceptible" and one had a "mild" response to 50% Mink Oil at the 48 hour reading. At the 72 hour reading, erythema was noted in 7 of 10 animals, which ranged from "barely perceptible" (4 animals), to "desquamation" (2 animals), and "mild with desquamation" (1 animal). The 50% Mink Oil in petrolatum was considered to produce a weak allergenic response (CTFA 1983).

The Buehler Technique was used by Wells Laboratories (1990a) in a guinea pig sensitization assay. During induction, nine occlusive patches containing 0.5 ml of a 50% dilution (in corn oil) of a Mink Oil cream was applied to the clipped back and flank of 10 animals. The patches remained for 6 hours of contact and readings were made at the time of patch removal and at the end of 24 hours. Positive controls were treated with dinitrochlorobenzene (DNCB). After a 2-week nontreatment period animals were challenged with two identical applications of the test material. One patch was applied at the site of induction, and the other on a previously unexposed site. Challenge patches were applied for 6 hours of contact and observations were made at the time of patch removal and at the end of 24 hours. A reading of "very slight" erythema was noted in

all animals during induction; “slight” erythema and “slight” edema were noted in positive controls. During challenge, no irritation was noted in animals of the Mink Oil group; marked erythema and very slight edema was noted in animals of the positive-control group.

Ocular Irritation

A commercial preparation of refined Mink Oil (0.1 ml) was instilled into the right conjunctival sac of three albino rabbits. The untreated left eye served as control. Eyes were evaluated every 24 hours for four days and then again on the seventh day. Scores were made according to the Draize scale. No irritation was noted (score 0) (Leberco Laboratories 1972c). Another ocular irritation test was conducted using three rabbits and the above described protocol (Wells Laboratories 1979a). Undiluted Mink Oil (commercial preparation, dose not reported) did not produce any observable irritation. A Mink Oil cream (0.1 ml) was applied to one eye of each of three New Zealand white rabbits. The contralateral eye was not treated and served as control. Eyes were unwashed for 24 hours and observations made 24, 48, and 72 hours after treatment. No irritation was noted (Draize score 0.01) (Wells Laboratories, Inc. 1990c).

CLINICAL ASSESSMENT OF SAFETY

Dermal Irritation and Sensitization

Twenty panelists received a single 24-hour occlusive patch containing a 25% aqueous dilution of a hair-rinse containing 3% Mink Oil (effective concentration of mink oil tested: 0.75%). No reactions were noted in 16 panelists, reactions of \pm (the mildest nonzero score possible) were noted in four panelists. The Primary Irritation Index (PII) was 0.10 out of a maximum possible score of 4.0 (CTFA 1989). Identical results were reported in a second patch test study in which a formulation containing 2.8% Mink Oil was tested at full-strength in 20 panelists (CTFA 1992a). A third human patch test study reported a PII of 0.08 after 19 panelists were tested with a (full-strength) formulation containing 2.8% Mink Oil (CTFA 1993).

A four-day minicumulative patch test of a body spray containing 1% Mink Oil had a PII of 0.50. Of 20 panelists, 5 had no response, 12 had a “barely perceptible” response, 2 had a “mild” response, and 1 had a “moderate” response (CTFA 1991).

In an exaggerated-use study, 19 female panelists applied to the lower arm a rinse-off hair product containing 2.8% Mink Oil once a day for 5 days with no rinse off. A commercially available rinse-off hair product

was applied to a different site to serve as a reference. Four subjects experienced some transient mild to moderate erythema (CTFA 1992b). In a second study using the same protocol, no clinical responses were observed in 24 females who also tested a rinse-off hair product containing 2.8% Mink Oil (CTFA 1994).

Sun-Protection Factor

Wells Laboratories (1988a) tested the sunscreens ability of Mink Oil using a modified version of the procedure described by FDA (1978) for evaluating the sun protection factor (SPF) of sunscreens (the modification involved use of less than the prescribed 20 subjects). The solutions tested contained Mink Oil in various concentrations with Octyldimethyl *para*-aminobenzoic acid (PABA) and isopropanol. Using six subjects, it was observed that 20% Mink Oil with 3% Octyldimethyl PABA and 77% Isopropanol has an "SPF value equal to that of a solution using the same amount of active ingredient." This SPF value was 5.36. Details and the specific light source used were not reported.

In an earlier study, a commercial preparation of Mink Oil had an SPF value of 2.2. The undiluted solution was tested on five healthy fair-skinned persons with skin types characterized as very sensitive, sensitive, or normal (Wells Laboratories 1988b).

SUMMARY

Mink Oil, obtained from the fatty tissues of minks, is a mixture of the natural glycerides of 14–20 carbon chain fatty acids. As of January 1995, it was reportedly used in 139 cosmetic formulations as a hair conditioning agent, an occlusive skin conditioning agent, and as a surfactant. In 1984 FDA data, Mink Oil was used at concentrations of 25% and less (1995 data from the cosmetics industry indicate use at $\leq 2\%$ in two product types).

An Attenuated Total Reflectance–InfraRed Spectroscopy study found that 1 hour after application, Mink Oil was detected on the skin surface of all 5 panelists; it was detected within the stratum corneum in 2 of 5 panelists.

Mink Oil has an oral LD₅₀ of >64.0 cc/kg in albino rats.

No erythema or edema was noted after refined Mink Oil was applied for 24 hours to intact and scarified area of albino rabbits. Although reactions were noted during induction, a 50% dilution of a Mink Oil cream did not sensitize guinea pigs using the Buehler technique. In a second study, using the Magnusson-Kligman Maximization Procedure, 50% Mink Oil in petrolatum induced sensitization reactions. Mink Oil was not an ocular irritant to albino rabbits.

Clinical studies using single occlusive patches found no irritation with up to 2.8% Mink Oil (PII scores of 0.08–0.10). Transient mild to no irritation was noted in two exaggerated-use studies.

DISCUSSION

Section 1, paragraph (p) of the CIR Procedures states that “A lack of information about an ingredient shall not be sufficient to justify a determination of safety.” In accordance with Section 30(j)(2)(A) of the Procedures, the Expert Panel informed the public of its decision that the data on Mink Oil were not sufficient for determination whether the ingredient, under relevant conditions of use, was either safe or unsafe. The Panel released a Notice of Insufficient Data on March 17, 1995, outlining the data needed to assess the safety of Mink Oil. Comments were received during the 90-day public comment period. Additional data needed to make a safety assessment are:

1. 28-day dermal toxicity;
2. UV absorption data; if the ingredient absorbs in the UVA or UVB region, then photosensitization data are also needed; and
3. method of manufacture/extraction.

In the event that the above data are received and are sufficient to complete the safety assessment, the CIR Expert Panel will limit the total PCB/pesticide contamination to not more than 40 ppm with not more than 10 ppm for any specific residue. These limitations are modeled after the United States Pharmacopeia standards for Lanolin (USP 1995).

CONCLUSION

The CIR Panel concludes that the available data are insufficient to support the safety of Mink Oil for use in cosmetic products.

REFERENCES

- Bio-Toxicology Laboratories, Inc. 1974. Acute oral LD₅₀ toxicity study on emulan oil of mink. Unpublished data provided by CTFA, 3 pages.*
- Cosmetic, Toiletry, and Fragrance Association (CTFA). 1983. Mink Oil: Modified Magnusson-Kligman Guinea pig maximization test for contact sensitization. Study project: GPA-07-83 B. Unpublished data provided by CTFA. 17 pages.*
- CTFA. 1989. Clinical evaluation report: Human patch test; Mink Oil. Unpublished data provided by CTFA, 1 page.*

*Available for review from Director, Cosmetic Ingredient Review, 1101 17th Street, NW, Suite 310, Washington, DC 20036 USA.

- CTFA. 1991. Four-day mini-cum patch test results; Mink Oil. Study Ref #: 731-91. Unpublished data provided by CTFA, 1 page.*
- CTFA. 1992a. Clinical evaluation report: Human patch test; Mink Oil. Unpublished data provided by CTFA 1 page.*
- CTFA. 1992b. Irritancy potential; Mink Oil. Unpublished data provided by CTFA, 4 pages.*
- CTFA. 1993. Clinical evaluation report: Human patch test; Mink oil. Unpublished data provided by CTFA, 1 page.*
- CTFA. 1994. Evaluation of hair improvement; Mink oil. Unpublished data provided by CTFA, 3 pages.*
- CTFA. 1995. Use levels for various ingredients. Unpublished data provided by CTFA 1 page.*
- Emulan, Inc. Unknown data. Information on the stability of Mink Oil with storage and acid content of two grades of Mink Oil. Unpublished data provided by CTFA, 2 pages.*
- Food and Drug Administration (FDA). 1978. Sunscreen drug products for over-the-counter human drugs. *Federal Register* 43:38206–38269.
- FDA. 1984. *Cosmetic product formulation and frequency of use data*. [FDA database.] Washington, DC: FDA.
- FDA. 1992. Modification in Voluntary Filing of Cosmetic Product Ingredient and Cosmetic Raw Composition Statements. Final rule. *Federal Register* 57:3128–3130.
- FDA. 1995. *Frequency of use of cosmetic ingredients*. [FDA database.] Washington, DC: FDA.
- Leberco Laboratories. 1972a. Dermal irritation assay on guinea pigs: New process EmulanTM multi-refined oil of mink. Assay no. 23872. Unpublished data provided by CTFA, 2 pages.*
- Leberco Laboratories. 1972b. Dermal sensitization assay on guinea pigs: New process EmulanTM multi-refined oil of mink. Assay No. 23873. Unpublished data provided by CTFA, 4 pages.*
- Leberco Laboratories. 1972c. Ocular Irritation assay on rabbits: New process EmulanTM multi-refined oil of mink. Assay No. 23871. Unpublished data provided by CTFA, 4 pages.*
- New Jersey Laboratories/ A. A. Labs Division. 1995. Results of analysis of Mink Oil for PCB's. Assay No. 25583. Unpublished data provided by CTFA, 1 page.*
- Nikitakis, J. M., and G. N. McEwen, Jr. ed. 1990. *CTFA compendium of cosmetic ingredient composition—Descriptions I*. Washington, D.C: CTFA.
- SGS Control Services Inc. 1994. Sample identification. Unpublished data provided by CTFA, 1 page.*
- U.S. Pharmacopeia. 1995. *The United States pharmacopeia: The national formulary*. 23rd rev, 869–871. Rockville, MD: United States Pharmacopeial Convention, Inc.
- Wells Laboratories, Inc. 1979a. Report on eye irritation testing in rabbits: Multi-refined oil of mink. Lab No. H-7676. Unpublished data provided by CTFA, 2 pages.*
- Wells Laboratories, Inc. 1979b. Report on primary skin irritation on rabbits: Multi-refined oil of mink. Lab No. H-7677. Unpublished data provided by CTFA, 2 pages.*

- Wells Laboratories, Inc. 1988a. Report on a comparative study of the sun protection factor determination screen employing a mink oil mixture with octyldimethyl PABA. Lab No. L-4730A. Unpublished data provided by CTFA, 4 pages.*
- Wells Laboratories, Inc. 1988b. Report on sun protection screening test using oil of mink. Lab No. L-4730. Unpublished data provided by CTFA, 2 pages.*
- Wells Laboratories, Inc. 1990a. Report on guinea pig sensitization test: Emulan sable cream. Lab No. W90-992. Unpublished data provided by CTFA, 10 pages.*
- Wells Laboratories, Inc. 1990b. Comedogenic assay: Emulan sable cream. Lab No. W90-994. Unpublished data provided by CTFA, 6 pages.*
- Wells Laboratories, Inc. 1990c. Report on eye irritation testing in rabbits: Emulan sable cream. Lab No. W90-990. Unpublished data provided by CTFA, 3 pages.*
- Wells Laboratories, Inc. 1990d. Report on primary skin irritation in rabbits: Emulan sable cream. Lab No. W90-991. Unpublished data provided by CTFA, 3 pages.*
- Wells Laboratories, Inc. 1990e. Report on acute oral toxicity: Emulan sable cream. Lab No. W90-993. Unpublished data provided by CTFA, 2 pages.*
- Wenninger, J. A., and G. N. McEwen, Jr. ed. 1997. *International cosmetic ingredient dictionary and handbook*. 7th ed, 822. Washington, DC: CTFA.
- Xienta Institute for Skin Research. 1988. Infrared penetration study for Emulan Incorporated. Study No. BM-01-88. Unpublished data provided by CTFA, 19 pages.*

Final Amended Report on the Safety Assessment of Mink Oil¹

Mink Oil, obtained from the fatty tissues of minks, is a mixture of the natural glycerides of 14 to 20 carbon chain fatty acids. There are 100 current reported uses as a hair-conditioning agent, an occlusive skin-conditioning agent, and as a surfactant; up to a maximum concentration of 3%. Mink Oil is manufactured by harvesting animal hides and scraping the fat layer from the hide. It is rendered and refined using high temperature processes (230°F to 240°F) and saponification to reduce free fatty acids. Analyses demonstrate that Mink Oil can be substantially free of impurities, including pesticides. Mink Oil does not absorb significant UVA or UVB radiation. In a clinical test of skin penetration, 1 h after application, Mink Oil was detected on the skin surface of all five panelists; it was detected within the stratum corneum in 2/5 panelists. Mink Oil has an oral LD₅₀ of >64.0 cc/kg in albino rats. No erythema or edema was noted after refined Mink Oil was applied for 24 h to intact and scarified area of albino rabbits. A 50% dilution of a Mink Oil cream did not sensitize guinea pigs in a maximization test. Mink Oil was not an ocular irritant to albino rabbits. Clinical studies using single occlusive patches found no irritation with up to 2.8% Mink Oil, although transient mild to no irritation was noted in two exaggerated-use studies. Mink Oil is used in aerosols and sprays. Although there are no inhalation toxicity data available on Mink Oil, the available data on particle sizes of cosmetic aerosols and sprays indicates diameters more than an order of magnitude larger than the diameter of respirable particles. Most of the glycerides in Mink Oil are triglycerides (glyceryl triesters), the safety of which has been substantiated in previous safety assessments; e.g., dermal absorption is nil to slight; there is little or no acute, subchronic, or chronic oral toxicity; dermal application was not associated with significant irritation or sensitization; ocular exposures were, at most, mildly irritating; most of the genotoxicity test systems are negative; use as vehicles in carcinogenicity testing of other chemicals has produced no adverse reaction; and clinical tests produce no irritation or sensitization reactions—but, they may enhance the of penetration of other chemicals. Formulators should be aware of the possible penetration-enhancing properties of Mink Oil. Although pesticide residues have been analyzed and found to be below levels of detection, the Panel is concerned that the available data suggesting the absence of pesticide residues in Mink Oil are limited. The Panel advised the industry that the total polychlorinated biphenyl (PCB)/pesticide contamination should be limited to not more than 40 ppm, with not more than 10 ppm for any specific residue.

Received 15 April 2005; accepted 12 July 2005.

Address correspondence to F. Alan Andersen, Director, Cosmetic Ingredient Review, 1101 17th Street, NW, Suite 310, Washington, DC 20036, USA. E-mail: andersena@cir-safety.org

¹Reviewed by the Cosmetic Ingredient Review Expert Panel.

INTRODUCTION

This amended safety assessment updates and supersedes an earlier Cosmetic Ingredient Review (CIR) safety assessment of Mink Oil (CIR 1995). New information is included on the manufacturing process; batch analysis data; pesticide residue analysis; updated concentration of use; and ultraviolet (UV) absorption spectra. Information provided by the Cosmetic, Toiletry, and Fragrance Association (CTFA) indicated that most glycerides in Mink Oil are triglycerides (CTFA 2001a). The safety of triglycerides in cosmetics was assessed by CIR, with the conclusion that these ingredients are safe as used in cosmetics (CIR 1999). In addition, CIR has published a safety assessment of Oleic, Lauric, Palmitic, Myristic, and Stearic Acids (Elder 1987). Uses of both the triglycerides group and fatty acid group of ingredients in cosmetics at maximum concentrations ranging from 25% to 50% were determined to be safe. On the basis of the available data the Panel issued this amended safety assessment.

CHEMISTRY

Method of Manufacture

Mink Oil is a mixture of the natural glycerides of 14 to 20 carbon chain fatty acids obtained from the subdermal fatty tissues of the mink (Pepe et al. 2002).

EMULAN, Inc. (1990) uses Northern Ranch Mink as the source for Mink Oil. Animals are raised in cages and kept free from disease and are harvested typically in late fall/early winter.

The fat is scraped from the hides of the mink and frozen until it is rendered and refined. The fat is rendered at temperatures ranging from 220°F to 240°F for a minimum of 5 h, after which the saturated fat is separated from the unsaturated fat using filtration. A saponification process then further separates the free fatty acids; i.e., the saponified free fatty acids form soap at the bottom. The oil is further bleached and clarified using diatomaceous earth, again at the high temperature described above. Impurities such as any remaining soap or any proteinaceous impurity (e.g., collagen) are removed at this stage (EMULAN, Inc. 1990; Complete Analysis Laboratories, Inc. 1996; CTFA 1999a).

Chemical Composition

According to the *Compendium of Cosmetic Ingredient Composition* (Nikitakis and McEwen 1990), the oil is a mixture of the natural glycerides of 14 to 20 carbon chain fatty acids; approximately 75% of the composition is 16 to 18 carbon chain

TABLE 1

Fatty acid composition of two Mink Oils (CTFA 1994)

Fatty acid	Carbon chain length/ number of unsaturated C:C bonds	Mink Wax Oil (as %)	Mink Crude Oil (as %)
Lauric	12/0	0.1	0.1
Myristic	14/0	4.0	3.5
Myristoleic	14/1	0.7	0.9
Pentadecanoic	15/0	0.2	0.1
Palmitic	16/0	28.0	17.2
Palmitic	16/1	13.3	17.0
Heptadecanoic	17/0	0.5	0.4
Heptadecanoic	17/1	0.6	0.5
Stearic	18/0	4.7	2.5
Oleic	18/1	35.3	40.9
Linoleic	18/2	10.6	15.0
Linolenic	18/3	0.4	0.6
Eicosanoic	20/0	0.1	—
Eicolenic	20/1	0.7	0.6
Total fatty acid		99.2	99.3
Unsaturated		61.6	75.5
Saturated		37.6	23.8

fatty acid glycerides. The detailed fatty acid composition of two forms of Mink Oil (Mink Wax Oil and Mink Crude Oil) provided by one supplier is given in Table 1 (CTFA 1994).

Physical and Chemical Properties

Mink Oil is a pale yellow liquid with a mild characteristic odor. It is soluble in chloroform, carbon tetrachloride, ether, benzene, acetone, and isopropanol and is insoluble in water. The refractive index at 25°C is between 1.4665 and 1.4675. The freezing point is 5°C maximum; the saponification value is 190 to 220; the iodine value is 80 to 90 (Nikitakis and McEwen 1990). SGS Control Services Inc. (1994) reported a specific gravity of 0.91437 and a refractive index at 40°C of 1.4623. Jennings Laboratories, Inc. (1999) reported a specific gravity at 25°C of 0.9110 and a refractive index at 25°C of 1.4680. One supplier reported that because of its iodine value, Mink Oil “exhibits very low oxidation and degradation” (CTFA 1994).

UV Absorption

Mink Oil prepared according to the manufacturing process described above has significant absorbance only in the UVC region. Absorption maxima in a 0.5% sample occur at 212 nm (2.59700 absorption units), 270 nm (0.55858 absorption units), and a small absorption at 316 nm (0.07776 absorption units). A 0.05% solution had no detectable absorption at 316 nm (CTFA 1999b).

Impurities

One analysis reported 0.0628% moisture, 0.0012% unsaponifiable matter, and 0.0002% insolubles (Jennings Laboratories, Inc. 1999).

Analysis of one sample of Mink Oil found no detectable levels of the following tradename polychlorinated biphenyls (PCBs): Arochlor 1016, 1221, 1232, 1242, 1248, 1254, and 1260. The assay had a minimum detection limit of 0.5 µg/L (New Jersey Laboratories/A.A. Labs Division 1995).

Pesticide analyses of two samples produced the results shown in Table 2. No explanation was provided for the different detection limits for the two sample analyses, although it was reported that they were done at different times (CTFA 2001a).

The United States Pharmacopeia (USP) standards for lanolin, for comparison purposes, limit PCB/pesticide contamination to not more than 40 ppm, with not more than 10 ppm for any specific residue (U.S. Pharmacopeia 1995).

COSMETIC USE

Mink Oil is used in formulations as a hair conditioning agent, an occlusive skin conditioning agent, and a surfactant (Pepe et al. 2002). Based on information received from industry, the Food and Drug Administration (FDA) lists 100 current uses of Mink Oil (FDA 2001). The uses of Mink Oil as a function of product type are given in Table 3. The product types and the total number of each product type reported to FDA (FDA 2001) are given, with the number of formulations of that product type that were reported to contain Mink Oil; e.g., of 187 mascara products, only one contains Mink Oil. Current concentration of use data provided by industry (CTFA 1999c, 2001b) as a function of product type are also included in Table 3. These data indicate that Mink Oil is reportedly used in a given product category, but the concentration of use is not available. In other cases, information regarding use concentration for specific product categories is provided, but the number of such products is not known.

Mink Oil is used in products that are or may be found in sprays or aerosols. Cosmetic product sprays and aerosols produce large particle sizes. For example, aerosol hair sprays have particles of 60 to 80 µ diameter, with <1% below 10 µ, and pump hair sprays have particle diameters of ≥80 µ (Bower 1999). These particle sizes are larger than the mean aerodynamic diameter of respirable particles, which is 4.25 ± 1.5 µ (Jensen and O'Brien 1993).

The European Commission (EC) has not included Mink Oil in the list of substances which cosmetic products must not contain or must not contain except subject to restrictions and conditions laid down (EC 2002). In the current Japanese Ministry of Health, Labor and Welfare (MHLW) regulations, Mink Oil is not included on a negative list (MHLW 2000a), on a list of ingredients for which there are restrictions to use in cosmetics (MHLW 2000b), or on a list of quasi-drugs for which listing is required (MHLW 2000c).

TABLE 2
Pesticide residues in Mink Oil (CTFA 2001b)

Pesticide	8/7/96 sample		10/7/96 sample	
	Level in Mink Oil (mg/kg)	Min. detectable level (mg/kg)	Level in Mink Oil (mg/kg)	Min. detectable level (mg/kg)
A-BHC	U*	0.020	U	0.020
B-Bhc	U	0.020	U	0.020
G-BHC (Lindane)	U	0.020	0.11	0.020
D-BHC	U	0.020	U	0.020
Heptachlor	U	0.020	U	0.020
Aldrin	U	0.020	U	0.020
Heptachlor Epoxide	U	0.020	U	0.020
Endosulfan I	U	0.020	U	0.020
A-Chlordane	U	0.020	U	0.020
G-Chlordane	U	0.020	U	0.020
Dieldrin	U	0.020	U	0.020
4,4'-DDE	0.27	0.020	U	0.020
Endrin	U	0.020	U	0.020
Endosulfan II	U	0.020	U	0.040
4,4'-DDD	U	0.020	U	0.040
Endrin Aldehyde	U	0.020	U	0.040
Endosulfan Sulfate	U	0.020	U	0.040
4,4'-DDT	U	0.020	U	0.040
Endrin Ketone	U	0.020	U	0.040
Methoxychlor	U	0.020	U	0.020
Toxaphene	U	0.020	U	1.00

*U = undetectable.

ABSORPTION, DISTRIBUTION, METABOLISM, AND EXCRETION

Absorption

Xienta Institute for Skin Research (1988) conducted an attenuated total reflectance–infrared spectroscopy study (ATR-IR) to determine the penetration of a commercial preparation of Mink Oil into human stratum corneum. Five healthy Caucasian females were instructed to refrain from applying any topical agents to their forearms 24 h prior to the study. A site on one forearm of each panelist was treated with 50 μ l of the test substance. The same site on the other forearm was not treated and served as control. Both sites were scanned prior to product application. Scanning consisted of placing the subject's forearm directly onto a germanium plate and conducting an IR spectra for wavelengths between 800 and 4000 cm^{-1} . The germanium plate served as an internal reflection crystal. One hour after application, the sites were again scanned. This was followed by tape stripping and scans after 5, 10, and 15 tape strippings. The same scans were conducted on the untreated sites.

Based on the IR spectra of the test substance, wavelengths of 1746, 1466, and 1163 cm^{-1} were chosen for analysis. Presence

of Mink Oil in each of the four scans (surface, 5, 10, 15 layers all done at 1 h post treatment) was defined as a greater absorbance value at the treated site versus the control. For each of the four scans, there were 15 absorbance pairs (a pair consisting of an absorbance value for the control and treated sites). Mink Oil was detected at the surface scan of all panelists. It was detected within 5 layers in 4/5 panelists, 10 layers in 2/5 panelists, and 15 layers in 2/5 panelists. The authors concluded that, after 1 h, Mink Oil remains on the skin surface (Xienta Institute for Skin Research 1988).

Distribution, Metabolism, and Excretion

No studies were available describing the distribution, metabolism, or excretion of Mink Oil.

ANIMAL TOXICOLOGY

Acute Oral Toxicity

A commercial preparation of Mink Oil had an oral LD_{50} of >64.0 cc/kg in albino rats (Bio-Toxicology Laboratories, Inc. 1974). In another study, a Mink Oil cream had an oral LD_{50} of >5.0 ml/kg in rats (Wells Laboratories, Inc. 1990e).

TABLE 3
Frequency of use of Mink Oil

Product category (number of formulations in category) (FDA 2001)	Number of formulations containing Mink Oil (FDA, 2001)	Concentration of use (CTFA 1999c, 2001) (%)
Baby lotions, oils, powders, and creams	—	1
Other bath preparations (193)	1	0.1
Eye shadow	—	0.1
Eye lotion (23)	1	—
Mascara (187)	1	0.3–1
Other fragrance preparations	—	1
Hair conditioner (630)	14	3
Hair spray—aerosol fixatives (267)	3	—
Hair straighteners (63)	4	—
Permanent waves (211)	1	—
Rinses—noncoloring	—	3
Shampoos—noncoloring (851)	4	—
Tonics, dressings, and other hair- grooming aids (577)	9	—
Other hair preparations (276)	4	—
Blushers—all types	—	0.1
Face powders (301)	1	0.1
Lipstick (942)	12	0.001
Makeup bases	—	0.5
Shaving cream (133)	5	—
Skin cleansing products (733)	7	—
Face and neck creams, lotions, powders and sprays—excluding shaving products (304)	2	0.2–0.5
Body and hand creams, lotions, powders and sprays—excluding shaving products (827)	10	0.1
Moisturizing (881)	8	0.2
Night products (200)	5	—
Paste masks—mud packs (269)	1	—
Other skin care preparations (715)	6	—
Suntan gels, creams, liquids (131)	1	0.3
Other suntan preparations	—	2
2001 total uses/ranges for Mink Oil	100	0.001–3

Dermal Irritation

Patches containing 0.5 ml of a commercial preparation of refined Mink Oil were applied to intact and scarified areas of guinea pig skin. The treatment areas had earlier been clipped free of hair. Animals were immobilized for a 24-h period; the sites were evaluated at the end of the treatment period and 24 h afterwards. Scores were made according to the Draize scale. No erythema or edema was noted (Leberco Laboratories 1972a).

Patches containing 0.5 ml of refined Mink Oil (neat) were applied to intact and abraded skin of six rabbits (strain not specified). No irritation was noted at observations made 24 and 72 h after application (Wells Laboratories, Inc. 1979b). Similar re-

sults were reported by Wells Laboratories, Inc. (1990d) when a cream containing Mink Oil was tested on three New Zealand white rabbits using the same protocol.

Dermal Sensitization

Over a period of 20 days, eight male guinea pigs received 10 subcutaneous injections of a 0.1% suspension of refined Mink Oil; the injections were administered to the clipped back and sides. The test material was suspended in physiological saline and was administered in 0.1 ml doses except for the first injection, which contained 0.05 ml. Two weeks after the final induction injection, a test injection of 0.05 ml was

administered. Reaction areas were measured 24 h after each injection. Irritation reactions were noted throughout the induction period. The reaction on challenge was less than the irritant reaction seen during induction and the authors concluded that the test material was not a sensitizer (Leberco Laboratories 1972b).

CTFA (1983) reported a study in which female Dunkin-Hartley guinea pigs were treated with Mink Oil using the Magnusson-Kligman maximization procedure: induction, dose range, booster, and challenge. During induction, sites on the upper back were injected intradermally with 50% Freund's complete adjuvant, 5% Mink Oil in propylene glycol, and 5% Mink Oil in 50% Freund's complete adjuvant. Control animals were injected with complete adjuvant, propylene glycol, and 1:1 propylene glycol in complete adjuvant. During the dose range phase, 10 animals were tested with 24 h occlusive patches containing 25%, 50%, and 100% Mink Oil. Sites were evaluated for irritation at 24, 48, and 72 h after patch application. Two of the animals reacted to the 25% concentration with a response graded as "barely perceptible"; by 72 h the condition cleared. These two animals also reacted to the higher concentrations of Mink Oil; the reactions cleared by the 72-h observation. Another four had "barely perceptible" or "mild" reactions to the 50% concentration. These reactions also cleared by 72 h and three of these four animals did not react to the 100% concentration. Three animals reacted for the first time to the 100% concentration with a "barely perceptible" response (two also had desquamation); these conditions cleared by 72 h.

During the booster phase (1 week after induction injections) undiluted Mink Oil was topically applied. Sodium lauryl sulfate (SLS) at 5% was applied to the induction site 24 h before the booster. Control animals were pretreated with SLS before receiving a booster of undiluted petrolatum; positive controls were treated with 25% phenylacetaldehyde without SLS pretreatment. During the booster phase, all test and control animals were wrapped occlusively for 48 h. Two weeks following the booster, test animals were challenged with a 24-h occlusive patch of 25% and 50% Mink Oil in petrolatum. Reactions were scored at 48 and 72 h after patch application. There were no reactions to 25% Mink Oil noted in the ten animals at the 48-h reading; one animal had a "barely perceptible" reaction at 72 h. Four animals had a "barely perceptible" and one had a "mild" response to 50% Mink Oil at the 48-h reading. At the 72-h reading, erythema was noted in 7 of 10 animals, which ranged from "barely perceptible" (4 animals) to "desquamation" (2 animals) and "mild with desquamation" (1 animal). Mink Oil at 50% in petrolatum was considered to produce a weak allergenic response (CTFA 1983).

The Buehler Technique was used by Wells Laboratories (1990a) in a guinea pig sensitization assay. During induction nine occlusive patches containing 0.5 ml of a 50% dilution (in corn oil) of a Mink Oil cream was applied to the clipped back and flank of 10 animals. The patches remained for 6 h of contact and readings were made at the time of patch removal and

at the end of 24 h. Positive controls were treated with dinitrochlorobenzene (DNCB). After a 2-week nontreatment period animals were challenged with two identical applications of the test material. One patch was applied at the site of induction, and the other on a previously unexposed site. Challenge patches were applied for 6 h of contact and observations were made at the time of patch removal and at the end of 24 h. A reading of "very slight" erythema was noted in all animals during induction; "slight" erythema and "slight" edema were noted in positive controls. During challenge, no irritation was noted in animals of the Mink Oil group; marked erythema and very slight edema was noted in animals of the positive-control group.

Ocular Irritation

A commercial preparation of refined Mink Oil (0.1 ml) was instilled into the right conjunctival sac of three albino rabbits. The untreated left eye served as control. Eyes were evaluated every 24 h for 4 days and then again on the 7th day. Scores were made according to the Draize scale. No irritation was noted (score 0) (Leberco Laboratories 1972c).

Another ocular irritation test was conducted using three rabbits and the above described protocol (Wells Laboratories 1979a). Undiluted Mink Oil (commercial preparation, dose not reported) did not produce any observable irritation.

A Mink Oil cream (0.1 ml) was applied to one eye of each of three New Zealand white rabbits. The contralateral eye was not treated and served as control. Eyes were unwashed for 24 h and observations made 24, 48, and 72 h after treatment. No irritation was noted (Draize score 0.01) (Wells Laboratories, Inc. 1990c).

Comedogenicity

A Mink Oil cream was considered to be noncomedogenic following repetitive applications (10 applications in 2 consecutive weeks) to the external ear canal of six albino rabbits. Observations noted include "minimal to mild" hyperemia and "very minimal" hyperkeratosis in the treated areas of all animals towards the end of the treatment period (Wells Laboratories, Inc. 1990b).

CLINICAL ASSESSMENT OF SAFETY

Dermal Irritation and Sensitization

Twenty panelists received a single 24-h occlusive patch containing a 25% aqueous dilution of a hair rinse containing 3% Mink Oil (effective concentration of Mink Oil tested: 0.75%). No reactions were noted in 16 panelists, reactions of \pm (the mildest non-zero score possible) were noted in four panelists. The primary irritation index (PII) was 0.10 out of a maximum possible score of 4.0 (CTFA 1989). Identical results were reported in a second patch test study in which a formulation containing 2.8% Mink Oil was tested at full strength in 20 panelists (CTFA 1992a). A third human patch test study reported a PII of

0.08 after 19 panelists were tested with a (full-strength) formulation containing 2.8% Mink Oil (CTFA 1993).

A 4-day minicumulative patch test of a body spray containing 1% Mink Oil had a PII of 0.50. Of 20 panelists, 5 had no response, 12 had a “barely perceptible” response, 2 had a “mild” response, and 1 had a “moderate” response (CTFA 1991).

In an exaggerated-use study, 19 female panelists applied to the lower arm, a rinse-off hair product containing 2.8% Mink Oil once a day for 5 days with no rinse off. A commercially available rinse-off hair product was applied to a different site to serve as a reference. Four subjects experienced some transient mild to moderate erythema (CTFA 1992b). In a second study using the same protocol, no clinical responses were observed in 24 females who also tested a rinse-off hair product containing 2.8% Mink Oil (CTFA 1994).

Sun-Protection Factor

Wells Laboratories (1988a, 1988b) tested the sunscreens ability of Mink Oil. In six subjects, 20% Mink Oil with 3% Octyl Dimethyl PABA and 77% Isopropanol had a sun-protection factor (SPF) value of 5.36. In five subjects, a preparation containing Mink Oil had an SPF value of 2.2. Further study details and the specific light source used were not reported.

SUMMARY

Mink Oil, obtained from the fatty tissues of minks, is a mixture of the natural glycerides of 14 to 20 carbon chain fatty acids. In 2001, it was reportedly used in 100 cosmetic formulations as a hair-conditioning agent, an occlusive skin-conditioning agent, and as a surfactant at concentrations no greater than 3%.

Mink Oil is manufactured by harvesting animal hides and scraping the fat layer from the hide. It is rendered and refined using high temperature processes (230°F to 240°F) and saponification to reduce free fatty acids.

Most of the glycerides in Mink Oil are triglycerides, the safety of which has substantially assessed in previous safety assessments; e.g., dermal absorption is nil to slight; there is little or no acute, subchronic, or chronic oral toxicity; dermal application was not associated with significant irritation or sensitization; ocular exposures were, at most, mildly irritating; most of the genotoxicity test systems are negative; use as vehicles in carcinogenicity testing of other chemicals has produced no adverse reaction; and clinical tests produce no irritation or sensitization reactions—but, they may enhance the penetration of other chemicals.

Analyses demonstrate that Mink Oil can be substantially free of impurities, including pesticides. Mink Oil does not absorb significant UVA or UVB radiation.

An attenuated total reflectance–infrared spectroscopy study found that 1-h after application, Mink Oil was detected on the skin surface of all five panelists; it was detected within the stratum corneum in 2/5 panelists.

Mink Oil has an oral LD₅₀ of >64.0 cc/kg in albino rats.

No erythema or edema was noted after refined Mink Oil was applied for 24 h to intact and scarified area of albino rabbits.

Although reactions were noted during induction, a 50% dilution of a Mink Oil cream did not sensitize guinea pigs using the Buehler technique. In a second study, using the Magnusson-Kligman maximization procedure, 50% Mink Oil in petrolatum-induced sensitization reactions.

Mink Oil was not an ocular irritant to albino rabbits.

Clinical studies using single occlusive patches found no irritation with up to 2.8% Mink Oil (PII scores of 0.08 to 0.10). Transient mild to no irritation was noted in two exaggerated-use studies.

DISCUSSION

The CIR Expert Panel had previously issued a final safety assessment of Mink Oil with an insufficient data conclusion. The Panel has now considered information on the manufacturing process; batch analysis data; pesticide residue analysis; updated concentration of use; and UV absorption spectra newly provided by industry.

The Panel noted that Mink Oil is used in aerosols and sprays. Although there are no inhalation toxicity data available on Mink Oil, the available data on particle sizes of cosmetic aerosols and sprays indicate diameters more than an order of magnitude larger than the diameter of respirable particles. Inhalation, therefore, is not believed to produce significant exposure of lung tissue.

It was noted that most of the glycerides in Mink Oil are triglycerides, and the Panel agreed that the safety test data in the triglycerides safety assessment are relevant to Mink Oil. The triglycerides data demonstrate, among other things, that dermal absorption of triglycerides is nil to slight; there is little or no acute, subchronic, or chronic oral toxicity; dermal application is not associated with significant irritation or sensitization; ocular exposures are, at most, mildly irritating; most genotoxicity test system results are negative; use as vehicles in carcinogenicity testing of other chemicals has produced no adverse reaction; and clinical tests produce no irritation or sensitization reactions. Because triglycerides may enhance the penetration of other chemicals, it has been recommended that care be exercised in using them in cosmetic products. That same caution should be applied to Mink Oil.

Although pesticide residues have been analyzed and found to be below levels of detection, the Panel is concerned that the available data suggesting the absence of pesticide residues in Mink Oil are limited. The Panel advised the industry that the total PCB/pesticide contamination should be limited to not more than 40 ppm, with not more than 10 ppm for any specific residue. As with any animal-derived ingredient, Mink Oil should be free of detectable pathogenic viruses or infectious agents.

The CIR Expert panel recognizes that Mink Oil is reportedly used in some product categories, but that the concentration of

use is not available. In other cases, information regarding use concentration for specific product categories is provided, but the number of such products is not known. Although there are gaps in knowledge about product use, the overall information available on the types of products in which Mink Oil is used and at what concentration indicate a pattern of use. Within this overall pattern of use, the Expert Panel considers Mink Oil to be safe.

CONCLUSION

Based on the available information, the CIR Expert Panel concludes that Mink Oil is safe as a cosmetic ingredient in the practices of use and concentration described in this safety assessment.

REFERENCES

- Bio-Toxicology Laboratories, Inc. 1974. Acute oral LD₅₀ toxicity study on Emulan Oil of Mink. Unpublished data submitted by CTFA on August 5, 1994. (3 pages.)²
- Bower, D. 1999. Unpublished information on hair spray particle sizes. Minutes of the September 9, 1999 meeting of the CIR Expert Panel.²
- Complete Analysis Laboratories, Inc. 1996. Analytical data report for EMULAN, Inc. Unpublished data submitted by CTFA, September 14, 2001. (6 pages.)²
- Cosmetic Ingredient Review (CIR). 1995. Final report on the safety assessment of Mink Oil. December, 1995.³
- CIR. 1999. Final report on the safety assessment of Trilaurin, Triarachidin, Tribehenin, Tricaprin, Tricaprylin, Trierucin, Triheptanoin, Triheptylundecanoin, Triisononanoin, Triisopalmitin, Triisostearin, Trilinolein, Trimyristin, Trioctanoin, Triolein, Tripalmitin, Tripalmitolein, Triricinolein, Tristearin, Triundecanoin, Glyceryl Triacetyl Hydroxystearate, Glyceryl Triacetyl Ricinoleate, and Glyceryl Stearate Diacetate. March 4, 1999.²
- Cosmetic, Toiletry, and Fragrance Association (CTFA). 1983. Mink Oil: Modified Magnusson-Kligman guinea pig maximization test for contact sensitization. Study project: GPA-07-83 B. Unpublished data submitted by CTFA, June 6, 1995. (17 pages.)²
- CTFA. 1989. Clinical evaluation report: Human patch test; Mink Oil. Unpublished data submitted by CTFA, June 6, 1995. (1 page.)²
- CTFA. 1991. Four-day mini-cum patch test results; Mink Oil. Study ref. no. 731-91. Unpublished data submitted by CTFA, June 6, 1995. (1 page.)²
- CTFA. 1992a. Clinical evaluation report: Human patch test; Mink Oil. Unpublished data submitted by CTFA, June 6, 1995. (1 page.)²
- CTFA. 1992b. Irritancy potential; Mink Oil. Unpublished data submitted by CTFA, June 6, 1995. (4 pages.)²
- CTFA. 1993. Clinical evaluation report: Human patch test; Mink oil. Unpublished data submitted by CTFA, June 6, 1995. (1 page.)²
- CTFA. 1994. Evaluation of hair improvement; Mink Oil. Unpublished data submitted by CTFA, June 6, 1995. (3 pages.)²
- CTFA. 1999a. Summary of manufacturing method for Mink Oil (information obtained from two suppliers in 1999). Unpublished data submitted by CTFA. (1 page.)²
- CTFA. 1999b. UV absorption data. Unpublished data submitted by CTFA. (6 pages.)²
- CTFA. 1999c. Use levels for Mink Oil. Updated February 11, 1999. Unpublished data submitted by CTFA. (1 page.)²
- CTFA. 2000. Mink Oil—Need for a 28-day dermal toxicity study (7/5/2000). Unpublished data submitted by CTFA. (1 page.)²
- CTFA. 2001a. Industry data regarding Mink Oil: Processing and pesticide analysis. Unpublished data submitted by CTFA, 9/14/01. (8 pages.)²
- CTFA. 2001b. Use levels for Mink Oil. Updated December 12, 2001. Unpublished data submitted by CTFA. (1 page.)²
- Elder, R.L. ed. 1987. Final report on the safety assessment of Oleic, Lauric, Palmitic, Myristic, and Stearic Acids. *J. Am. Coll. Toxicol.* 6:321–401.
- EMULAN, Inc. 1990. Manufacturing process. Unpublished data provided by CTFA, September 14, 2001. (1 page.)²
- European Commission (EC). 2002. Twenty-sixth Commission Directive 2002/34/EC of 15 April 2002 adapting to technical progress Annexes II, III, and VII to Council Directive 76/768/EEC. Brussels: EC.
- Food and Drug Administration (FDA). Sunscreen drug products for over-the-counter human drugs. *Federal Register*, August 25, 1978.
- FDA, 2001. Frequency of use of cosmetic ingredients. *FDA database*. Washington, DC: FDA.
- Jennings Laboratories, Inc. 1999. Certificate of analysis: Initially processed Mink Oil. Unpublished data submitted by CTFA. (2 pages.)²
- Jenson, P. A., and D. O'Brien. 1993. Industrial hygiene. In *Aerosol Measurement. Principles, Techniques, and Applications*, ed. K. Willeke and P. A. Baron, New York: John Wiley and Sons. 538–540.
- Leberco Laboratories 1972a. Dermal irritation assay on guinea pigs: New process Emulan™ Multi-Refined Oil of Mink. Assay No. 23872. Unpublished data submitted by CTFA, August 5, 1994. (2 pages.)²
- Leberco Laboratories 1972b. Dermal sensitization assay on guinea pigs: New process Emulan™ Multi-Refined Oil of Mink. Assay No. 23873. Unpublished data submitted by CTFA, August 5, 1994. (4 pages.)²
- Leberco Laboratories 1972c. Ocular irritation assay on rabbits: New process Emulan™ Multi-Refined Oil of Mink. Assay No. 23871. Unpublished data submitted by CTFA, August 5, 1994. (4 pages.)²
- Ministry of Health, Labor and Welfare (MHLW). 2000a. Unofficial translation of MHW Ordinance No. 331, Attached Table 1 [Negative List]. Ministry of Health Labor and Welfare, Pharmaceutical and Medical Safety Bureau, Inspection and Guidance Division, 2-2,1-chome, Kasumigaseki, Chiyoda-ku, Tokyo 100-8045, Japan.
- MHLW. 2000b. Unofficial translation of MHW Ordinance No. 331, Attached Table 2 [Restricted List]. Ministry of Health Labor and Welfare, Pharmaceutical and Medical Safety Bureau, Inspection and Guidance Division, 2-2,1-chome, Kasumigaseki, Chiyoda-ku, Tokyo 100-8045, Japan.
- MHLW. 2000c. Unofficial translation of MHW Ordinance No. 331, Appendix 4 [Ingredients of quasi-drugs] Ministry of Health Labor and Welfare, Pharmaceutical and Medical Safety Bureau, Inspection and Guidance Division, 2-2,1-chome, Kasumigaseki, Chiyoda-ku, Tokyo 100-8045, Japan.
- New Jersey Laboratories/A.A. Labs Division. 1995. Results of analysis of Mink Oil for PCB's. Assay No. 25583. Unpublished data submitted by CTFA, June 6, 1995. (1 page.)²
- Nikitakis, J. M., and G. N., McEwen, Jr., eds. 1990. *CTFA compendium of cosmetic ingredient composition—Descriptions I*. Washington, DC: Cosmetic, Toiletry, and Fragrance Association (CTFA).
- Pepe, R. C., J. A. Wenninger, and G. N. McEwen, Jr., eds. 2002. *International Cosmetic Ingredient Dictionary and Handbook*, 9th ed., 997. Washington, DC: CTFA.
- SGS Control Services Inc. 1994. Sample Identification. Unpublished data submitted by CTFA, August 5, 1994. (1 page.)¹
- U.S. Pharmacopeia. 1995. *The United States Pharmacopeia: The National Formulary*, 23rd rev., 869–871. Rockville, MD: United States Pharmacopeial Convention.
- Wells Laboratories, Inc. 1979a. Report on eye irritation testing in rabbits: Multi-refined Oil of Mink. Lab No. H-7676. Unpublished data submitted by CTFA, August 5, 1994. (2 pages.)²
- Wells Laboratories, Inc. 1979b. Report on primary skin irritation on rabbits: Multi-refined Oil of Mink. Lab No. H-7677. Unpublished data submitted by CTFA, June 6, 1995. (2 pages.)²

²Available from the Director, Cosmetic Ingredient Review, 1101 17th Street NW, Suite 310, Washington, DC 20777, USA.

³Not available—superseded by this report.

- Wells Laboratories, Inc. 1988a. Report on a comparative study of the sun protection factor determination screen employing a Mink Oil mixture with Octyl Dimethyl PABA. Lab No. L-4730A. Unpublished data submitted by CTFA, August 5, 1994. (4 pages.)²
- Wells Laboratories, Inc. 1988b. Report on sun protection screening test using Oil of Mink. Lab No. L-4730. Unpublished data submitted by CTFA, August 5, 1994. (2 pages.)²
- Wells Laboratories, Inc. 1990a. Report on guinea pig sensitization test: Emulan Sable Cream. Lab No. W90-992. Unpublished data submitted by CTFA, June 6, 1995. (10 pages.)²
- Wells Laboratories, Inc. 1990b. Comedogenic assay: Emulan Sable Cream. Lab No. W90-994. Unpublished data submitted by CTFA, June 6, 1995. (6 pages.)²
- Wells Laboratories, Inc. 1990c. Report on eye irritation testing in rabbits: Emulan Sable Cream. Lab No. W90-990. Unpublished data submitted by CTFA, June 6, 1995. (3 pages.)²
- Wells Laboratories, Inc. 1990d. Report on primary skin irritation in rabbits: Emulan Sable Cream. Lab No. W90-991. Unpublished data submitted by CTFA, June 6, 1995. (3 pages.)²
- Wells Laboratories, Inc. 1990e. Report on acute oral toxicity: Emulan Sable Cream. Lab No. W90-993. Unpublished data submitted by CTFA, June 6, 1995. (2 pages.)²
- Xienta Institute for Skin Research. 1988. Infrared penetration study for Emulan Incorporated. Study No. BM-01-88. Unpublished data submitted by CTFA, August 5, 1994. (19 pages.)²