
Safety Assessment of PEGylated Alkyl Glycerides as Used in Cosmetics

Status: Draft Final Report for Panel Review
Release Date: November 14, 2014
Panel Meeting Date: December 8-9, 2014

The 2014 Cosmetic Ingredient Review Expert Panel members are: Chair, Wilma F. Bergfeld, M.D., F.A.C.P.; Donald V. Belsito, M.D.; Ronald A. Hill, Ph.D.; Curtis D. Klaassen, Ph.D.; Daniel C. Liebler, Ph.D.; James G. Marks, Jr., M.D., Ronald C. Shank, Ph.D.; Thomas J. Slaga, Ph.D.; and Paul W. Snyder, D.V.M., Ph.D. The CIR Director is Lillian J. Gill, D.P.A. This safety assessment was prepared by Monice M. Fiume, Assistant Director/Senior Scientific Analyst and Bart Heldreth, Ph.D., Chemist.



Commitment & Credibility since 1976

Memorandum

To: CIR Expert Panel Members and Liaisons
From: Monice M. Fiume *MMF*
Assistant Director/Senior Scientific Analyst
Date: November 14, 2014
Subject: Safety Assessment of PEGylated Alkyl Glycerides as Used in Cosmetics

Enclosed is the Draft Final Report on the Safety Assessment of PEGylated Alkyl Glycerides as Used in Cosmetics. (It is identified as *PEGgly122014rep* in the pdf document.) At the September meeting, the Panel determined that it was appropriate to include the PEG glyceryl cocoates in this report and concluded that the 60 PEGylated alkyl glycerides included in this assessment are safe in the present practices of use and concentration in cosmetics when formulated to be non-irritating.


A summary of a developmental toxicity study on PEG-8 caprylic/capric glycerides was received and added to the report (*PEGgly122014data*). Comments also were received from the Council and have been addressed (*PEGgly122014PCPC_1*; *PEGgly122014PCPC_2*).

The Panel should be prepared to issue a Final Report.

PEGylated
Alkyl Glycerides

Distributed for Comment Only -- Do Not Cite or Quote
SAFETY ASSESSMENT FLOW CHART

Dec 2014



*The CIR Staff notifies of the public of the decision not to re-open the report and prepares a draft statement for review by the Panel. After Panel review, the statement is issued to the Public.
 **If Draft Amended Report (DAR) is available, the Panel may choose to review; if not, CIR staff prepares DAR for Panel Review.

PEGylated Alkyl Glycerides - History

Scientific Literature Review: June 5, 2014

Concentration of use data were received and included in the SLR.

Draft Report: September 8-9, 2014

The following unpublished data submissions were received after the SLR was issued:

1. Specifications of PEG-11 Avocado Glycerides and PEG-10 Olive Glycerides; memo dated June 16, 2014.
 - a. Anonymous. 2013. Specification/certificate of analysis PEG-11 Avocado Glycerides.
 - b. Anonymous. 2013. Specification/certificate of analysis PEG-10 Olive Glycerides.
2. Information on PEG-10 Olive Glycerides, PEG-11 Avocado Glycerides, and PEG-9 Cocoglycerides; memo dated June 24, 2014.
 - a. Zschimmer & Schwarz GmbH & Co KG. 2013. Raw material information PEG-10 Olive Glycerides.
 - b. Zschimmer & Schwarz GmbH & Co KG. 2012. Raw material information PEG-11 Avocado Glycerides.
 - c. Zschimmer & Schwarz GmbH & Co KG. 2013. Raw material information PEG-9 Cocoglycerides.
3. Information on Products Containing PEG Alkyl Glycerides; memo dated July 9, 2014.
 - a. Peritesco. 2004. Ocular and cutaneous tolerance study of self foaming cleansers containing 1% PEG-75 Shea Butter Glycerides.
 - b. Consumer Product Testing Co. 2013. The hen's egg test- utilizing the chorioallantoic membrane (HET-CAM) of a hair styling product containing 11% PEG-7 Glyceryl Cocoate.
 - c. TKL Research. 2013. Repeated insult patch test of a hair styling product containing 11% PEG-7 Glyceryl Cocoate.
 - d. Harrison Research Laboratories, Inc. 2008. Repeated insult patch test of a hair setting lotion containing 1% PEG-75 Shea Butter Glycerides.
 - e. Consumer Product Testing Co. 2005. Safety in use of a female hygiene product containing 0.1% PEG-75 Shea Butter Glycerides.
4. Information on PEG Alkyl Glycerides; memo dated July 24, 2014.
 - a. Anonymous. 2005. Assessment of the eye irritation potential of PEG-7 Glyceryl Cocoate in the hen's egg test on the chorioallantoic membrane (HET-CAM)
 - b. Anonymous. 2005. Assessment of the eye irritation potential of PEG-7 Glyceryl Cocoate by cytotoxicity measurement in the neutral red uptake assay (NRU) on human.
 - c. Hazleton Laboratories America, Inc. 1989. Summary of 28-day subchronic oral toxicity study in rats with a material containing 0.8% PEG-7 Glyceryl Cocoate.
 - d. Research Toxicology Centre S.p.A. 1995. Delayed dermal sensitization study in the guinea pig (PEG-7 Glyceryl Cocoate).
 - e. TKL Research, Inc. 2004. Repeated insult patch test of a styling pomade (BCS578-094) containing 8% PEG-7 Glyceryl Cocoate.
 - f. TKL Research, Inc. 2004. Repeated insult patch test of a styling pomade (BCS578-098) containing 8% PEG-7 Glyceryl Cocoate.
 - g. Pharmaco UK Ltd. 1994. Human repeat insult patch test of PEG-20 Almond Glycerides, 100% neat tested at 5%.
 - h. Leberco-Celsis Testing. 1997. Ocular irritation screen of PEG-10 Sunflower Glycerides using the chorioallantoic membrane vascular assay (CAMVA).
 - i. Raltech Scientific Services. 1981. Skin corrosivity (DOT procedure) of a perfume containing 4% PEG-6 caprylic/capric glycerides.
 - j. Springborn Institute for Bioresearch, Inc. 1981. Rabbit eye irritation (perfume containing 4% PEG-6 caprylic/capric glycerides).
 - k. Anonymous. 1981. Human sensitization test of a perfume containing 4% PEG-6 caprylic/capric glycerides.

Comments on the SLR also were received from the Council.

The Panel incorporated five previously reviewed PEG glyceryl cocoates into the report and issued a tentative report for public comment with the conclusion that the PEGylated alkyl glycerides are safe as used in cosmetics when formulated to be non-irritating.

Draft Final Report: December 8-9, 2014

A summary of a developmental toxicity study on PEG-8 caprylic/capric glycerides was received and added to the report.

PEGylated Alkyl Glycerides – Dec 2014 – Monice Fiume

[illegible]

PEGylated Alkyl Glycerides - Dec 2014 - Monice Fiume

	Reported Use	Method of Mfg	Toxicokinetics	Dermal Penetration	Animal Tox - Acute, Dermal	Animal Tox - Acute, Oral	Animal Tox, Acute, Inhal.	Animal Tox - Rptd Dose, Derm	Animal Tox, Rptd Dose, Oral	Animal Tox - Rptd Dose, Inhal	Repro/Dev Tox	Genotox	Carcinogenicity	PhotocarC	Dermal Irr/Sens	Phototoxicity	Ocular Irritation	Mucous Membrane Irr
PEG-45 Safflower Glycerides																		
PEG-60 Shea Butter Glycerides																		
PEG-75 Shea Butter Glycerides	X														X		X	X
PEG-75 Shorea Butter Glycerides																		
PEG-35 Soy Glycerides	X																	
PEG-75 Soy Glycerides																		
PEG-2 Sunflower Glycerides																		
PEG-7 Sunflower Glycerides																		
PEG-10 Sunflower Glycerides	X																X	
PEG-13 Sunflower Glycerides	X																	
PEG-5 Tsubakiate Glycerides																		
PEG-10 Tsubakiate Glycerides																		
PEG-20 Tsubakiate Glycerides																		
PEG-60 Tsubakiate Glycerides																		

*"X" indicates that data were available in a category for the ingredient

PEGylated Alkyl Glycerides – Search Terms

67762-35-0
70377-91-2
103819-44-9
103819-45-0
103819-46-1
124046-50-0
124046-52-2
180254-52-8
186511-05-7

PubMed – April 10, 2014

(PEG OR (polyethylene AND glycol) OR (polyoxyethylene AND oxide) OR polyoxyethylene OR POE OR PEO) AND ((almond OR (apricot AND kernel) OR avocado OR babassu OR (caprylic AND capric) OR ((cocoa OR shea OR shorea) AND butter) OR corn OR (evening AND primrose) OR (hydrogenated AND fish) OR macadamia OR mango OR mink OR moringa OR mushroom OR olive OR palm OR (passiflora AND seed) OR safflower OR soy OR sunflower OR tsubakiate) AND glyceride*) OR cocoglyceride OR (coco AND glyceride)) – 127 hits/0 useful

PEGylated AND alkyl AND glyceride – 0 hits

*glyceride AND ((ethylene AND glycol) OR epoxide) – 219 hits/1 useful

labrasol – 190 hits/3 useful

SciFinder – April 14, 2014

Substance Identifier search – 11 hits via names and CAS numbers/no references found

Research Topic search – PEG alkyl glycerides; glyceride ethylene glycol; glyceride epoxide; PEG Glycerides; PEG Almond Glycerides; PEG Apricot Kernel Glycerides; PEG Avocado Glycerides; PEG Babassu Glycerides; PEG Caprylic/Capric Glycerides; Labrasol; PEG Cocoa Butter Glycerides; PEG Coco glycerides; PEG Corn Glycerides; PEG Evening Primrose Glycerides; PEG Hydrogenated Corn Glycerides; PEG Macadamia Glycerides; PEG Mango Glycerides; PEG Mink Glycerides; PEG Moringa Glycerides; PEG Mushroom Glycerides; PEG Olive Glycerides; PEG Palm Glycerides; PEG Palm Kernel Glycerides; PEG Passiflora Glycerides; PEG Passiflora Edulis Seed Glycerides; PEG Passiflora Incarnata Seed Glycerides; PEG Safflower Glycerides; PEG Shea Butter Glycerides; PEG Shorea Butter Glycerides; PEG Soy Glycerides; PEG Sunflower Glycerides; PEG Tsubakiate Glycerides – all with document-type limiters – 1461 hits/ 14 useful

ECHA – REACH - April 15, 2014

by CAS No – 0

by name – 0

ChemPortal

by CAS No – 0

by name – 0

COSMOS – May 1, 2014

by CAS No – 0

by name – 0

PEGYLATED ALKYL GLYCERIDES TRANSCRIPTS – Sept 8-9, 2014

Belsito Team

DR. BELSITO: Okay. So PEGylated alkyl glycerides, it's the first time we're seeing the report. 60 ingredients in conditioning agents and surfactants and there were similarities to the glyceryl cocoates in 1999 that were found safe up to 10 percent and there's PEG-3 glyceryl cocoate that's in this report, right?

MS. FIUME: Yes.

DR. BELSITO: And then we got a huge amount of data in wave two and Monice was nice enough to actually create a whole new document for us to look at after I'd already looked at the first one.

DR. LIEBLER: Same thing.

DR. BELSITO: So my comments are in the first one and my spelling corrections to the additions are in wave two. So I will move to the first document here. PEGylated (inaudible) okay. Safe as used, plant -- in a discussion, plant derived not absorbed, so we're not concerned about lack of systemic toxicity. Penetration enhancer and not concerned regarding the irritant effects that occurred at high concentrations beyond those that would be used in cosmetics as part of the discussion. That's all I have. And then a few points that we should have put the oat seed oil here but that's okay.

MS. FIUME: So it's just safe as used not when formulated to be non-irritating?

DR. BELSITO: No. I thought the irritation were at levels that were far beyond what we would see in cosmetic products. Yeah, I mean, and that should go in the discussion since we are going to see that data on irritation. But I don't think we need to put it -- so the discussion is the plant-derived boilerplate, lack of systemic toxicity.

We're not concerned because basically there's not going to be significant absorption. Penetration enhancer boilerplate and the irritant effects were at concentrations higher than are used in cosmetic products.

MS. FIUME: Can I ask at this Dr. Bergfeld had alluded -- I did a test run trying to create a road map for you with what the filings were compared to if you were finding them in the pdf. Did you find it useful, confusing and did you like that version? In the transmittal I have a parentheses in italicized text what each of the parts of the pdf were called so that if you wanted to find the data that referred -- I talked about in a certain study, it told you where it was at.

And I was just curious if you found that useful, confusing, it didn't matter? So that -- because I know Dr. Bergfeld will often ask for a way to help navigate the documents because we really can't do any type of table of contents in the pdf.

DR. BELSITO: I mean, I actually just go to edit, find and do it that way. So, I mean, I just do it myself. But I mean, I -- so I'm neutral on that position.

MS. FIUME: Okay.

DR. BELSITO: If it helps Dr. Bergfeld, then continue to do it. If she felt it was confusing and prefers not to, I'm fine with not doing it. I mean, I just --

MS. FIUME: She did like it. She said that it was helpful for her.

DR. BELSITO: Then keep it.

MS. FIUME: Okay.

DR. GILL: I think the issue is it'll be a uniform (inaudible) writers.

DR. BELSITO: I see. I mean, if it's very helpful to -- I mean, we're also -- I, well, accept for Dan I think most of us are struggling to get used to these documents and we're all finding our ways of making them a little more friendly. I just find the find tab to be very easy for me.

The one thing that I do find difficult a little bit is with the administrative documents where it almost would be nice if when we were doing re-review or boilerplates, rather than lumping them into an administrative document like we used to before a buff document, if we actually had them as individuals. So like this time, like, the for re-review priorities were just a separate document. So I wasn't having to scroll through this administrative document. That's the only thing that I --

DR. SNYDER: Just print the agenda and those re-reviews. It'll be like 10 pages. I mean, just print and include them with it.

DR. BELSITO: No, I mean, I just think separating the items out rather doing, you know, in the administrative I'd like to just see the agenda and the minutes. And then, if there are other issues like a re-review document just have a separate tab for re-review so I can go there and do it rather than having re-reviews mixed in with boilerplates, mixed in with this, mixed in with that where I have to, you know, sort of scroll through stuff that I didn't look at all 346 pages of the priority re-review. But I had to scroll and search and get to the questions Bart had and it just took me a lot of time.

DR. GILL: It may be easier in the electronic form.

DR. BELSITO: Right.

DR. GILL: I know they did it in the past because they were trying to minimize how many books you guys had, so.

DR. BELSITO: Right.

DR. GILL: It may be easier.

DR. BELSITO: But electronically, I think separating them out makes it much easier for me. That would be the only thing that I find a hassle.

DR. LIEBLER: No, I wholeheartedly agree. I mean, it's just the way that their package is to hold over from having the books.

DR. GILL: Right.

DR. LIEBLER: Yeah.

DR. BELSITO: Okay, anything else on these PEG glycerides? No? Okay.

Marks Team

DR. MARKS: Okay. If there are more than that it should come, obviously, back to the panel. Okay. Any other comments about this?

So the next ingredient are the PEG glycerides. So this is the first time we've seen this report. There are 60 ingredients, and, of course, I'm going to be asking Ron, Ron, and Tom are these ingredients okay. Now we're in the combination mode. Do we combine the five PEGylated glyceryl cocoates?

They actually are fractions of the oil so the PEGylated oils, so. My question is do you combine the oils with these too? But what needs do we have? So should we go to the long list at this point in combining? How do you want to proceed, Ron, Ron, and Tom? Ingredients?

DR. SLAGA: So we're really talking about adding a large number.

DR. MARKS: Yes.

DR. SLAGA: Even the PEG 3 cocoate? They're all, definitely, related.

DR. MARKS: Anything stand out that shouldn't be included when you look down the list? I don't --

DR. HILL: No, everything fit from where I stand.

DR. MARKS: Okay. So all 60 ingredients are okay? That included the combination, right, Barb or Monice?

MS. FIUME: It did.

DR. MARKS: And Monice, tell me more about this. If the PEGylated alkyl glycerides are a sub-fraction of the PEGylated oils. And we said the oils are safe when formulated to be non-irritating. I kind of like the reasoning, you know, do we subdivide this? Have we done that before with these 60 ingredients or should they be under the oils?

It's sort of a hypothetical question when I saw that they're a sub-fraction. What's your feeling Ron Hill, and Ron Shank, and Tom?

DR. HILL: Could you ask the question again while they're preparing the thing?

DR. MARKS: If the --

DR. HILL: I'm sorry. I was looking at something else, and I was half-listening, and now I'm (inaudible).

DR. MARKS: This is from in the fourth paragraph where it says the glycerides are sub-fractions of the oils, and we said the oils were safe when formulated to be non-irritating. If we're combing reports should we be combining it with the oils?

DR. HILL: These are PEGylated, right?

DR. MARKS: Yes.

DR. HILL: So no.

DR. BERGFELD: I'm sorry, no you should combine them, no you shouldn't?

DR. HILL: No, do not combine them.

DR. BERGFELD: Okay. Thank you.

DR. MARKS: Because the oils are PEGylated too.

DR. HILL: What oils are we talking about that are -- okay, wait a minute. I'm making sure I'm looking at the right there. Where --

DR. MARKS: It's the fourth paragraph.

DR. HILL: Of what page?

DR. MARKS: The memo.

DR. HILL: The memo?

DR. MARKS: That's Page 2.

DR. HILL: Yeah. Okay. I think the problem is I'm looking at the Wave 2 version, so let me go back. That's the problem.

DR. MARKS: And we don't --

DR. HILL: I just shifted to Wave 2 because they did a complete revision of the report.

DR. MARKS: We certainly don't have to do that. I just was raising if it's a sub-fraction of oils, and we've already said the oils were safe when formulated non-irritating. Should we reopen the oils and just add this group to it? No?

MR. PEREIRA: Sorry. I have a --

DR. MARKS: Do you want to come up? Yeah, thank you for your comments.

MR. PEREIRA: (inaudible) for clarity.

DR. HILL: Could you say who you are again, sir, please?

MR. PEREIRA: Yes, Abel Pereira, research director. I'm with CRODA Incorporated. There's a term, there's PEGylated oils, and then there's PEGylated glycerides. Is that what you're trying to --

DR. MARKS: Yes.

MR. PEREIRA: They, in the industry, unfortunately, both end up being the same. So, in essence, one takes a triglyceride, an oil, and you either do a reaction with ethylene oxide, in which case it breaks apart, and you end up with PEGylated glycerides.

You could do a trans esterification with glycerin beforehand, and then detoxilate it, and you end up with -- you know, they're slightly different in the quality and so forth, but in the end they're all basically the same. I hope that answers the question.

DR. HILL: The major difference is just the sequence in the method of manufacture?

MR. PEREIRA: Correct.

DR. HILL: Yeah, that's what I thought I understood.

DR. MARKS: Bart?

DR. HELDRETH: Yeah, we simply sectioned them off separately here because typically when the INCI committee gives a more specific name it's because they looked that submission package that they've gotten, and seen that there was more specifics there.

But if that's not the case in reality, and these are actually virtually identical to PEGylated oils then, I mean, we don't necessarily have to combine them. But it doesn't sound as if there's anything as far as scientifically from combining them. It may be an administrative headache to push the whole thing back for combination purposes. I mean, administratively we could always go forward with this report as is, and combine them when we do a re-review. You'll see a strategy like you saw today to combine PEGylated oils and these. At that point we'll make them all the same now knowing that they're virtually identical structurally.

DR. HILL: Yes, because that would, in the interim, provide more time to find out, well, given that the sequence is different. Okay, the composition's pretty similar, but how similar is similar? In the end it may, not in practicality matter, but maybe by that point in time they'll be some additional analytical data.

We're starting to see the sorts of things I would like to see and hear like the NMR shows that here's the nature of the mixture, and we haven't, in the past, been seeing enough of that from where I sit. So now we are.

So maybe, you know, going forward people will better characterize some of these mixtures, and we'll have a better idea of what the distribution is, and then we can note, does combining them make great sense. It sounds like it might.

DR. MARKS: Okay. So we're at the 60 ingredients. It's interesting, the discussion on these, when you added that paragraph it triggered maybe should we put them together, obviously, even more.

MS. FIUME: The purpose was to support the safety of this group.

DR. MARKS: Right.

MS. FIUME: Because you've looked at a very similar group and what you found, and that was the purpose of letting you know that we've looked at something very similar in the past, not, hey, do you want to combine these? Because we did just review them in 2012.

DR. MARKS: That's known as the unintended consequences.

MS. FIUME: I've learned my lesson.

DR. SLAGA: So should we table to handle all that, no?

DR. MARKS: No. I think we decided we'll do the 60, and then in the future with a re-review perhaps combine them at that point, is my understanding.

Now, why was PEG-3 glyceryl cocoate not included in the original report?

MS. FIUME: At that time it just may not have been listed as an ingredient. It may not have had use. It may have just been overlooked.

I was going to add, to the purpose of that subgroup being folded into this new list is we noticed as we were developing the SLR that up for re-review this year were the PEG glyceryl cocoates, which were fitting, as we looked at it, to this group. So we thought rather than do a separate re-review and a separate SLR we would fold them in, and add that one ingredient that was either missed or not in use or not listed at the time of the original PEG glyceryl cocoate report.

DR. MARKS: I just wanted to make sure there wasn't some reason it wasn't stated in the report why it wasn't included.

MS. FIUME: Not to my knowledge.

DR. MARKS: Ron Shank, do you remember back then?

DR. SLAGA: (inaudible).

DR. MARKS: Tom or --

DR. SLAGA: I don't think anybody remembers.

DR. MARKS: Yeah, exactly. So we have 60 ingredients. We've agreed on that. The ones listed are all okay. Any needs, Ron, Ron, and Tom to move this forward?

DR. SHANK: No needs. I think you can go safe as used. Pardon me, safe as used when formulated to be non-irritating.

DR. MARKS: Ron Hill?

DR. HILL: The only thing I felt like I lacked was possibility of any residual plant originating substances of toxicological concern. So we don't have any data speaking to that in our method of manufactured data, at least for a whole lot of these. So I thought, well, okay, any of the ones that we don't have that specifically for that compound, insufficient. That was just sort of a not a real way of thinking, but just to put it out there as why don't we have this information. It would be nice to try and --

DR. SLAGA: I have a question. When the PEGylated oils were found safe to be non-irritating, refresh my memory, what was the smallest one in that grouping?

MS. FIUME: PDF page --

DR. SLAGA: I mean, we have data down to PEGylated 6 which --

DR. HILL: Yes, and based on those concerned --

DR. SLAGA: -- down to that, but if we had three, and we have a two do we need anything on these lower or do we have that related to the PEGylated oils?

MS. FIUME: So you can double check me, the PEGylated oils that were reviewed are listed on a PDF Page 69, but there was a PEG-2 castor oil, a PEG-3 castor oil, a PEG-

Hydrogenated castor oil. So there were low PEGs.

DR. SLAGA: Yeah, thanks. I agree with Ron Shank.

DR. HILL: Let me go back one step from where I was to the thing I was just talking about because I remembered where my comfort level came from, sort of. Because what I had written here was if we know, in capital letters, that they started with a plant derived oil that meets cosmetic grade or food grade standards then the concern goes away. Can we infer that for all of the vegetable names ingredients in this report? That's what I kind of need an answer to.

DR. SHANK: What was your question specifically?

DR. HILL: What I said was in interest and concern of assessing the safety is do we know that -- well, I'd written down heavy metals, but I kind of wrote that off. But also, do we know about the possibility or residual plant originating substances of toxicological concern because they are plant derived oils.

Then what I had written after that I didn't look at when I was talking because apparently I can't walk and chew gum at the same time. Can we infer that they're all using cosmetic grade or food grade standard oils as a starting material? Can we get information that, more or less, confirms that, and any of the ones that we don't know for sure? Maybe that's insufficient until we get it.

MS. FIUME: Dr. Hill, I don't know if we can ever, if we ever do speak to that, whenever we review something that's plant derived. But previous CIR reports are available on all component oils except hydrogenated corn oil, although we do have a corn oil report, hydrogenated fish oil, mushroom oil, and then the passiflora incarnate oil.

DR. HILL: Actually, those were the four that were on my mind when I wrote this comment.

MS. FIUME: But, I believe, that should be something we can address in the discussion, correct?

DR. HILL: I don't know. That's why I'm asking it.

DR. HELDRETH: I can't speak to whether or not it's true for all of these definitions, but many of these, when they list that it's a combination of the PEG and the oil, although it's removed from our table because it wouldn't make sense, but in the actual dictionary it says quo vida after the name of the oil saying that it is actually an ingredient in the dictionary. So they're proposing, in that definition, that it actually starts with the cosmetic ingredient.

DR. HILL: Would that other cosmetic ingredient automatically be, if they use that, as their feed stock would they -- I don't think that that necessarily implies that they would be starting with food grade or cosmetic grade material where we wouldn't have the -- and so the concern.

I've noticed in the transcripts I have a lot of incomplete sentences. So, yes, maybe Mr. Steinberg could speak to that.

MR. STEINBERG: With the exception of the use of any of these oils to make soap they're all chemically treated before they were even put into industrial commerce. The treatment is basically to remove any free fatty acids.

So all of the vegetable oils, whether it be corn oil, soybean oil, or castor oil are first treated with a wash, and then they are, I think, clay treated, and therefore, you just have the triglyceride present or the phytosterols which are soluble in the triglyceride.

DR. HILL: Yeah, so if the phytosterols are there that's the point.

MR. STEINBERG: Yes. They would still be in there --

DR. HILL: But they have potential for things like linalool and such?

MR. STEINBERG: Yeah.

DR. HILL: Because the fatty acid wash would probably be an alkaline wash --

MR. STEINBERG: Yes.

DR. HILL: -- and that won't remove that sort of thing.

MR. STEINBERG: Right.

DR. SHANK: Many of these are used as incipients in pharmaceuticals and food additives. So I think, perhaps, the answer lies in that.

DR. HILL: Yes. When they are used as -- then they will be using pharmaceutical grade material for those incipients because that's legally required, if I understand correctly. But I don't know if that's true if we're talking about a cosmetic product. So it seemed reasonable to ask it, particularly for those small number of oils that were not in previous reviews.

DR. MARKS: So tomorrow I'm going to move that a tentative report be issued with a conclusion that all 60 ingredients are safe when formulated to be non-irritating. Ron Hill, do you want tomorrow to make a comment about that? It sounds like Ron Shank and Tom Slaga you're fine with that.

DR. HILL: I mean, this is not the final report, right?

DR. MARKS: Yeah.

DR. HILL: So that'll be captured in the transcript, and for me right now that's good enough.

DR. MARKS: Right. Yes, okay.

DR. BERGFELD: Could I ask a question? For the record, I wonder, Carol, if you would define cosmetic grade?

DR. EISENMANN: No, there really isn't such a thing as cosmetic grade.

DR. HILL: I didn't think that there was, I just wrote it down to be sure. I know there is such a thing as food grade.

Full Panel

DR. BERGFELD: Moving on to the PEG glycerides, Dr. Marks?

DR. MARKS: This is the first time we've evaluated these ingredients. And we reviewed a draft report of the 60 ingredients and our team felt that we could move forward with a tentative report, having a conclusion of safe when formulated to be non irritating. So that's a motion.

DR. BELISTO: Well, we felt that the irritation was really at very high concentrations that weren't used in cosmetics, so we just went as a safe as used, but I'm not going to argue with you if you want to say safe when formulated to be non irritating. I don't think it puts any further restrictions on how they're currently being used.

DR. BERGFELD: Any other discussion?

DR. BELISTO: Yeah.

DR. BERGFELD: Don't

DR. BELISTO: So these are plant derived, so we need the boilerplate there. There's a relative lack of systemic toxicity data in the document, but they're large molecules, they're not absorbed. They're penetration enhancers, so the usual penetration boilerplate, and that was it. And I want to thank Monice for rewriting to include all of the data we had in Wave 2 and give us a finished report.

DR. BERGFELD: She also gave us the roadmap

DR. BELISTO: Right.

DR. BERGFELD: of this particular ingredient, so thank you for that, as well.

All right, no other comments? We've had a motion, it's been seconded, to proceed with safe, and I gather it's going to say non irritating, is that the agreement?

DR. BELISTO: Yes, that's fine.

DR. BERGFELD: All those in favor, please indicate by raising your hands?

(All hands raised.)

DR. BERGFELD: Thank you, unanimous.

Safety Assessment of PEGylated Alkyl Glycerides as Used in Cosmetics

Status: Draft Final Report for Panel Review
Release Date: November 14, 2014
Panel Meeting Date: December 8-9, 2014

The 2014 Cosmetic Ingredient Review Expert Panel members are: Chair, Wilma F. Bergfeld, M.D., F.A.C.P.; Donald V. Belsito, M.D.; Ronald A. Hill, Ph.D.; Curtis D. Klaassen, Ph.D.; Daniel C. Liebler, Ph.D.; James G. Marks, Jr., M.D.; Ronald C. Shank, Ph.D.; Thomas J. Slaga, Ph.D.; and Paul W. Snyder, D.V.M., Ph.D. The CIR Director is Lillian J. Gill, D.P.A. This safety assessment was prepared by Monice M. Fiume, Assistant Director/Senior Scientific Analyst and Bart Heldreth, Ph.D., Chemist.

ABSTRACT

The CIR Expert Panel assessed the safety of 60 PEGylated alkyl glycerides. PEGylated alkyl glycerides are mono-, di-, and/or tri-glycerides that have been modified with a specific number of ethylene glycol repeat units (in the starting material form as epoxide). Most of the PEGylated alkyl glycerides are reported to function as skin conditioning agents or surfactants. The Panel reviewed the available animal and clinical data, as well as data from the 1999 report of PEG glyceryl cocoates and the 2012 report of PEGylated oils, to determine the safety of these ingredients. The Panel concluded these ingredients are safe in the current practices of use and concentration when formulated to be non-irritating; this conclusion supersedes the 1999 conclusion issued on five PEG glyceryl cocoate ingredients.

INTRODUCTION

This report assesses the safety of the following 60 PEGylated alkyl glycerides as used in cosmetic formulations:

PEG-6 Almond Glycerides	PEG-6 Hydrogenated Palm/Palm Kernel Glyceride
PEG-20 Almond Glycerides	PEG-16 Macadamia Glycerides
PEG-35 Almond Glycerides	PEG-70 Mango Glycerides
PEG-60 Almond Glycerides	PEG-13 Mink Glycerides
PEG-192 Apricot Kernel Glycerides	PEG-25 Moringa Glycerides
PEG-11 Avocado Glycerides	PEG-42 Mushroom Glycerides
PEG-14 Avocado Glycerides	PEG-2 Olive Glycerides
PEG-11 Babassu Glycerides	PEG-6 Olive Glycerides
PEG-42 Babassu Glycerides	PEG-7 Olive Glycerides
PEG-4 Caprylic/Capric Glycerides	PEG-10 Olive Glycerides
PEG-6 Caprylic/Capric Glycerides	PEG-40 Olive Glycerides
PEG-7 Caprylic/Capric Glycerides	PEG-18 Palm Glycerides
PEG-8 Caprylic/Capric Glycerides	PEG-12 Palm Kernel Glycerides
PEG-11 Cocoa Butter Glycerides	PEG-45 Palm Kernel Glycerides
PEG-75 Cocoa Butter Glycerides	PEG-60 Passiflora Edulis Seed Glycerides
PEG-7 Cocoglycerides	PEG-60 Passiflora Incarnata Seed Glycerides
PEG-9 Cocoglycerides	PEG-45 Safflower Glycerides
PEG-20 Corn Glycerides	PEG-60 Shea Butter Glycerides
PEG-60 Corn Glycerides	PEG-75 Shea Butter Glycerides
PEG-20 Evening Primrose Glycerides	PEG-75 Shorea Butter Glycerides
PEG-60 Evening Primrose Glycerides	PEG-35 Soy Glycerides
PEG-3 Glyceryl Cocoate	PEG-75 Soy Glycerides
PEG-7 Glyceryl Cocoate*	PEG-2 Sunflower Glycerides
PEG-30 Glyceryl Cocoate*	PEG-7 Sunflower Glycerides
PEG-40 Glyceryl Cocoate*	PEG-10 Sunflower Glycerides
PEG-78 Glyceryl Cocoate*	PEG-13 Sunflower Glycerides
PEG-80 Glyceryl Cocoate*	PEG-5 Tsubakiate Glycerides
PEG-5 Hydrogenated Corn Glycerides	PEG-10 Tsubakiate Glycerides
PEG-8 Hydrogenated Fish Glycerides	PEG-20 Tsubakiate Glycerides
PEG-20 Hydrogenated Palm Glycerides	PEG-60 Tsubakiate Glycerides

*originally reviewed in 1999

PEGylated alkyl glycerides are mono-, di-, and/or tri-glycerides that have been modified with a specific number of ethylene glycol repeat units (in the starting material form as epoxide). Most of the PEGylated alkyl glycerides are reported to function as skin conditioning agents or surfactants¹ (Table 1).

Five PEG glyceryl cocoate ingredients, indicated above, previously have been reviewed, and in 1999 the Cosmetic Ingredient Review (CIR) Expert Panel (Panel) concluded those ingredients were safe as used in rinse-off products and safe at up to 10% in leave-on products. According to the Discussion of that report, the basis of the concentration limit for leave-on products was the ocular and sensitization data.² At the September 2014 meeting, the Panel determined that it is appropriate to include those ingredients in this report, and where appropriate, information from the 1999 report is included in this safety assessment, as indicated by *italicized text*.

In 2012, the Panel reviewed a very similar family of ingredients, the PEGylated oils, and concluded the PEGylated oils are safe as used in cosmetics when formulated to be non-irritating.³ PEGylated oils are complex mixtures of the etherification and transesterification products of fatty acid glycerides and fatty acids derived from identified plant sources with the equivalents of ethylene oxide to reach the desired polyethylene glycol (PEG) length. The only difference between the PEGylated

oils and the PEGylated alkyl glycerides is that prior to the PEGylation step in the production of PEGylated alkyl glycerides, a further refinement of the oil has been performed to retain only the glycerides, and in some cases refined (or partially de-esterified) to only the mono-, di-, or triglycerides. In other words, PEGylated alkyl glycerides are a sub-fraction of PEGylated oils. PEGylated oils mostly contain PEGylated triglycerides and some PEGylated fatty acids (i.e., not attached to a molecule of glycerin), while PEGylated alkyl glycerides contain only specific PEGylated glycerides and no (or virtually no) PEGylated fatty acids.

As an example, the PEGylated oil, almond oil PEG-6 esters, is comprised mostly of *triglycerides*, wherein the six stoichiometric equivalents of ethylene oxide have been inserted between the glycerin core and the attached almond fatty acids. Free almond fatty acids that have been PEGylated to some degree *may also be present*. In contrast, the PEGylated alkyl glyceride, PEG-6 almond glycerides, is comprised mostly of *mono-* and *diglycerides*, wherein six stoichiometric equivalents of ethylene oxide have 1) been inserted between the glycerin core and the attached almond fatty acids, 2) *etherified the free alcohol(s) of the glyceride, or 3) both*. No appreciable free almond fatty acids should be present.

The materials that are used in the manufacture of the PEGylated alkyl glycerides, i.e., ethylene oxide (and any incidentally generated free PEG chains) and the applicable glycerides (and the oils from which they are derived), may be present as impurities in the final ingredient. The CIR has reviewed the safety of the majority of the component ingredients, and although the information from those safety assessments may not be relevant for read-across, the reports have some relevance because starting material may be present as residual impurities. Therefore, summary information and the conclusion of each relevant report are provided⁴⁻¹¹ (Table 2). Previous CIR reports were available on all component oils except hydrogenated corn oil (although a report on corn oil was available), hydrogenated fish oil, mushroom oil, and *Passiflora incarnata* oil. Additionally, the conclusion and a data summary from the PEGylated oils³ report are provided (Table 2). Finally, CIR currently is reviewing the safety of glycerin.

CHEMISTRY

Definition and Structure

PEGylated alkyl glycerides are mono-, di-, and/or tri-glycerides that have been modified with a specific number of ethylene glycol repeat units (in the starting material form as epoxide).




Figure 1. generic PEGylated alkyl glyceride structure, wherein R is hydrogen or the fatty acids declared in the name or found in the source material, and the sum of all cases of n is "X" in PEG-"X"




Figure 2. PEG-6 caprylic/capric glycerides, wherein the sum of all instances of “n” is 6 (i.e., “PEG-6”).

Just as oils and other PEGylated materials are mixtures, the PEGylated alkyl glycerides are mixtures. PEGylation of glycerides occurs not only as an etherification of the free alcohol groups of the glycerides with ethylene oxide groups, but also as a transesterification which results in net insertion of PEG groups between the glyceryl and fatty acid components of the glyceride.^{12,13} For example, the glycerin core of PEG-6 caprylic/capric glycerides is ethoxylated (Figure 2), wherein the sum of all n is equal to the number of ethylene oxide repeat units and need not be the same at all places of the molecule.

It should be stated that n is not equal to X. In other words, wherein the value of X in PEG-X is equal to 6 (e.g., PEG-6 caprylic/capric glycerides), n is not equal to 6. Instead, X represents the number of stoichiometric equivalents of ethylene oxide that were added to one stoichiometric equivalent of caprylic/capric glycerides. Therefore, the sum of all of the different n values in the mixture may be no more than X. Indeed, when one mole of ethylene oxide is reacted with one mole of fatty alcohol, adducts having **no** added ethylene oxide are the predominate material in the mixture.¹² Furthermore, wherein ethylene oxide reacts with mono- or diglyceride, it is approximately twice as likely that it will react at an ester site versus an alcohol site. Moreover, a percentage (13% in one specific case) of the ethylene oxide simply reacts with other molecules of ethylene oxide, resulting in some polyethylene glycols, unattached to glycerides.

Most of the ingredients in this report share in common a specific nomenclature, generically PEG-X “source”/alkyl-chain-common-name glycerides (e.g., PEG-6 caprylic/capric glycerides). The PEG glyceryl cocoates have a slightly different order of nomenclature, but they are not structurally distinct from the other ingredients in this report.

Chemical and Physical Properties

Much like the oils these ingredients are derived from, the PEGylated alkyl glycerides range from oily liquids to waxy solids, depending on alkyl chain length and degree of PEGylation (Table 3). Owing to the combination of polar and non-polar functional groups, most of these ingredients are soluble in some organic solvents and are at least dispersible in water, if not soluble. The fatty acid compositions of the oils that comprise the alkylated glycerides are provided in Table 4.

Methods of Manufacture

PEG-11 avocado glycerides,¹⁴ PEG-9 cocoglycerides,¹⁵ and PEG-10 olive glycerides¹⁶ are produced by the reaction of their respective oil with ethylene oxide. PEG-6 caprylic/capric glycerides is derived completely from a vegetable/synthetic source, and palm kernel oil and/or rapeseed oil are the starting materials.¹⁷

One source states that PEG-6 caprylic/capric glycerides is manufactured in a two-step procedure starting with the preparation of a mono/diglyceride of a C8/C10 fatty acid blend; the C8:C10 ratio is approximately 60:40.¹⁸ The raw materials are charged with an excess of glycerin, the blend is purged with nitrogen and heated under pressure, and the mixture is then cooled, treated with Fuller’s earth, and filtered. Following filtration, step two is initiated, and includes the addition of an alkaline catalyst in a solvent, heating, and the removal of the solvent under pressure. The pressure in the reactor is then normalized with nitrogen, and ethylene oxide is added until the desired saponification value is reached; the mixture is then stirred until the ethylene oxide value is ≤ 1 ppm. After cooling, the product is neutralized with lactic or acetic acid. Finally,

the raw ethoxylate is treated with steam to remove the volatile impurities, and if necessary, the product may be bleached with Fuller's earth.

PEG-8 caprylic/capric glycerides can be obtained by partial alcoholysis of medium-chain triglycerides using PEG, by esterification of glycerol and PEG with caprylic acid and capric acid, or as a mixture of glycerol esters and condensates of ethylene oxide with caprylic acid and capric acid.^{19,20}

Methods of manufacture of the other PEGylated alkyl glycerides were not found in the published literature, nor were unpublished data provided.

Composition and Impurities

Residual PEG may be present in the PEGylated alkyl glycerides. PEGs are the condensation products of ethylene oxide and water, with the chain length controlled by number of moles of ethylene oxide that are polymerized. PEGs may contain trace amounts of 1,4-dioxane, a by-product of ethoxylation; 1,4-dioxane is a known animal carcinogen.²¹ The Food and Drug Administration (FDA) has been periodically monitoring the levels of 1,4-dioxane in cosmetic products, and the cosmetic industry reported that it is aware that 1,4-dioxane may be an impurity in PEG-containing ingredients and, thus, uses additional purification steps to limit it in the ingredient before blending into cosmetic formulations.^{22,23}

Raw material information sheets for PEG-11 avocado glycerides,¹⁴ PEG-9 cocoglycerides,¹⁵ and PEG-10 olive glycerides¹⁶ state that these compounds contain <5 ppm 1,4-dioxane, <1 ppm ethylene oxide, and <1 ppm residual monomers. Solvents are not added to these materials. PEG-6 caprylic/capric glycerides contain ≤5 ppm dioxane and ≤1 ppm ethylene oxide, and it meets the valid regulatory requirements for limits on agricultural pesticides.^{17,24} Residual solvents, monomers, free amines, and nitrosamines are not by-products of the production process.

Nuclear magnetic resonance spectroscopy indicated PEG-8 caprylic/capric glycerides consists of 30% mono-, di-, and triglycerides of C₈ and C₁₀ fatty acids, 50% of mono- and diesters of PEG-8 (i.e., esters of capric or caprylic acid with PEG-8; not glycerides), and 20% free PEG-8.²⁵ It is reported to contain ≤10 ppm 1,4-dioxane, <1 ppm ethylene oxide, and <10 ppm heavy metals (Pb).^{26,27}

USE

Cosmetic

Most of the PEGylated alkyl glycerides are reported to function as a skin conditioning agent – emollient and as a surfactant – emulsifying agent¹ (Table 1). Emollients function by their ability to remain on the skin surface or in the stratum corneum to act as lubricants, to reduce flaking, and to improve the skin's appearance. For the surfactant – emulsifying agent function, the efficacy of an emulsifying agent depends on its ability to reduce surface tension, to form complex films on the surface of emulsified droplets, and to create a repulsive barrier on emulsified droplets to prevent their coalescence. A few of the PEGylated alkyl glycerides also reported to act as surfactant - solubilizing agents or surfactant – cleansing agents.

The FDA collects information from manufacturers on the use of individual ingredients in cosmetics as a function of cosmetic product category in its Voluntary Cosmetic Registration Program (VCRP). In 2014, VCRP data obtained from the FDA,²⁸ and data received in response to surveys of the maximum reported use concentration by category that were conducted by the Personal Care Products Council (Council),²⁹⁻³¹ indicate that 21 of the 60 ingredients included in this safety assessment are used in cosmetic formulations (Table 5). (The ingredients not currently reported to be used are listed in Table 6.)

According to 2014 VCRP data, PEG-7 glyceryl cocoate has the most reported uses, i.e., 858; the majority of these uses (713) are in rinse-off formulations.²⁸ PEG-6 caprylic/capric glycerides is reported to be used in 548 formulations, the majority of which are rinse-off formulations, and PEG-60 almond glycerides is reported to be used in 163 cosmetic formulations. All other in-use PEGylated alkyl glycerides are reported to be used in less than 45 formulations. The PEG glyceryl cocoates have increased in use. In 1996, PEG-7 glyceryl cocoate was reported to be used in 173 formulations, and it is now reported to be used in 858 formulations; however, both then and now, PEG-7 glyceryl cocoate is used mostly in rinse-off formulations.^{2,28} The frequency of use also has increased for PEG-30 glyceryl cocoate (from 10 to 44 uses), PEG-40 glyceryl cocoate (from 5 to 23 uses), and PEG-80 glyceryl cocoate (from 2 to 11 uses).

The results of the concentration of use survey conducted by the Council indicate the highest leave-on concentrations reported are 11.3% and 6% for PEG-7 glyceryl cocoate in tonics, dressings, and other hair grooming aids and "other" fragrance preparations.²⁹ PEG-7 glyceryl cocoate also has the highest rinse-off concentration of use reported, i.e., 10% in skin cleansing products.

At the time of the original safety assessment, concentration of use data were not available from the FDA. However, based on ocular and sensitization data, a concentration limit of 10% was placed on leave-on products containing PEG glyceryl cocoates. The current reported concentration of use in leave-on products is slightly higher, i.e., 11.3% in tonics, dressings, and other hair grooming aids.

One ingredient is used in products that could be incidentally ingested (i.e., PEG-75 cocoa butter is reported to be used in two lipstick formulations) and a few of the ingredients are used near the eye (e.g., PEG-20 hydrogenated palm glycerides is used

at 1.2% in an eyebrow pencil) or mucous membranes (PEG-7 glyceryl cocoate is used at up to 6.9% in other personal cleanliness products). Five PEGylated alkyl glycerides were reported to the VCRP as used in baby products, but concentration of use data were not submitted by industry for these uses. Additionally, according to the VCRP, several ingredients are used in products that can be incidentally inhaled, and results of the Council survey reported concentrations of up to 6% PEG-7 glyceryl cocoate in "other" fragrance preparations, 2% PEG-7 glyceryl cocoate in a pump spray deodorant, and up to 1.9% PEG-45 palm kernel glyceride in a pump spray suntan product. In practice, 95 to 99% of the droplets/particles released from cosmetic sprays have aerodynamic equivalent diameters $>10\text{ }\mu\text{m}$.^{32,33} Therefore, most droplets/particles incidentally inhaled from cosmetic sprays would be deposited in the nasopharyngeal and bronchial regions and would not be respirable (i.e., they would not enter the lungs) to any appreciable amount.^{34,35}

The PEGylated alkyl glycerides named in this safety assessment are listed in the European Union inventory of cosmetic ingredients.³⁶

Non-Cosmetic

PEG-8 caprylic/capric glycerides has use as a pharmaceutical excipient.³⁷ It has a maximum potency of 70 mg in oral soft gelatin capsules and 6.12% in oral solutions.³⁸

A mixture of mono-, di- and tri-glycerides and PEG mono- and di-esters of fatty acids (polyglycides) of hydrogenated vegetable oils may be safely used as an excipient in dietary supplement tablets, capsules, and liquid formulations that are intended for ingestion in daily quantities measured in drops or similar small units of measure when the total ester content is $>90\%$, the acid value is <2 , the hydroxyl value is <56 , the lead content is $<0.1\text{ mg/kg}$, and $<10\text{ mg/kg}$ 1,4-dioxane and $<1\text{ mg/kg}$ ethylene oxide is present (21CFR172.736).

Fats, oils, fatty acids, and fatty alcohols derived from coconut, palm, and soybean oils and reacted with 400-3000 molecular weight PEG are approved for use as indirect food additives in polymers for use as components of articles intended for repeated use in textiles and textile fibers (21CFR177.2800). Also, the fatty acids and alcohols derived from coconut, corn, palm, and soybean oils reacted with 200-4600 molecular weight PEG, with or without dehydration, to form an ester are approved for use as an indirect food additive in defoaming agents used in the manufacture of paper and paperboard components (21CFR176.210).

TOXICOKINETICS

Penetration Enhancement

Surfactants can enhance penetration of some compounds through the skin. The ability of PEG-8 caprylic/capric glycerides to be a dermal penetration enhancer has been investigated in a number of studies; a few examples of those studies are provided. In one study in which transdermal delivery of lidocaine (lipophilic) and prilocaine hydrochloride (hydrophilic) was measured *in vitro* in Franz-type diffusion cells, several microemulsion systems that contained 11.5-53% PEG-8 caprylic/capric glycerides, 3-70% isostearyl isostearate, 8-35% polyglyceryl-6 isostearate, and 7-65% water increased delivery of lidocaine up to four times compared to an oil/water emulsion vehicle, and increased the delivery of prilocaine hydrochloride almost ten times compared to a hydrogel.²⁵ The increase in delivery was attributed to increased solubility of the drug, and appeared to be dependent on drug mobility in the vehicle.

Another example is the effect of PEG-8 caprylic/capric glycerides on the transdermal flux of carvedilol, a non-selective β -adrenergic antagonist, through porcine skin. The flux of carvedilol from a PEG-8 caprylic/capric glycerides vehicle (i.e., $14.28\text{ }\mu\text{g/cm}^2/\text{h}$) was 8.6 times greater than that obtained with the control vehicle (i.e., $1.66\text{ }\mu\text{g/cm}^2/\text{h}$) composed of phosphate buffered saline containing 40% v/v PEG 400.³⁹ However, when placed in the donor compartment, flux was only 4.7 times greater with 5% PEG-8 caprylic/capric glycerides compared to controls. In another study, when used as a vehicle, PEG-8 caprylic/capric glycerides was only a weak penetration enhancer of cyclosporin A, a non-polar cyclic oligopeptide, through excised rat skin.⁴⁰

A formulation containing 40% PEG-8 caprylic/capric glycerides, 39.75% ethoxydiglycol, 10% ethanol, 5% ethyl oleate, and 5% sorbitan oleate improved the skin delivery of kahalalide F, a lipophilic cyclic molecule with a large molecular weight; however, the amount of kahalalide F that penetrated and permeated through the skin was much greater with dimethyl sulfoxide.⁴¹ Additionally, a formulation containing 39.75% PEG-8 caprylic/capric glycerides, 35% propylene glycol, 10% ethanol, 10% *n*-methyl-2-pyrrolidone, and 5% isopropyl myristate did not improve skin delivery of kahalalide F.

TOXICOLOGICAL STUDIES

Single Dose (Acute) Toxicity

Oral

PEG-6 caprylic/capric glycerides has an oral LD_{50} of $>5\text{ g/kg}$ in rats; five males and five females were given a single dose of 5.5 g/kg , and none of the animals died.⁴² PEG-8 caprylic/capric glycerides has an oral LD_{50} of 22 g/kg in rats.⁴³ *The oral LD_{50} of PEG-7 glyceryl cocoate in rats was $>19.9\text{ mg/g}$, the highest dose administered.*²

Repeated Dose Toxicity

Oral

PEG-8 Caprylic/Capric Glycerides

The oral toxicity of a blend of three solvents was evaluated in Wistar rats; the blend consisted of 40% PEG-8 caprylic/capric glycerides, 40% apricot kernel oil PEG-6 esters, and 20% ethoxydiglycol.⁴⁴ (In the published paper, the blend was described using trade names; trade name and other information suggests that the second portion of the blend is apricot kernel oil PEG-6 esters.) Groups of 10 male and 10 female rats were given by gavage a volume of 5, 10, or 20 ml/kg/day test article for 4 wks. A negative control group of 10 rats/sex received 20 ml/kg/day 1% (w/v) hydroxyethylcellulose in purified water. The animals were examined daily for signs of toxicity, and hematology, clinical chemistry, and urinalysis parameters were measured at study termination; at necropsy, organs were examined grossly and microscopically, and hepatic cytochrome P450 content and related activities were evaluated.

The test article was relatively well-tolerated at 5 ml/kg/day, and this volume was considered the no-observable adverse effect level (NOAEL). Adverse effects were observed with the greater doses, and the main effects were observed in the liver, kidneys, and adrenal glands. One female of the 20 ml/kg/day group died on day 28; the death was considered test-article related because the animal had worsening clinical condition prior to death and necropsy results included enlarged and pale kidneys, a dilated stomach with multiple dark red areas on the glandular mucosa, a dilated caecum, and several whitish areas on the liver. In the surviving 20 ml/kg/day animals, there was an increase in clinical signs of toxicity, including ptialism, partial blepharoptosis, and piloerection, and a statistically significant decrease in feed consumption and water intake in males and females and in body weights in males, when compared to control animals. At study termination, most of the high-dose animals had dilated caecums containing soft brownish content, and 12 of the animals had pale adrenal glands; five of the ten males and three of the nine surviving females had enlarged kidneys. Absolute and relative organ-to-body weights were also statistically significantly affected; increases were reported in absolute and relative kidney weights in males and females, relative liver weights in males, and absolute and relative liver weights in females, and decreases were reported in absolute and relative spleen weights in males and females, absolute thymus weights in males, and absolute and relative thymus weights in females. Microscopic findings at this dose included mild centrilobular hepatocellular hypertrophy, mostly mild bilateral diffuse tubular dilation and bilateral multifocal tubular epithelium degeneration and necrosis in the kidney, and vacuolation of adrenal gland cortex. Changes in some hematological parameters were also reported. A volume-related statistically significant increase in induction of hepatic CYP was reported. Some similar effects were reported in animals given 10 ml/kg/day, but not at the same rate of incidence.

No adverse effects were observed in dogs that were dosed orally for 13 wks with 1.0 g/kg/day PEG-8 caprylic/capric glycerides.⁴³ (Details were not provided.)

PEG-7 Glyceryl Cocoate

Groups of 10 male rats were fed a diet containing 0, 2.5, 5.0, or 7.5% of a formulation containing 0.8% PEG-7 glyceryl cocoate for 28 days.⁴⁵ All animals survived until study termination. For all groups, select tissues were weighed at necropsy. Select tissues of animals in the control and high-dose group were examined microscopically. Spleen weights were significantly decreased in high-dose animals; although there were no associated microscopic changes, the researchers stated the change could be associated with dosing. Soft feces, a distended cecum, and enlarged mesenteric lymph nodes were observed at necropsy in the mid- and high-dose animals. The no-observable effect level (NOEL) was 2.5% of the formulation containing 0.8% PEG-7 glyceryl cocoate.

REPRODUCTIVE AND DEVELOPMENTAL TOXICITY

A segment II developmental toxicity test was performed in rats with PEG-8 caprylic/capric glycerides.⁴⁶ The animals were dosed with 0, 1000, 2000, or 3000 mg/kg/day by gavage on days 6 through 17 of gestation. The maternal NOAEL was 2000 mg/kg/day; effects on body weight were reported at 3000 mg/kg/day. The embryo/fetal NOAEL was 3000 mg/kg/day. No signs of embryotoxicity, fetotoxicity or teratogenicity were noted at any dose level. (No other details were provided.)

GENOTOXICITY

PEG-6 caprylic/capric glycerides, PEG-7 glyceryl cocoate, and PEG-10 olive glycerides were not mutagenic, with or without metabolic activation. *Salmonella typhimurium* strains TA98, TA100, TA1535, TA1537, and TA102 were exposed to concentrations of 50-5000 µg/plate PEG-6 caprylic/capric glycerides in one study,⁴² and *S. typhimurium* strains TA98, TA100, TA1535, TA1537, and TA1538 were exposed to 0.04% PEG-6 caprylic/capric glycerides in 0.9% saline in another.⁴⁷ *S. typhimurium* strains TA98, TA100, TA1535, TA1537, and TA1538 also were exposed to 0.625% PEG-7 glyceryl cocoate in 0.9% saline.⁴⁸ Details of testing with PEG-10 olive glycerides, including strains used and concentrations tested, were not provided.¹⁶

Genotoxicity studies of other PEGylated alkyl glycerides were not found in the published literature, nor were additional unpublished data provided.

CARCINOGENICITY

Carcinogenicity data of PEGylated alkyl glycerides were not found in the published literature, nor were unpublished data provided.

IRRITATION AND SENSITIZATION

Dermal

The irritation and sensitization potentials of several PEGylated alkyl glycerides were evaluated in alternative,^{25,49-55} non-human,^{2,16,42,56-62} and human studies,^{2,16,42,63-72} and these ingredients generally were not irritants or sensitizers in the alternative and human studies (Table 7). Mixed results were observed in the non-human studies; specifically, undiluted PEG-20 almond glycerides,⁶¹ PEG-60 almond glycerides,⁶⁰ PEG-12 palm kernel glycerides,⁵⁹ and PEG-45 palm kernel glycerides⁵⁸ were irritating to rabbit skin in primary skin irritation studies, and *PEG-7 glyceryl cocoate (concentration not specified) was mildly irritating to rabbit skin.*² The only reaction reported in clinical testing was PEG-10 olive glycerides (2% active matter) was non-irritating to mildly irritating.¹⁶ In other clinical studies, up to 20% PEG-60 almond glyceride,^{64,69} undiluted PEG-6 caprylic/capric glycerides,^{42,63,66} undiluted PEG-7 glyceryl cocoate,^{2,65,70-72} and a formulation containing 1% PEG-75 shea butter glycerides^{67,68} were not irritants or sensitizers.

Phototoxicity

PEG-7 glyceryl cocoate was not phototoxic in mice at a concentration of 50%.² Formulations containing 0.3% PEG-7 glyceryl cocoate were not photosensitizers in clinical testing.

Ocular

In a hen's egg test utilizing the chorioallantoic membrane (HET-CAM), a formulation containing 11% PEG-7 glyceryl cocoate had practically no irritation potential,⁷³ and concentrations of 10, 50, and 100% PEG-7 glyceryl cocoate were slightly irritating;⁷⁴ in a neutral red uptake (NRU) assay, the ocular irritation potential of PEG-7 glyceryl cocoate was not classified⁷⁵ (Table 8). PEG-10 sunflower glyceride was a non-irritant in a chorioallantoic membrane vascular assay (CAMVA).⁷⁶ In rabbits, 10% aq. PEG-7 glyceryl cocoate² and up to 3% PEG-8 caprylic/capric glycerides were not ocular irritants,⁷⁷ and 5% PEG-8 caprylic/capric glycerides was a slight irritant to rabbit eyes.⁷⁷ Undiluted PEG-6 caprylic/capric glycerides was a strong ocular irritant; however, aq. solutions of up to 5%⁴² and formulations containing 4% PEG-6 caprylic/capric glycerides⁷⁸ were not ocular irritants in rabbits. In clinical testing, a formulation containing 1% PEG-75 shea butter did not induce ocular irritation in a tolerance test.⁶⁸

Mucous Membrane

PEG-10 olive glycerides, 3% active matter, did not irritate the mucous membrane.¹⁶ (No details were provided.)

A female hygiene product containing 0.1% PEG-75 shea butter glycerides did not induce any evidence of vaginovulvar irritation in a use study.⁷⁹ Twenty-one female subjects bathed in a bubble bath using a "quarter-size amount" of the test material for 10-20 min daily for one week.

SUMMARY

This report addresses the safety of 60 PEGylated alkyl glycerides as used in cosmetics. PEGylated alkyl glycerides are mono-, di-, and/or tri-glycerides that have been modified with a specific number of ethylene glycol repeat units (in the starting material form as epoxide). Most of the PEGylated alkyl glycerides are reported to function as skin conditioning agents or surfactants.

VCRP data obtained from the FDA, and data received in response to surveys of the maximum reported use concentration by category that were conducted by the Council, indicate that 21 of the 60 ingredients included in this safety assessment are used in cosmetic formulations. PEG-7 glyceryl cocoate has the most reported uses, i.e., 858, followed by PEG-6 caprylic/capric glycerides with 548 reported uses; the majority of uses for both of these ingredients are in rinse-off formulations. The highest concentrations of use reported for products resulting in leave-on dermal exposure are 11.3% and 6% PEG-7 glyceryl cocoate in tonics, dressings, and other hair grooming aids and other fragrance preparations. PEG-7 glyceryl cocoate also has the highest rinse-off concentration of use reported, i.e., 10% in skin cleansing products.

Since the original safety assessment was published, the frequency of use of PEG-7 glyceryl cocoate increased from 173 reported uses in 1996 to 858 reported uses in 2014. At the time of the original safety assessment, concentration of use data were not available from the FDA. However, based on ocular and sensitization data, a concentration limit of 10% was placed on leave-on products containing PEG glyceryl cocoates. The current reported concentration of use in leave-on products is slightly higher, i.e., 11.3% PEG-7 glyceryl cocoate in tonics, dressings, and other hair grooming aids.

PEG-8 caprylic/capric glycerides can be a dermal penetration enhancer.

In rats, PEG-6 caprylic/capric glycerides had an oral LD₅₀ of >5 g/kg, PEG-8 caprylic/capric glycerides had an oral LD₅₀ of 22 g/kg, and PEG-15 glyceryl isostearate has an oral LD₅₀ of 29.76 ml/kg. In a 4-wk study in rats, the NOAEL for a blend of 40% PEG-8 caprylic/capric glycerides, 40% apricot kernel oil PEG-6 esters, and 20% ethoxydiglycol was 5 ml/kg/day. Test-article related effects were reported in the kidneys, livers, and gastrointestinal systems of animals dosed with 20 ml/g/day; effects were also observed in animals given 10 ml/kg/day, but they did not occur at the same rate of incidence. In a 4-wk dietary study in rats, the NOEL of a formulation containing 0.8% PEG-7 glyceryl cocoate was 2.5%; at higher concentrations, soft feces, a distended cecum, and enlarged mesenteric lymph nodes were observed, and a decrease in spleen weights was reported. No adverse effects were observed in dogs that were dosed orally for 13 wks with 1.0 g/kg/day PEG-8 caprylic/capric glycerides.

PEG-8 caprylic/capric glycerides, up to 3000 mg/kg in purified water, did not produce embryotoxicity, fetotoxicity or teratogenicity. The maternal embryo/fetal NOAELs were 2000 and 3000 mg/kg/day, respectively.

PEG-6 caprylic/capric glycerides, PEG-7 glyceryl cocoate, and PEG-10 olive glycerides were not mutagenic in the Ames test.

The irritation and sensitization potentials of several PEGylated alkyl glycerides were evaluated in alternative, non-human, and human studies, and these ingredients generally were not irritants or sensitizers in the alternative and human studies. Mixed results were observed in the non-human studies; specifically, undiluted PEG-20 almond glycerides, PEG-60 almond glycerides, PEG-12 palm kernel glycerides, and PEG-45 palm kernel glycerides were irritating to rabbit skin in primary skin irritation studies, and *PEG-7 glyceryl cocoate was mildly irritating to rabbit skin in one study*. The only reactions reported in clinical testing was PEG-10 olive glycerides (2% active matter) was non-irritating to mildly irritating. In other clinical studies, up to 20% PEG-60 almond glyceride, undiluted PEG-6 caprylic/capric glycerides, undiluted PEG-7 glyceryl cocoate, and a formulation containing 1% PEG-75 shea butter glycerides were not irritants or sensitizers.

In a HET-CAM, a formulation containing 11% PEG-7 glyceryl cocoate had practically no ocular irritation potential, and concentrations of 10, 50, and 100% PEG-7 glyceryl cocoate were slightly irritating. PEG-10 sunflower glyceride was classified as a non-irritant in a CAMVA. In rabbits, formulations containing 4% PEG-6 caprylic/capric glycerides and up to 3% PEG-8 caprylic/capric glycerides were not ocular irritants and 5% PEG-8 caprylic/capric glycerides was a slight irritant. Undiluted PEG-6 caprylic/capric glycerides was a strong ocular irritant; however, aq. solutions of up to 5%, and formulations containing 4% PEG-6 caprylic/capric glycerides were not ocular irritants in rabbits. In clinical testing, a formulation containing 1% PEG-75 shea butter did not induce ocular irritation in a tolerance test.

PEG-10 olive glycerides, 3% active matter, did not irritate the mucous membrane. In a use study, a female hygiene product containing 0.1% PEG-75 shea butter glycerides did not induce any evidence of vaginovulvar irritation.

Data on the carcinogenicity of PEGylated alkyl glycerides were not found in the published literature, nor were unpublished data provided.

DISCUSSION

Five of the ingredients included in this review, i.e. PEG-7 glyceryl cocoate, PEG-30 glyceryl cocoate, PEG-40 glyceryl cocoate, PEG-78 glyceryl cocoate, and PEG-80 glyceryl cocoate, were reviewed previously; in 1999, the Panel concluded these five ingredients are safe as used in rinse-off products and safe at up to 10% in leave-on products. The safety assessment of these ingredients was scheduled for re-review this year; however, the Panel found it appropriate to include the PEG glyceryl cocoates in this report.

The Panel used the 1999 PEG glyceryl cocoates report, as well as the 2012 safety assessment of the PEGylated oils and the findings on skin irritation and sensitization included in this current report, in its assessment of the safety of the PEGylated alkyl glycerides.

There are no data on skin penetration by the PEGylated alkyl glycerides, but many are approved for use as excipients for pharmaceuticals and food additives, and therefore there is little concern for systemic toxicity; thus, the Panel focused its review on the safety of topical application. Some ingredients demonstrated irritation, but at concentrations above those used in cosmetic formulations; however because the potential exists for dermal irritation with the use of products formulated using PEGylated alkyl glycerides, the Panel specified that products containing these ingredients must be formulated to be non-irritating.. It should be noted that although the conclusion of the 1999 report of PEG glyceryl cocoates limited their use to 10% in leave-on products, data contained in this report alleviated the need for that concentration limit.

The Panel recognized that these ingredients can enhance the penetration of other ingredients through the skin. The Panel cautioned that care should be taken in formulating cosmetic products that may contain these ingredients in combination with any ingredients whose safety was based on their lack of dermal absorption data, or when dermal absorption was a concern.

The Panel noted that some of the PEGylated alkyl glycerides are used in products that could be incidentally inhaled. Although reported concentrations of use included 6% PEG-7 glyceryl cocoate in "other" fragrance preparations and 2% PEG-7 glyceryl cocoate in a pump spray deodorant, and there were no inhalation data available, the Panel was not concerned with

the use of these ingredients in such formulations. The Panel noted that in aerosol products, 95% – 99% of droplets/particles would not be respirable to any appreciable amount. Furthermore, droplets/particles deposited in the nasopharyngeal or bronchial regions of the respiratory tract present no toxicological concerns based on the chemical and biological properties of these ingredients. Coupled with the small actual exposure in the breathing zone and the concentrations at which the ingredients are used, the available information indicates that incidental inhalation would not be a significant route of exposure that might lead to local respiratory or systemic effects. A detailed discussion and summary of the Panel's approach to evaluating incidental inhalation exposures to ingredients in cosmetic products is available at <http://www.cir-safety.org/cir-findings>.

Finally, because these ingredients are obtained from plant sources, the Panel expressed concern regarding pesticide residues and heavy metals that may be present in botanical ingredients. The Panel stressed that the cosmetics industry should continue to use current good manufacturing practices to limit these impurities in the ingredient before blending into cosmetic formulation.

CONCLUSION

The CIR Expert Panel concluded the following 60 PEGylated alkyl glycerides are safe in the present practices of use and concentration in cosmetics when formulated to be non-irritating:

PEG-6 Almond Glycerides*	PEG-6 Hydrogenated Palm/Palm Kernel Glyceride*
PEG-20 Almond Glycerides	PEG-16 Macadamia Glycerides
PEG-35 Almond Glycerides*	PEG-70 Mango Glycerides
PEG-60 Almond Glycerides	PEG-13 Mink Glycerides*
PEG-192 Apricot Kernel Glycerides	PEG-25 Moringa Glycerides*
PEG-11 Avocado Glycerides*	PEG-42 Mushroom Glycerides*
PEG-14 Avocado Glycerides*	PEG-2 Olive Glycerides*
PEG-11 Babassu Glycerides*	PEG-6 Olive Glycerides*
PEG-42 Babassu Glycerides*	PEG-7 Olive Glycerides*
PEG-4 Caprylic/Capric Glycerides*	PEG-10 Olive Glycerides
PEG-6 Caprylic/Capric Glycerides	PEG-40 Olive Glycerides*
PEG-7 Caprylic/Capric Glycerides	PEG-18 Palm Glycerides*
PEG-8 Caprylic/Capric Glycerides	PEG-12 Palm Kernel Glycerides*
PEG-11 Cocoa Butter Glycerides*	PEG-45 Palm Kernel Glycerides
PEG-75 Cocoa Butter Glycerides	PEG-60 Passiflora Edulis Seed Glycerides*
PEG-7 Cocoglycerides*	PEG-60 Passiflora Incarnata Seed Glycerides*
PEG-9 Cocoglycerides*	PEG-45 Safflower Glycerides*
PEG-20 Corn Glycerides*	PEG-60 Shea Butter Glycerides
PEG-60 Corn Glycerides*	PEG-75 Shea Butter Glycerides
PEG-20 Evening Primrose Glycerides*	PEG-75 Shorea Butter Glycerides*
PEG-60 Evening Primrose Glycerides*	PEG-35 Soy Glycerides
PEG-3 Glyceryl Cocoate	PEG-75 Soy Glycerides*
PEG-7 Glyceryl Cocoate	PEG-2 Sunflower Glycerides*
PEG-30 Glyceryl Cocoate	PEG-7 Sunflower Glycerides*
PEG-40 Glyceryl Cocoate	PEG-10 Sunflower Glycerides
PEG-78 Glyceryl Cocoate*	PEG-13 Sunflower Glycerides
PEG-80 Glyceryl Cocoate	PEG-5 Tsubakiate Glycerides*
PEG-5 Hydrogenated Corn Glycerides*	PEG-10 Tsubakiate Glycerides*
PEG-8 Hydrogenated Fish Glycerides*	PEG-20 Tsubakiate Glycerides*
PEG-20 Hydrogenated Palm Glycerides	PEG-60 Tsubakiate Glycerides*

**Not reported to be in current use. Were ingredients in this group not in current use to be used in the future, the expectation is that they would be used in product categories and at concentrations comparable to others in this group.*

TABLES**Table 1. Definitions and functions of the ingredients in this safety assessment**

Ingredient (CAS No.)	Definition ¹	Function(s) ¹
PEG-6 Almond Glycerides [124046-50-0; generic]	a polyethylene glycol derivative of the mono- and diglycerides from almond oil with an average of 6 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from almond oil, where at least one R (and no more than two) is a fatty acid, and the sum of all n is 6.</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-20 Almond Glycerides [124046-50-0; generic]	a polyethylene glycol derivative of the mono- and diglycerides from almond oil with an average of 20 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from almond oil, where at least one R (and no more than two) is a fatty acid, and the sum of all n is 20</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-35 Almond Glycerides 124046-50-0 [generic]	a polyethylene glycol derivative of the mono- and diglycerides from almond oil with an average of 35 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from almond oil, where at least one R (and no more than two) is a fatty acid, and the sum of all n is 35</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-60 Almond Glycerides 124046-50-0 [generic]	a polyethylene glycol derivative of the mono- and diglycerides from almond oil with an average of 60 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from almond oil, where at least one R (and no more than two) is a fatty acid, and the sum of all n is 60.</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-192 Apricot Kernel Glycerides	a polyethylene glycol derivative of the mono- and diglycerides from apricot kernel oil with an average of 192 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from apricot kernel oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 192</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-11 Avocado Glycerides 103819-44-9 (generic)	a polyethylene glycol derivative of mono- and diglycerides from avocado oil with an average of 11 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from avocado oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 11</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-14 Avocado Glycerides 103819-44-9 (generic)	a polyethylene glycol derivative of mono- and diglycerides from avocado oil with an average of 14 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from avocado oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 14.</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-11 Babassu Glycerides	a polyethylene glycol derivative of the mono- and diglycerides derived from babassu oil with an average of 11 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from babassu oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 11</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-42 Babassu Glycerides	a polyethylene glycol derivative of the mono- and diglycerides derived from babassu oil with an average of 42 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from babassu oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 42.</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-4 Caprylic/Capric Glycerides	a polyethylene glycol derivative of a mixture of mono-, di-, and triglycerides of caprylic and capric acids with an average of 4 moles of ethylene oxide; <i>wherein the sum of all cases of n in Figure 2 is 4.</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-6 Caprylic/Capric Glycerides 361459-38-3 ⁸⁰	a polyethylene glycol derivative of a mixture of mono-, di-, and triglycerides of caprylic and capric acids with an average of 6 moles of ethylene oxide; <i>wherein the sum of all cases of n in Figure 2 is 6.</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-7 Caprylic/Capric Glycerides	a polyethylene glycol derivative of a mixture of mono-, di-, and triglycerides of caprylic and capric acids with an average of 7 moles of ethylene oxide; <i>wherein the sum of all cases of n in Figure 2 is 7.</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-8 Caprylic/Capric Glycerides	a polyethylene glycol derivative of a mixture of mono-, di-, and triglycerides of caprylic and capric acids with an average of 8 moles of ethylene oxide; <i>wherein the sum of all cases of n in Figure 2 is 8.</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-11 Cocoa Butter Glycerides	a polyethylene glycol derivative of the mono- and diglycerides derived from theobroma cacao (cocoa) seed butter with an average of 11 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from theobroma cacao (cocoa) seed butter, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 11</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-75 Cocoa Butter Glycerides	a polyethylene glycol derivative of the glycerides derived from theobroma cacao (cocoa) seed butter with an average of 75 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from theobroma cacao (cocoa) seed butter, where at least one R is a fatty acid, and the sum of all cases of n is 75</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-7 Cocoglycerides [67762-35-0; generic]	the polyethylene glycol derivative of the mono- and diglycerides of coconut oil with an average ethoxylation value of 7; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from coconut oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 7</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent

Table 1. Definitions and functions of the ingredients in this safety assessment

Ingredient (CAS No.)	Definition ¹	Function(s) ¹
PEG-9 Cocoglycerides 67762-35-0 (generic)	the polyethylene glycol derivative of the mono- and diglycerides of coconut oil with an average ethoxylation value of 9; wherein R in Figure 1 is hydrogen or the fatty acid residues from coconut oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 9	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-20 Corn Glycerides	a polyethylene glycol derivative of corn glycerides with an average of 20 moles of ethylene oxide; wherein R in Figure 1 is hydrogen or the fatty acid residues from ze mays (corn) oil, where at least one R is a fatty acid, and the sum of all cases of n is 20 <i>Corn Glycerides is a mixture of mono, di and triglycerides derived from Zea Mays (Corn) Oil.</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-60 Corn Glycerides	PEG-60 Corn Glycerides is a polyethylene glycol derivative of Corn Glycerides with an average of 60 moles of ethylene oxide; wherein R in Figure 1 is hydrogen or the fatty acid residues from ze mays (corn) oil, where at least one R is a fatty acid, and the sum of all cases of n is 60 <i>Corn Glycerides is a mixture of mono, di and triglycerides derived from Zea Mays (Corn) Oil.</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent; surfactant – solubilizing agent
PEG-20 Evening Primrose Glycerides	a polyethylene glycol derivative of the mono and diglycerides from evening primrose oil with an average of 20 moles of ethylene oxide; wherein R in Figure 1 is hydrogen or the fatty acid residues from evening primrose oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 20	skin-conditioning agent - emollient
PEG-60 Evening Primrose Glycerides	a polyethylene glycol derivative of the mono and diglycerides from evening primrose oil with an average of 60 moles of ethylene oxide; wherein R in Figure 1 is hydrogen or the fatty acid residues from evening primrose oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 60	surfactant – emulsifying agent; surfactant – solubilizing agent
PEG-3 Glyceryl Cocoate	the polyethylene glycol ether of glyceryl cocoate that conforms generally to the structure below; the polyethylene glycol derivative of the monoglyceride of coconut oil with an average ethoxyla-tion value of 3	skin-conditioning agent - emollient; surfactant – emulsifying agent
	<p style="text-align: center;">wherein RC(O) represents the fatty acid residues from coconut oil, and the sum of all cases of n is 3.</p>	
PEG-7 Glyceryl Cocoate 66105-29-1 68201-46-7 (generic)	the polyethylene glycol ether of glyceryl cocoate that conforms generally to the structure below; the polyethylene glycol derivative of the monoglyceride of coconut oil with an average ethoxyla-tion value of 7	skin-conditioning agent - emollient; surfactant – emulsifying agent
	<p style="text-align: center;">wherein RC(O) represents the fatty acid residues from coconut oil, and the sum of all cases of n is 7.</p>	
PEG-30 Glyceryl Cocoate 68201-46-7(generic)	the polyethylene glycol ether of glyceryl cocoate that conforms generally to the structure below; the polyethylene glycol derivative of the monoglyceride of coconut oil with an average ethoxyla-tion value of 30	surfactants - cleans-ing agent; surfac-tants - solubilizing agent
	<p style="text-align: center;">wherein RC(O) represents the fatty acid residues from coconut oil, and the sum of all cases of n is 30.</p>	

Table 1. Definitions and functions of the ingredients in this safety assessment




Ingredient (CAS No.)	Definition ¹	Function(s) ¹
PEG-40 Glyceryl Cocoate 68201-46-7(generic)	the polyethylene glycol ether of glyceryl cocoate that conforms generally to the structure below; the polyethylene glycol derivative of the monoglyceride of coconut oil with an average ethoxyla- tion value of 40	surfactants - cleans- ing agent; surfac- tants - solubilizing agent
 <p>wherein $RC(O)$ represents the fatty acid residues from coconut oil, and the sum of all cases of n is 40.</p>		
PEG-78 Glyceryl Cocoate 68201-46-7(generic)	the polyethylene glycol ether of glyceryl cocoate that conforms generally to the structure below; the polyethylene glycol derivative of the monoglyceride of coconut oil with an average ethoxyla- tion value of 78	surfactants - cleans- ing agent; surfac- tants - solubilizing agent
 <p>wherein $RC(O)$ represents the fatty acid residues from coconut oil, and the sum of all cases of n is 78.</p>		
PEG-80 Glyceryl Cocoate 68201-46-7(generic)	the polyethylene glycol ether of glyceryl cocoate that conforms generally to the structure below; the polyethylene glycol derivative of the monoglyceride of coconut oil with an average ethoxyla- tion value of 80	surfactants - cleans- ing agent; surfac- tants - solubilizing agent
 <p>wherein $RC(O)$ represents the fatty acid residues from coconut oil, and the sum of all cases of n is 80.</p>		
PEG-5 Hydrogenated Corn Glycerides	the polyethylene glycol derivative of mixed glycerides derived from hydrogenated corn oil. It has an average of 5 moles of ethylene oxide; wherein R in Figure 1 is hydrogen or the fatty acid residues from hydrogenated corn oil, where at least one R is a fatty acid, and the sum of all cases of n is 5	skin-conditioning agent - emollient; surfactant – emulsi- fying agent
PEG-8 Hydrogenated Fish Glycerides	a polyethylene glycol derivative of hydrogenated fish oil with an average of 8 moles of ethylene oxide; wherein R in Figure 1 is hydrogen or the fatty acid residues from hydrogenated fish oil, where at least one R is a fatty acid, and the sum of all cases of n is 8	skin-conditioning agent - emollient; surfactant – emulsi- fying agent
PEG-20 Hydrogenated Palm Glycerides	a polyethylene glycol ether of hydrogenated palm glycerides with an average of 20 moles of ethylene oxide; wherein R in Figure 1 is hydrogen or the fatty acid residues from hydrogenated palm glycerides, where at least one R is a fatty acid, and the sum of all cases of n is 20. (Hy- drogenated palm glyceride is the end product of the controlled hydrogenation of palm glycer- ide.)	skin-conditioning agent - emollient; surfactant – emulsi- fying agent
PEG-6 Hydrogenated Palm/Palm Kernel Gly- ceride	a polyethylene glycol derivative of a mixture of hydrogenated palm glyceride and hydrogenated palm kernel glyceride containing an average of 6 moles of ethylene oxide; wherein R in Figure 1 is hydrogen or the fatty acid residues from hydrogenated palm glycerides or hydrogenated palm kernel glycerides, where at least one R is a fatty acid, and the sum of all cases of n is 6 (Palm kernel glycerides is a mixture of mono, di and triglycerides derived from <i>Elaeis guineen- sis</i> (palm) kernel oil.)	skin-conditioning agent - emollient; surfactant – emulsi- fying agent
PEG-16 Macadamia Gly- cerides	the polyethylene glycol derivative of the mono- and diglycerides derived from macadamia nut oil with an average of 16 moles of ethylene oxide; wherein R in Figure 1 is hydrogen or the fatty acid residues from macadamia nut oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 16	skin-conditioning agent - emollient; surfactant – emulsi- fying agent

Table 1. Definitions and functions of the ingredients in this safety assessment

Ingredient (CAS No.)	Definition ¹	Function(s) ¹
PEG-70 Mango Glycerides	a polyethylene glycol derivative of the mono- and diglycerides from mango seed oil containing an average of 70 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from mango seed oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 70</i>	skin-conditioning agent - emollient; surfactant – cleansing agent; surfactant – solubilizing agent
PEG-13 Mink Glycerides 103819-45-0	a polyethylene glycol derivative of mono and diglycerides derived from mink oil with an average of 13 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from mink oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 13</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-25 Moringa Glycerides	a polyethylene glycol derivative of the fatty oil obtained from the seeds of <i>Moringa pterygosperma</i> containing an average of 25 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from Moringa pterygosperma seed oil, where at least one R is a fatty acid, and the sum of all cases of n is 25</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-42 Mushroom Glycerides	the polyethylene glycol derivative of the mono- and diglycerides of mushroom oil with an average of 42 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from mushroom oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 42</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-2 Olive Glycerides 103819-46-1 [generic]	a polyethylene glycol derivative of mono- and diglycerides derived from olive oil with an average of 2 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from olive oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 2</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-6 Olive Glycerides 103819-46-1 [generic]	a polyethylene glycol derivative of mono- and diglycerides derived from olive oil with an average of 6 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from olive oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 6</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-7 Olive Glycerides [103819-46-1; generic]	a polyethylene glycol derivative of mono- and diglycerides derived from olive oil with an average of 7 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from olive oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 7</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-10 Olive Glycerides [103819-46-1; generic]	a polyethylene glycol derivative of mono- and diglycerides derived from olive oil with an average of 10 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from olive oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 10</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-40 Olive Glycerides [103819-46-1; generic]	a polyethylene glycol derivative of mono- and diglycerides derived from olive oil with an average of 40 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from olive oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 40</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-18 Palm Glycerides	is a polyethylene glycol derivative of palm glycerides with an average of 18 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from palm glycerides, where at least one R is a fatty acid, and the sum of all cases of n is 18 (Palm glycerides is a mixture of mono, di and triglycerides derived from Elaeis guineensis (palm) oil)</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-12 Palm Kernel Glycerides 124046-52-2 [generic]	a polyethylene glycol derivative of the mono and diglycerides of palm kernel oil with an average of 12 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from palm kernel oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 12</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-45 Palm Kernel Glycerides 68606-12-2; ⁸¹ 124046-52-2 [generic]	a polyethylene glycol derivative of the mono and diglycerides of palm kernel oil with an average of 45 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from palm kernel oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 45</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-60 Passiflora Edulis Seed Glycerides	the polyethylene glycol derivative of the mono- and diglycerides derived from the oil of the seeds of <i>Passiflora edulis</i> containing an average of 60 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from Passiflora edulis seed oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 60</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-60 Passiflora Incarnata Seed Glycerides	the polyethylene glycol derivative of the mono- and diglycerides derived from the oil of the seeds of <i>Passiflora incarnata</i> ; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from Passiflora incarnata seed oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 60</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-45 Safflower Glycerides	a polyethylene glycol derivative of the mono- and diglycerides derived from safflower oil with an average of 45 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from safflower oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 45</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-60 Shea Butter Glycerides	a polyethylene glycol derivative of the glycerides derived from butyrospermum parkii (shea) butter with an average ethoxylation value of 60; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from oil obtained from butyrospermum parkii (shea) butter, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 60</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent; surfactant – solubilizing agent

Table 1. Definitions and functions of the ingredients in this safety assessment

Ingredient (CAS No.)	Definition ¹	Function(s) ¹
PEG-75 Shea Butter Glycerides	a polyethylene glycol derivative of the glycerides derived from butyrospermum parkii (shea) butter with an average ethoxylation value of 75; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from oil obtained from butyrospermum parkii (shea) butter, where at least one R is a fatty acid, and the sum of all cases of n is 75</i>	skin-conditioning agent - emollient; surfactant – cleansing agent; surfactant – solubilizing agent
PEG-75 Shorea Butter Glycerides	a polyethylene glycol derivative of the glycerides derived from shorea stenoptera seed butter with an average of 75 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from oil obtained from shorea stenoptera seed butter, where at least one R is a fatty acid, and the sum of all cases of n is 75</i>	skin-conditioning agent - emollient; surfactant – cleansing agent; surfactant – solubilizing agent
PEG-35 Soy Glycerides 61791-23-9 ⁸²	a polyethylene glycol derivative of the mono- and diglycerides derived from glycine soja (soybean) oil containing an average of 35 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from glycine soja (soybean) oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 35</i>	skin-conditioning agent – miscellaneous; surfactant – emulsifying agent; surfactant – solubilizing agent
PEG-75 Soy Glycerides	a polyethylene glycol derivative of the mono- and diglycerides derived from glycine soja (soybean) oil containing an average of 75 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from glycine soja (soybean) oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 75</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-2 Sunflower Glycerides 180254-52-8 (generic) 186511-05-7 (generic)	a polyethylene glycol derivative of the mono- and diglycerides derived from sunflower seed oil with an average of 2 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from sunflower seed oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 2</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-7 Sunflower Glycerides [180254-52-8; generic 186511-05-7; generic]	a polyethylene glycol derivative of the mono- and diglycerides derived from sunflower seed oil with an average of 7 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from sunflower seed oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 7</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-10 Sunflower Glycerides 180254-52-8 (generic) 186511-05-7 (generic)	a polyethylene glycol derivative of the mono- and diglycerides derived from sunflower seed oil with an average of 10 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from sunflower seed oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 10</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-13 Sunflower Glycerides 186511-05-7 (generic) 70377-91-2 (generic)	a polyethylene glycol derivative of the mono- and diglycerides derived from sunflower seed oil with an average of 13 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from sunflower seed oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 13</i>	skin-conditioning agent - emollient; surfactant – emulsifying agent
PEG-5 Tsubakiate Glycerides	the polyethylene glycol derivative of the mono- and diglycerides derived from camellia japonica seed oil containing an average of 5 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from camellia japonica seed oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 5</i>	surfactant – emulsifying agent
PEG-10 Tsubakiate Glycerides	the polyethylene glycol derivative of the mono- and diglycerides derived from camellia japonica seed oil containing an average of 10 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from camellia japonica seed oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 10</i>	surfactant – emulsifying agent
PEG-20 Tsubakiate Glycerides	the polyethylene glycol derivative of the mono- and diglycerides derived from camellia japonica seed oil containing an average of 20 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from camellia japonica seed oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 20</i>	surfactant – emulsifying agent
PEG-60 Tsubakiate Glycerides	the polyethylene glycol derivative of the mono- and diglycerides derived from camellia japonica seed oil containing an average of 60 moles of ethylene oxide; <i>wherein R in Figure 1 is hydrogen or the fatty acid residues from camellia japonica seed oil, where at least one R (and no more than two) is a fatty acid, and the sum of all cases of n is 60</i>	surfactant – cleansing agent

Table 2. Summaries of previous CIR safety assessments

Ingredient(s)	Conclusion	Summary	Reference
Triethylene Glycol and Polyethylene Glycols (PEGs) ≥ 4	Safe for use in cosmetics in the present practices of use and concentration	In general, PEGs are not oral toxicants, with acute oral LD ₅₀ values in rodents ranging from 15-22 g/kg. The LC ₅₀ of aerosolized triethylene glycol in rats was greater than 3.9 mg/L. Minimal dermal irritation and sensitization with undiluted PEGs has been observed. Ocular exposure to triethylene glycol in rabbits produced no corneal injury, however all rabbits displayed acute iritis and minor transient conjunctival irritation; overall, PEGs cause mild, transient ocular irritation in rabbits. PEGs are not genotoxic (up to ~5700 mg/kg/ day in a	⁴

Table 2. Summaries of previous CIR safety assessments

Ingredient(s)	Conclusion	Summary	Reference
		rat dominant lethal assay) or carcinogenic (up to 0.25 ml/week in a rat subcutaneous injection study). PEGs are not reproductive or developmental toxicants at doses up to 6.78 g/kg in mice. Use of antimicrobial creams with a PEG vehicle was associated with renal toxicity when applied to burned skin, but studies of extensively tape stripped skin demonstrated that the levels of PEGs that could penetrate in a worst case analysis are >100 times less than the renal toxicity no observable effect level of 1.1 g/kg/day, providing a margin of safety.	
Plant-Derived Fatty Acid Oils includes: almond oil; apricot kernel oil; avocado oil; babassu oil; <i>Theobroma cacao</i> (cocoa) seed butter; coconut oil; evening primrose oil; moringa oleifera seed oil; moringa pterygosperma seed oil; olive oil; palm oil; palm kernel oil; safflower oil; <i>Butyrospermum parkii</i> (shea) butter; soybean oil; <i>Carthamus tinctorius</i> (safflower) seed oil; sunflower seed oil; <i>Camellia japonica</i> seed oil	Safe for use in cosmetics in the present practices of use and concentration	Oils are used in a wide variety of cosmetic products for their skin conditioning, occlusive, emollient, and moisturizing properties. Undiluted, technical grade, <i>Arachis hypogaea</i> (Peanut) Oil was moderately irritating to rabbits and guinea pig skin, and 5% aq. solutions of a bar soap containing 13% sodium cocoate had irritation scores of 1.6-4.0/8 in animal studies. However, the remaining animal and clinical irritation and/or sensitization studies conducted on a large number of the oils included in the safety assessment report, primarily in formulation, did not report any significant irritation or sensitization reactions, indicating that refined oils derived from plants are not ocular or dermal irritants or sensitizers.	5
Caprylic/Capric Triglyceride	Safe as used (maximum concentration of use, >50% in eye products, perfumes, and other dermal products)	Caprylic/capric triglyceride is an oily mixed ester predominantly composed of caprylic and capric fatty acids derived from coconut oil. When absorbed from the digestive tract, it is hydrolyzed, and the fatty acids are catabolized to C2 fragments which may be further metabolized either to CO ₂ or to form long-chain fatty acids. It has a very low toxicity to man and animals as shown by tests involving oral ingestion, intraperitoneal and intramuscular injection, skin and eye irritation tests, skin sensitization, percutaneous toxicity and finally, by two generation feeding studies. Application of undiluted test material to rabbits produced no to mild irritation. A 4% solution in ethanol was not a sensitizer in guinea pigs.	6
Coconut Oil includes cocoglycerides	see Plant-Derived Fatty Acid Oils: safe as used (maximum concentration of use, 80% coconut oil in eye lotion; 14% cocoglycerides in lipstick)	The genotoxic potential of saponified coconut oil was evaluated in several prokaryote systems; the cytotoxic, anti-oxidant, and mutagenic effects of saponified coconut oil can be influenced by the aggregational state. Coconut oil, 100%, was not an allergen in a double-blind randomized study in 12 subjects.	7
Corn Oil includes corn glycerides	Safe as used (maximum use, 14% corn oil in dermal products; corn glycerides had no reported uses)	Corn oil is commonly used as a vehicle in studies of lipid-soluble substances carried out in numerous animal species; corn oil was not found to have effect when evaluated as a vehicle/control.	8
Palm Oil includes palm kernel oil	see Plant-Derived Fatty Acid Oils	Undiluted palm oil has an oral LD ₅₀ in rats of > 5g/kg. Short-term and subchronic feeding studies showed no evidence of toxicity. Chronic feeding studies produced results suggestive of metabolic hyperactivity. Minimal ocular irritation and no skin irritation (tested undiluted), sensitization (tested at 5%), or photosensitization (1.5% in formulation) were reported in animals studies. Anomalies in 30% of the live fetuses delivered by female albino rats fed commercial grade palm oil were reported; other studies including multigenerational tests of crude palm oil and heated palm oil (as would occur in cooking) reported no reproductive toxicity, developmental toxicity, or differences in endocrine function. Although some data show that palm oil can be mutagenic in certain Ames test <i>Salmonella</i> strains, it was negative in other strains and negative in an assay of chromosomal aberrations in bone marrow samples taken from mice dosed orally. Several studies suggesting an inhibitory effect of palm oil on 7,12-dimethylbenz(o)anthracene tumorigenesis have attributed the effect to the high vitamin E content of the oil. There was no evidence of irritation or sensitization in clinical tests; use testing of products containing palm oil produced no ocular or skin irritation.	9
Mink Oil	Safe for use in cosmetics in the	Mink oil does not absorb significant UVA or UVB	10

Table 2. Summaries of previous CIR safety assessments

Ingredient(s)	Conclusion	Summary	Reference
	present practices of use and concentration	radiation. It is not an acute oral toxicant (LD ₅₀ > 64 cc/kg) in rats. Non-human test data indicate that mink oil is not a dermal or ocular irritant or a dermal sensitizer up to 50%. No irritation was observed in clinical studies of mink oil up to 2.8%, although some transient irritation was noted in exaggerated-use studies. Because mink oil is a mixture of glycerides, namely triglycerides, it may enhance the penetration of other chemicals.	
Safflower Oil	see Plant-Derived Fatty Acid Oils	The pure oil produced slight to moderate comedogenicity. However, products containing up to 5 percent Safflower Oil were not comedogenic in rabbits. Results of animal tests indicated that undiluted safflower oil was not an eye or skin irritant or contact sensitizer. The oil increased the incidence of mammary tumors induced by 12-dimethylbenz(a)anthracene and large bowel tumors induced by dimethylhydrazine in rats.	11
PEGylated Oils	PEGylated oils are safe in the present practices of use and concentration in cosmetics when formulated to be non-irritating. (This conclusion supersedes the earlier conclusion for PEG castor oils issued by the Expert Panel in 1997.)	PEGylated oils are mixtures that are the condensation products of ethylene oxide and water, with the chain length controlled by number of moles of ethylene oxide that are polymerized. PEGs may contain trace amounts of 1,4-dioxane, a by-product of ethoxylation. PEG-40 hydrogenated castor oil had a maximum use concentration 22% in leave-on non-coloring hair products. PEG-30 castor oil and PEG-40 hydrogenated castor oil may be used as nonionic surfactants in oral, topical, and parenteral drug delivery systems. PEG-30 and -40 hydrogenated castor oil are approved as direct and indirect food additives. Various types of PEGs are hydrophilic penetration enhancers and are used in topical dermatological preparations. Unspecified PEG castor oil did not elicit irritation at the site of injection in rabbits when evaluated as a vehicle in an intravenous drug. PEG-60 hydrogenated castor oil produced toxicity, including decreased blood pressure, flush, swelling, itching, and increase histamine levels, in dogs injected intravenously with a single 1.25, 2.5 or 10 mg/kg dose of the test material. No signs of toxicity were observed with PEG-60 hydrogenated castor oil in monkeys, rabbits, guinea pigs or rats. Several rat studies investigated the effects of the use of PEG-35 castor oil as a vehicle in intravenous drugs; 100 µg/L had effects on cardiac and skeletal muscle function. PEG-60 hydrogenated castor oil was not genotoxic in a reverse mutation test, in a chromosome aberration study, and in a mouse micronucleus test. A microemulsion containing 20% w/w PEG-35 castor oil was not an irritation in guinea pigs. A formulation containing 20% PEG-40 hydrogenated castor oil tested in mouse skin would probably not irritate human skin. A formulation containing 20.66% PEG-40 hydrogenated castor oil was not a skin irritant. Nanoemulsions containing up to 13.5% PEG-35 castor oil were non-irritating in rabbit eyes.	3

Table 3. Chemical and Physical Properties

Property	Description	Reference
PEG-11 Avocado Glycerides		
physical appearance	brown liquid	83
	yellow, liquid to pasty	84
	yellowish to brown liquid with characteristic odor	85
solubility	soluble in water	84
pH value (10% in water; 25°C)	5.0-7.0	85
saponification value	60-80 mg KOH/g	85
iodine value	25-37 g iodine/100 g	85
surfactant-type	non-ionic	84
PEG-20 Almond Glycerides		
physical appearance	yellow liquid with a characteristic odor	86

Table 3. Chemical and Physical Properties

Property	Description	Reference
PEG-42 Babassu Glycerides		
physical appearance	clear to slightly hazy yellow liquid	87
PEG-6 Caprylic/Capric Glycerides		
physical appearance	clear yellow liquid	80,88
solubility (25°C; 10% concentration)	soluble in water, cosmetic alcohol, 1,2-propylene glycol; soluble in sunflower seed oil (slight turbidity); insoluble in paraffin oil and isopropyl myristate; dispersible in glycerol	89
hydroxyl value	200-220 mg KOH/g	24
iodine value	≤1 g I/100 g	24
saponification value	85-105 mg KOH/g	24
acid value	<2	80,88
PEG-8 Caprylic/Capric Glycerides		
physical appearance	pale yellow oily liquid with a faint odor	20,26
mean molecular wt	200-400	19,20
solubility	very soluble in ethanol (96°), chloroform, methylene chloride; soluble in water; insoluble in mineral oils freely soluble in methylene chloride; dispersible in hot water	26 20
specific gravity (20°C; D20/4)	1.060 – 1.070	26
refractive index (20°C)	1.450 – 1.470	26
saponification value	85 – 105 mg KOH/g	26
acid value	≤2.0	26,27
fatty acid composition	caproic acid (C6): ≤2.0% caprylic acid (C8): 50-80% capric acid (C10): 20-50% lauric acid (C12): ≤3.0% myristic acid (C14): ≤1.0%	26,27
free glycerol content	≤5%	26
hydrophilic-lipophilic balance (HLB)	14	43
PEG-9 Cocoglycerides		
physical appearance	yellow, liquid to pasty	90
solubility	soluble in water	90
HLB	13	90
surfactant-type	non-ionic	90
PEG-60 Corn Glycerides		
physical appearance	liquid	91
surfactant-type	non-ionic	91
PEG-60 Evening Primrose Glycerides		
physical appearance	soft, pale paste	92
PEG-7 Glyceryl Cocoate		
physical appearance	clear, pale yellow oil with a mild fatty odor	2
solubility	soluble in water and ethanol; insoluble in mineral oil 25°C; 10% concentration: soluble in water (slight turbidity) and cosmetic alcohol; insoluble in paraffin oil, isopropyl myristate, and glycerol; dispersible in sunflower seed oil and 1,2-propylene glycol	2 89
hydrophile-lipophile balance	15	2
hydroxyl value	172-187	2
saponification value	90-100	2
acid value	5.0 max	2
iodine value	5.0 max	2
PEG-16 Macadamia Glycerides		
physical appearance	liquid	93
affinity	hydrophilic	94
HLB	12-14	94
hydroxyl value	90-130 mg KOH/g	93
saponification value	45-60 mg KOH/g	93
PEG-70 Mango Glycerides		
physical appearance	solid	95
solubility	water-dispersible	96
surfactant-type	non-ionic	96
HLB	16-18	95,96
acid value	1.5	96

Table 3. Chemical and Physical Properties

Property	Description	Reference
saponification value	14-20 mg KOH/g	⁹⁶
PEG-10 Olive Glycerides		
physical appearance	yellow, liquid to pasty	⁹⁷
	yellowish-brown paste with characteristic odor	⁹⁸
solubility	soluble in water	⁹⁷
	solubility, 2.5% in water – almost clear to turbid	⁹⁸
	solubility, 10% in isopropyl alcohol - clear	⁹⁸
pH value (10% in water; 25°C)	5.0-7.0	⁹⁸
acid value	max 4.0 mg KOH/g	⁹⁸
saponification value	52-72 mg KOH/g	⁹⁸
iodine value	26-36 g iodine/100 g	⁹⁸
		⁹⁸
affinity	hydrophilic	⁹⁹
surfactant-type	ionic	⁹⁷
HLB	13	⁹⁷
PEG-45 Palm Kernel Glycerides		
physical appearance	clear liquid	⁸¹
surfactant-type	non-ionic	¹⁰⁰
PEG-75 Shea Butter Glycerides		
physical appearance	waxy solid	¹⁰¹
solubility	water-dispersible	⁹⁵
pH	5.5-8.5	¹⁰¹
surfactant-type	non-ionic	⁹⁵
HLB	16-18	¹⁰¹
PEG-35 Soy Glycerides		
physical appearance	solid or paste	⁸²
solubility	soluble in water	⁸²
surfactant-type	non-ionic	⁸²
PEG-10 Sunflower Glycerides		
physical appearance	liquid	¹⁰²
solubility	soluble in water; dispersible in oil	¹⁰³
refractive index	1.460	¹⁰⁴
pH	6.9	¹⁰⁴
surfactant-type	hydrophilic	¹⁰³
HLB	8-10	¹⁰⁴
hydroxyl value	135-160 mg KOH/g	¹⁰²
acid value	0-1.5 mg KOH/g	¹⁰²
saponification value	60-80 mg KOH/g	¹⁰²

Table 4 Total fatty acid composition of plant-derived fatty acid oils and of butters (%)

Fatty Acids	Prunus Amygdalus (Sweet Almond) Oil ⁵	Prunus Armeniaca (Apricot) Kernel Oil ⁵	Persea Gratissima (Avocado) Oil ⁵	Orbignya Oleifera Seed Oil [Babassu] ⁵	Theobroma Cacao (Cocoa) Seed Butter ⁵	Cocos Nucifera (Co- conut) Oil ⁵	Zea Mays (Corn) Oil ⁵
Caproic (C6)						0-1	
Caprylic (C8)				4 to 8		5-9	
Capric (C10)				4 to 8		6-10	
Lauric (C12)*				44 - 47		44-52	
Myristic (C14)	1			15 - 20		13-19	0.1 - 1.7
Myristoleic (C14:1)							
Palmitic (C16)	4-9	4.6-6	13-17	6 to 9	24-29	8-11	8-16.5
Palmitoleic (C16:1)	0.8	1-2	3 - 5.1			0-1	0.2 - 1.6
Heptadecanoic (C17:0)	0.2						
Stearic (C18)	2-3	0.5-1.2		3 to 5	34-36	1-3	0-4.5
Oleic (C18:1)	62-86	58-65.7 (total 18:1)	67-72	10 to 12	30-40	5-8	19 – 49
Linoleic (C18:2)	20-30	29-33	10 to 12	1 to 3	2.4	Trace-2.5	34-66
		28.5 (undef. 18:2)					
Linolenic (C18:3)	0.4	05-1.0 (undef 18:3)					0-2
Arachidic (C20)	0.2	0.2					1
Eicosenoic (C20:1)	0.3						1
Behenic (C22)	0.2						
Erucic (C22:1)	0.1						
Lignoceric (C24)							
Others	<C16:0 = 0.1	Oleic/Linoleic = 90-93%					

	Oenothera Biennis (Evening Primrose) Oil ⁵	Fish Oil	Macadamia Integri- folia Seed Oil ⁵	Mangifera Indica (Mango) Seed Oil ⁵	Mink Oil ¹⁰		Moringa Oleifera Seed Oil	Mushroom Oil
					Mink Wax Oil	Mink Crude Oil		
Caproic (C6)								
Caprylic (C8)								
Capric (C10)								
Lauric (C12)*			0.1-1.4		0.1	0.1		
Myristic (C14)			0.7-1.5		4.0	3.5	trace	
Myristoleic (C14:1)					0.7	0.9		
Pentadecanoic (15/0)					0.2	0.1		
Palmitic (C16)	4-10		6-12	5-8	28.0	17.2	5-9.3	
Palmitoleic (C16:1)			12-25		13.3	17.0	1.5-3	
Heptadecanoic (C17:0)					0.5	0.4		
Heptadecanoic (C17:1)					0.6	0.5		
Stearic (C18)	2-4		0.5-8	33-48	4.7	2.5	3-8	
Oleic (C18:1)	5-12		50-67	35-50	35.3	40.9	65-80	
Linoleic (C18:2)	60-85		1.5-5	4.0-8	10.6	15.0	1.5-5	
Linolenic (C18:3)			0.5-1.9		0.4	0.6	1-1.5	
Arachidic (C20)			1.5-5	1-7	0.1		2-5	
Eicosenoic (C20:1)			1.5-3.1		0.7	0.6	2.5-4	
Behenic (C22)			0.3-1				8-8.6	
Erucic (C22:1)			1				3	
Lignoceric (C24)							trace	
Others	α -Linolenic (C18:3) = 1% γ -Linolenic = 7-12%							

Table 4 Total fatty acid composition of plant-derived fatty acid oils and of butters (%)

Fatty Acids	Olea Europaea (Olive) Oil ⁵	Elaeis Guineensis (Palm) Oil ⁵	Elaeis Guineensis (Palm) Kernel Oil ⁵	Passiflora Edulis Seed Oil ⁵	Passiflora Incarnata Seed Oil ¹⁰⁵	Carthamus Tinctorius (Safflower) Seed Oil ⁵	Butyrospermum Par-kii (Shea) Butter ⁵
Caproic (C6)			0.3				
Caprylic (C8)			4.4				
Capric (C10)			3.7				
Lauric (C12)*		0.2	48.3				
Myristic (C14)	Trace	1.1	15.6	0.03			0.5
Myristoleic (C14:1)							
Palmitic (C16)	7.5 - 20	44	7.8	8.57	8	2	3-9
Palmitoleic (C16:1)	0.3 - 3.5	0.1		0.23			
Heptadecanoic (C17:0)							
Stearic (C18)	0.5 - 3.5	4.5	2	1.66	2		30-50
Oleic (C18:1)	53 - 86	39.2	15.1	16.25	12	26	38-50
Linoleic (C18:2)	3.5 - 20	10.1	2.7	72.69	77	68	3-8
Linolenic (C18:3)	0 - 1.5	0.4		0.26	1	Trace	0.5 max
Arachidic (C20)	Trace	0.4				Trace	2.5-3
Eicosenoic (C20:1)							
Behenic (C22)	Trace						
Erucic (C22:1)							
Lignoceric (C24)	Trace						
Others			0.2	Unspecified other fatty acids = 0.31			

	Shorea Stenoptera Seed Butter ¹⁰⁶	Glycine Soja (Soy-bean) Oil ⁵	Helianthus Annuus (Sunflower) Seed Oil ⁵	Camellia Japonica Seed Oil ⁵
Caproic (C6)				
Caprylic (C8)				
Capric (C10)				
Lauric (C12)*				
Myristic (C14)				
Myristoleic (C14:1)				
Palmitic (C16)	19.5		5.0 - 7.2	7.9
Palmitoleic (C16:1)				0.16
Heptadecanoic (C17:0)				
Stearic (C18)	42.4		2.0 - 6.5	2.46
Oleic (C18:1)	36.9	11.5 - 60.0	14.7 - 37.2	84.99
Linoleic (C18:2)	0.2		51.5 - 73.5	3.76
Linolenic (C18:3)		2.9 - 12.1	Trace - 0.3	
Arachidic (C20)	1.0		0.3 - 1	0.49
Eicosenoic (C20:1)				
Behenic (C22)				
Erucic (C22:1)				
Lignoceric (C24)				
Others				

Table 5. Frequency and concentration of use according to duration and type of exposure

	# of Uses ²⁸	Max. Conc. of Use (%) ^{29,31}	# of Uses ²⁸	Max. Conc. of Use (%) ^{29,31}	# of Uses ²⁸	Max. Conc. of Use (%) ^{29,31}
	PEG-20 Almond Glycerides		PEG-60 Almond Glycerides		PEG-192 Apricot Kernel Glycerides	
Totals*	11	0.01	163	0.0001-6.5	24	0.1
Duration of Use						
Leave-On	3	0.01	75	0.0001-3.2	24	0.1
Rinse Off	8	NR	87	0.001-6.5	NR	NR
Diluted for (Bath) Use	NR	NR	1	NR	NR	NR
Exposure Type						
Eye Area	NR	NR	6	0.01-0.2	NR	NR
Incidental Ingestion	NR	NR	NR	NR	NR	NR
Incidental Inhalation-Spray	2 ^a	0.01 ^a	2; 30 ^a ; 24 ^b	aerosol: 0.5; pump: 0.2; 0.001-1 ^a	4	NR
Incidental Inhalation-Powder	NR	NR	24 ^b	0.005-3.2 ^c	NR	NR
Dermal Contact	4	NR	91	0.0001-6.5	NR	0.1
Deodorant (underarm)	NR	NR	NR	NR	NR	NR
Hair - Non-Coloring	7	0.01	65	0.05-1	24	NR
Hair-Coloring	NR	NR	6	0.25	NR	NR
Nail	NR	NR	NR	NR	NR	NR
Mucous Membrane	2	NR	21	0.05-1	NR	NR
Baby Products	NR	NR	2	NR	NR	NR
	PEG-6 Caprylic/Capric Glycerides		PEG-7 Caprylic/Capric Glycerides		PEG-8 Caprylic/Capric Glycerides	
Totals*	548	0.000002-5	2	2	22	NR
Duration of Use						
Leave-On	96	0.1-3.5	NR	2	16	NR
Rinse Off	434	0.000002-5	2	NR	4	NR
Diluted for (Bath) Use	18	0.38-5	NR	NR	2	NR
Exposure Type						
Eye Area	17	0.35	NR	NR	NR	NR
Incidental Ingestion	NR	NR	NR	NR	NR	NR
Incidental Inhalation-Spray	3; 17 ^a ; 10 ^b	0.1-3.5 ^a	NR	NR	3 ^a ; 11 ^b	NR
Incidental Inhalation-Powder	10 ^b	0.13-0.75 ^c	NR	2 ^c	11 ^b	NR
Dermal Contact	476	0.0002-5	2	2	21	NR
Deodorant (underarm)	NR	NR	NR	NR	NR	NR
Hair - Non-Coloring	71	0.000002-3.5	NR	NR	1	NR
Hair-Coloring	NR	1.9	NR	NR	NR	NR
Nail	NR	NR	NR	NR	NR	NR
Mucous Membrane	340	0.0002-5	NR	NR	4	NR
Baby Products	2	NR	NR	NR	NR	NR
	PEG-75 Cocoa Butter Glycerides (as (PEG-75 Cocoa Butter in VCRP))		PEG-3 Glyceryl Cocoate		PEG-7 Glyceryl Cocoate	
Totals*	20	NR	2	NR	858	0.01-11.3
Duration of Use						
Leave-On	15	NR	2	NR	124	0.02-11.3
Rinse Off	5	NR	NR	NR	713	0.01-10
Diluted for (Bath) Use	NR	NR	NR	NR	21	NR
Exposure Type						
Eye Area	NR	NR	NR	NR	12	0.02
Incidental Ingestion	2	NR	NR	NR	NR	NR
Incidental Inhalation-Spray	1; 10 ^a	NR	2 ^b	NR	1; 25 ^a ; 50 ^b	6; 0.18-11.3 ^a
Incidental Inhalation-Powder	NR	NR	2 ^b	NR	50 ^b ; 1 ^c	0.35-3.5 ^c
Dermal Contact	13	NR	2	NR	700	0.01-10
Deodorant (underarm)	NR	NR	NR	NR	14 ^a	not spray: 3; pump spray: 2
Hair - Non-Coloring	5	NR	NR	NR	146	0.18-11.3
Hair-Coloring	NR	NR	NR	NR	11	1.8-4.6
Nail	NR	NR	NR	NR	1	0.5
Mucous Membrane	3	NR	NR	NR	453	0.5-6.9
Baby Products	NR	NR	NR	NR	10	NR

Table 5. Frequency and concentration of use according to duration and type of exposure

	# of Uses ²⁸	Max. Conc. of Use (%) ^{29,31}	# of Uses ²⁸	Max. Conc. of Use (%) ^{29,31}	# of Uses ²⁸	Max. Conc. of Use (%) ^{29,31}
	PEG-30 Glyceryl Cocoate		PEG-40 Glyceryl Cocoate		PEG-80 Glyceryl Cocoate	
Totals*	44	0.09-3	23	0.88-4.8	11	0.1-1
Duration of Use						
<i>Leave-On</i>	2	0.1-2.4	1	NR	0	NR
<i>Rinse Off</i>	42	0.09-2.4	21	0.88-4.8	11	0.1-1
<i>Diluted for (Bath) Use</i>	NR	NR	1	NR	0	NR
Exposure Type						
Eye Area	NR	NR	2	0.88	NR	NR
Incidental Ingestion	NR	NR	NR	NR	NR	NR
Incidental Inhalation-Spray	NR	0.1 ^a	1	NR	NR	NR
Incidental Inhalation-Powder	NR	0.5-2.4 ^c	NR	NR	NR	NR
Dermal Contact	22	0.09-3	20	0.88-4.8	10	0.1-1
Deodorant (underarm)	NR	NR	NR	NR	NR	NR
Hair - Non-Coloring	22	0.1-1	3	NR	1	NR
Hair-Coloring	NR	NR	NR	NR	NR	NR
Nail	NR	NR	NR	NR	NR	NR
Mucous Membrane	11	3	7	NR	2	0.1
Baby Products	1	NR	NR	NR	NR	NR
	PEG-20 Hydrogenated Palm Glycerides		PEG-16 Macadamia Glycerides		PEG-70 Mango Glycerides	
Totals*	NR	0.23-1.2	8	0.05-1	31	0.01-0.1
Duration of Use						
<i>Leave-On</i>	NR	0.23-1.2	3	NR	30	0.05
<i>Rinse Off</i>	NR	NR	5	0.05-1	1	0.01-0.1
<i>Diluted for (Bath) Use</i>	NR	NR	NR	NR	NR	NR
Exposure Type						
Eye Area	NR	0.26-1.2	NR	NR	NR	NR
Incidental Ingestion	NR	NR	NR	NR	NR	NR
Incidental Inhalation-Spray	NR	NR	1 ^a	NR	4; 4 ^a	0.05 ^a
Incidental Inhalation-Powder	NR	NR	NR	NR	NR	NR
Dermal Contact	NR	0.23-1.2	8	0.05-1	1	NR
Deodorant (underarm)	NR	NR	NR	NR	NR	NR
Hair - Non-Coloring	NR	NR	NR	NR	30	0.01-0.1
Hair-Coloring	NR	NR	NR	NR	NR	NR
Nail	NR	NR	NR	NR	NR	NR
Mucous Membrane	NR	NR	4	0.05	1	NR
Baby Products	NR	NR	NR	NR	NR	NR
	PEG-10 Olive Glycerides		PEG-45 Palm Kernel Glycerides		PEG-75 Shea Butter Glycerides (includes PEG-75 Shea Butter in VCRP)	
Totals*	1	0.02-0.5	43	0.5-5	29	0.005-1
Duration of Use						
<i>Leave-On</i>	1	NR	15	0.9-5	9	0.005-1
<i>Rinse Off</i>	NR	NR	19	0.5	19	0.1-1
<i>Diluted for (Bath) Use</i>	NR	NR	9	NR	1	NR
Exposure Type						
Eye Area	NR	NR	NR	NR	NR	NR
Incidental Ingestion	NR	NR	NR	NR	NR	NR
Incidental Inhalation-Spray	1 ^a	0.02 ^a	4; 7 ^b	pump:1.9	1; 2 ^a ; 2 ^b	0.005; pump: 0.15; 0.1 ^b
Incidental Inhalation-Powder	NR	NR	7 ^b	0.92 ^c	2 ^b	0.1 ^b ; 1 ^c
Dermal Contact	NR	0.02	36	0.5-1.9	23	0.005-1
Deodorant (underarm)	NR	0.02	NR	NR	NR	NR
Hair - Non-Coloring	1	0.2-0.5	7	5	5	0.1-0.4
Hair-Coloring	NR	NR	NR	NR	NR	NR
Nail	NR	NR	NR	NR	1	NR
Mucous Membrane	NR	NR	25	NR	16	0.1-0.4
Baby Products	NR	NR	NR	NR	1	NR

Table 5. Frequency and concentration of use according to duration and type of exposure

	# of Uses ²⁸	Max. Conc. of Use (%) ^{29,31}	# of Uses ²⁸	Max. Conc. of Use (%) ^{29,31}	# of Uses ²⁸	Max. Conc. of Use (%) ^{29,31}
	PEG-35 Soy Glycerides		PEG-10 Sunflower Glycerides		PEG-13 Sunflower Glycerides	
Totals*	1	NR	11	0.8-5	1	NR
Duration of Use						
<i>Leave-On</i>	<i>NR</i>	<i>NR</i>	<i>2</i>	<i>5</i>	<i>1</i>	<i>NR</i>
<i>Rinse Off</i>	<i>1</i>	<i>NR</i>	<i>8</i>	<i>0.8-2.5</i>	<i>NR</i>	<i>NR</i>
<i>Diluted for (Bath) Use</i>	<i>NR</i>	<i>NR</i>	<i>1</i>	<i>NR</i>	<i>NR</i>	<i>NR</i>
Exposure Type						
Eye Area	NR	NR	NR	NR	NR	NR
Incidental Ingestion	NR	NR	NR	NR	NR	NR
Incidental Inhalation-Spray	NR	NR	2 ^a	5 ^a	1 ^a	NR
Incidental Inhalation-Powder	NR	NR	NR	NR	NR	NR
Dermal Contact	1	NR	4	2.5	NR	NR
Deodorant (underarm)	NR	NR	NR	NR	NR	NR
Hair - Non-Coloring	NR	NR	7	0.8-5	1	NR
Hair-Coloring	NR	NR	NR	NR	NR	NR
Nail	NR	NR	NR	NR	NR	NR
Mucous Membrane	NR	NR	2	NR	NR	NR
Baby Products	NR	NR	NR	NR	NR	NR

* Because each ingredient may be used in cosmetics with multiple exposure types, the sum of all exposure types may not equal the sum of total uses

^a Includes products that can be sprays, but it is not known whether the reported uses are sprays

^b Not specified whether this product is a spray or a powder or neither, but it is possible it may be a spray or a powder, so this information is captured for both categories of incidental inhalation

^c Includes products that can be powders, but it is not known whether the reported uses are powders

NR – none reported

Table 6. Ingredients Not Reported to be Used

PEG-6 Almond Glycerides	PEG-42 Mushroom Glycerides
PEG-35 Almond Glycerides	PEG-2 Olive Glycerides
PEG-11 Avocado Glycerides	PEG-6 Olive Glycerides
PEG-14 Avocado Glycerides	PEG-7 Olive Glycerides
PEG-11 Babassu Glycerides	PEG-40 Olive Glycerides
PEG-42 Babassu Glycerides	PEG-18 Palm Glycerides
PEG-4 Caprylic/Capric Glycerides	PEG-12 Palm Kernel Glycerides
PEG-11 Cocoa Butter Glycerides	PEG-60 Passiflora Edulis Seed Glycerides
PEG-7 Cocoglycerides	PEG-60 Passiflora Incarnata Seed Glycerides
PEG-9 Cocoglycerides	PEG-45 Safflower Glycerides
PEG-20 Corn Glycerides	PEG-60 Shea Butter Glycerides
PEG-60 Corn Glycerides	PEG-75 Shorea Butter Glycerides
PEG-20 Evening Primrose Glycerides	PEG-75 Soy Glycerides
PEG-60 Evening Primrose Glycerides	PEG-2 Sunflower Glycerides
PEG-78 Glyceryl Cocoate	PEG-7 Sunflower Glycerides
PEG-5 Hydrogenated Corn Glycerides	PEG-5 Tsubakiate Glycerides
PEG-8 Hydrogenated Fish Glycerides	PEG-10 Tsubakiate Glycerides
PEG-6 Hydrogenated Palm/Palm Kernel Glyceride	PEG-20 Tsubakiate Glycerides
PEG-13 Mink Glycerides	PEG-60 Tsubakiate Glycerides
PEG-25 Moringa Glycerides	

Table 7. Dermal irritation and sensitization

Test Article	Concentration/Dose	Test Population	Procedure	Results	Reference
ALTERNATIVE STUDIES					
PEG-60 almond glycerides	100%; 100 µl	NHEK	EpiDerm <i>in vitro</i> toxicity test; MTT was used to measure cell viability	ET ₅₀ > 24 h; expected <i>in vivo</i> dermal irritancy is non-irritating	52
PEG-60 almond glycerides	100%; 100 µl	NHEK	EpiDerm <i>in vitro</i> toxicity test	ET ₅₀ = 5.3 h; expected <i>in vivo</i> dermal irritancy is moderate to mild	51
PEG-6 caprylic/capric glycerides	1% and 10%; 50 µl	skin ² (a 3-D human skin tissue)	Model ZK1300 <i>in vitro</i> toxicity test; MTT was used to measure cell viability	mild to moderate dermal irritation potential	49
PEG-8 caprylic/capric glycerides	35% in an ME also containing 10% isostearyl isostearate and 35% polyglyceryl-6 isostearate	male Wistar rat skin	1 ml of the ME or water was applied to the skin sample for 20 h in a Franz-type diffusion cell	low potential for skin irritancy the ME did not perturb the skin barrier	25
PEG-7 glyceryl cocoate	100%; 100 µl	NHEK	EpiDerm <i>in vitro</i> toxicity test	ET ₅₀ > 24 h; expected <i>in vivo</i> dermal irritancy is non-irritating	53
PEG-7 glyceryl cocoate	1% and 10%; 50 µl	skin ²	Model ZK1300 <i>in vitro</i> toxicity test	mild dermal irritation potential	50
PEG-13 sunflower glycerides	100%; 100 µl	NHEK	EpiDerm <i>in vitro</i> toxicity test	ET ₅₀ > 24 h; expected <i>in vivo</i> dermal irritancy is non-irritating	54
PEG-13 sunflower glycerides	100%	skin ²	Model ZK1350 MTT assay for corrosivity	expected to be non-corrosive	55
NON-HUMAN					
PEG-20 almond glycerides	neat; 0.5 ml	6 female NZW rabbits	24-h semi-occlusive patches (2.5 cm ²) applied to an approximately 6 sq. in. area of intact and abraded skin that was clipped free of hair test sites were scored 1 and 48 h after patch removal	PII = 4.5/8; marked irritant effect, but not a primary irritant (which is PII≥5) 24 h: moderate erythema at both sites of 3 rabbits, and well-defined erythema at both sites of the other 3 rabbits; slight edema at 1 abraded and 2 intact sites, moderate edema at 3 abraded and 4 intact sites, and severe edema at 2 abraded sites 72 h: some decline in edema, but not in erythema	61
PEG-60 almond glycerides	neat; 0.5 ml	6 female NZW rabbits	24-h semi-occlusive patches (2.5 cm ²) applied to an approximately 6 sq. in. area of intact and abraded skin that was clipped free of hair test sites were scored 1 and 48 h after patch removal	PII = 2.6/8; some irritant effect, that was reversible 24 h: defined erythema at all treated sites, except 1 intact site had moderate erythema; moderate edema at both sites of 2 rabbits, and slight edema in the other other animals 72 h: reactions declined; very slight erythema at both sites of 4 rabbits; very slight edema at the abraded site of 1 rabbit; desquamation or scaling at the abraded sites of 2 rabbits and the intact site of 1 rabbit	60
PEG-6 caprylic/capric glycerides	4% in formulation, 0.5 ml	3 NZW rabbits/sex	4-h occlusive patch applied to shaved intact skin	not corrosive; mildly irritating	56

Table 7. Dermal irritation and sensitization

Test Article	Concentration/Dose	Test Population	Procedure	Results	Reference
PEG-6 caprylic/capric glycerides	undiluted, 0.5 g	3 rabbits	Draize test; 4-h semi-occlusive patches were applied to intact and abraded skin the test sites were scored at 24 and 72 h	not an irritant	42
PEG-6 caprylic/capric glycerides	undiluted at induction 50% aq. at challenge	not specified	Bühler test, according to OECD Guideline 406	not an irritant or a sensitizer	42
PEG-8 caprylic/capric glycerides	ME containing 1.3% diclofenac epolamine w/w in 30% propylene glycol monocaprylate, 50% PEG-8 caprylic/capric glycerides/ethoxydiglycol (1:2 w/w), and 20% water	3 albino Wistar rats	100 µl was applied for 24 h to the left ear of each rat	not an irritant no erythema	62
PEG-7 glyceryl cocoate	5, 10, 20, and 50% in sterile water, and undiluted, 0.2 ml	5 female Dunkin-Hartley guinea pigs	topical application tolerance test using 24-h occlusive patches; each animal was patched with 2 concentrations	no irritation at any concentration	57
PEG-7 glyceryl cocoate	50% in Vaseline	4 male and female mutant hairless mice	fixed patches applied with porous leucoplastic to shaved skin for 24 h	no reaction	2
PEG-7 glyceryl cocoate	50% in Vaseline	4 male NZW rabbits	24-h patch test	slight reaction; reaction score of 3/4	2
PEG-7 glyceryl cocoate	50% in Vaseline	4 male Pirbright White guinea pigs	fixed patches applied with porous leucoplastic to shaved skin for 24 h	no reaction	2
PEG-7 glyceryl cocoate	not specified	3 albino rabbits	Draize primary irritation test; patches were applied to intact and abraded skin	mild dermal irritant; PII = 1.66 moderate erythema and mild edema were reported	2
PEG-7 glyceryl cocoate	not specified	2 albino rabbits	Draize primary irritation test sites were rinsed after 1 h and scored at and 48 24 h	skin had reddening directly after rinsing; no irritation observed after 24 h	2
PEG-7 glyceryl cocoate	1% in sterile water at intradermal induction, 0.1 ml 0 and 100% for both topical induction (0.4 ml) and challenge (0.2 ml)	10 female Dunkin-Hartley guinea pigs/group	Magnusson-Kligman maximization study, with FCA at intradermal induction topical induction was performed 1 wk later; 48-h occlusive patches were applied over the intradermal injection sites; the test site was pretreated with 10% SLS in petrolatum the day before patching 2 wks after topical induction, the challenge was performed with 24-h occlusive patches applied to a previously untreated site of test animals and to control animals	not a sensitizer	57
PEG-7 glyceryl cocoate	10% solution	5 male Pirbright White W58 guinea pigs	ten 0.1 ml intracutaneous injections at 2-day intervals; the challenge was performed 14 days after the last induction injection	not a sensitizer sharply demarcated necrotic changes of the skin (about the size of a pea), but no general signs, that subsided during the 14-day interval	2
PEG-10 olive glycerides	3% active matter	rabbits, number not provided	no details provided	not an irritant	16
PEG-10 olive glycerides	50% active matter	guinea pigs, number not provided	sensitization study; no details provided	not an irritant or sensitizer	16

Table 7. Dermal irritation and sensitization

Test Article	Concentration/Dose	Test Population	Procedure	Results	Reference
PEG-12 palm kernel glycerides	neat; 0.5 ml	6 female NZW rabbits	24-h occlusive patches (2.5 cm ²) applied to intact and abraded skin that was clipped free of hair; test sites were scored 1 and 48 h after patch removal	PII = 4.5/8; definite irritant effect 24 h: well-defined erythema at all test sites; moderate edema at both test sites of 3 rabbits, and slight edema at the remaining test sites 48 h: some increase in erythema and some decrease in edema; erythema increased to moderate to severe at both test sites of one animal and the abraded site of another; edema was moderate at both sites of 2 animals, slight at both sites of 3 animals, and very slight at both sites of 1 animal	59
PEG-45 palm kernel glycerides	neat; 0.5 ml	6 female NZW rabbits	24-h occlusive patches (2.5 cm ²) applied to intact and abraded skin that was clipped free of hair; test sites were scored 1 and 48 h after patch removal	PII = 2.6/8; definite irritant effect 24 h: well-defined erythema at all test sites; slight edema at both test sites of 3 rabbits, and very slight edema at the remaining test sites 72 h: reactions declined; well-defined erythema in one animal, very slight erythema at both sites of 3 animals, very slight erythema with desquamation at the abraded site and no erythema at the intact site of 1 animal, and no erythema at either site of 1 animal; slight edema at both test sites of 1 animal (this was an increase), very slight edema at both test sites of 1 animal, very slight edema at the abraded sites and no edema at the intact sites of 3 animals, and no edema at either site of 1 animal	58
HUMAN					
PEG-20 almond glyceride	5.0% aq; 0.4 ml	93 subjects	occlusive HRIPT; nine 24-h patches were applied to a site on the upper arm 3x/wk for 3 wks; the 24-h challenge patch was applied after a 17 day non-treatment period	not a sensitizer	69
PEG-60 almond glycerides	20% aq; 0.2 ml	51 subjects	occlusive HRIPT; 24-h semi-occlusive patches (1 sq. in.) were applied to a site on the upper back 3x/wk for a total of 10 applications; challenge patches were applied to the original and untreated sites after a 14-day non-treatment period, and the sites were scored 24-h and 48-h after application	not an irritant or a sensitizer	64
PEG-6 caprylic/capric glycerides	2% aq	50 subjects	48-h patch was applied to the back of each subject; it was not specified whether the patch was occlusive; the test site was observed at 48 and 72 h	not an irritant	42
PEG-6 caprylic/capric glycerides	concentration not specified; assumed undiluted	112 subject; 48 had sensitive skin	single 48-h patch; the type of patch was not specified	not irritating	42
PEG-6 caprylic/capric glycerides	4% in formulation	81 subjects	HRIPT; 24-h patch were applied to a site on the arm 3x/wk for 3 wks; the challenge patch was applied after a 10 day non-treatment period	not a sensitizer	63

Table 7. Dermal irritation and sensitization

Test Article	Concentration/Dose	Test Population	Procedure	Results	Reference
PEG-6 caprylic/capric glycerides	10% aq; 0.2 ml	57 subjects	occlusive HRIPT; 24-h semi-occlusive patches (1 in. x 0.75 in.) were applied to a site on the upper back 3x/wk for a total of 10 applications challenge patches were applied to the original and untreated sites after a 14-day non-treatment period, and the sites were scored 24-h and 48-h after application	not an irritant or a sensitizer	66
PEG-6 caprylic/capric glycerides	concentration not specified; assumed undiluted	10 subjects	five 48-h patches were applied at 48-h intervals; a 48-h challenge patch was applied after a 10-day non-treatment period	no reactions were observed	42
PEG-7 glyceryl cocoate	50%	4 subjects	24-h patches (not specified if occlusive or not)	no reactions	2
PEG-7 glyceryl cocoate	undiluted	40 subjects	primary irritation patch test (details not provided)	no signs of irritation	2
PEG-7 glyceryl cocoate	not specified	5 subjects	test plaster applied to forearm for 1 h	no signs of irritation	2
PEG-7 glyceryl cocoate	8% in formulation applied neat; 0.1 g	103 subjects	semi-occlusive HRIPT; nine 24-h patches were applied to a site on the back 3x/wk for 3 wks; the 24-h challenge patch was applied after a 10-15 day non-treatment period	not a sensitizer	70
PEG-7 glyceryl cocoate	8% in formulation applied neat; 0.1 g	103 subjects (same subjects as above)	semi-occlusive HRIPT, followed the same protocol as above	not a sensitizer	71
PEG-7 glyceryl cocoate	11% in formulation; 0.2 g	205 subjects	semi-occlusive HRIPT; nine 24-h patches were applied to a site on the back 3x/wk for 3 wks; the 24-h challenge patch was applied after a 10-15 day non-treatment period	not a sensitizer one subject had definite erythema and edema (++) at the 2 nd induction reading and was removed from the study	72
PEG-7 glyceryl cocoate	neat; 0.2 ml	57 subjects	occlusive HRIPT; 24-h semi-occlusive patches (1 in. x 0.75 in.) were applied to a site on the upper back 3x/wk for a total of 10 applications challenge patches were applied to the original and untreated sites after a 14-day non-treatment period, and the sites were scored 24-h and 48-h after application	not an irritant or a sensitizer	65
PEG-10 olive glycerides	2% active matter	not stated	Duhring-chamber test; no details provided	not irritating to mildly irritating	16
PEG-75 shea butter glycerides	1% in formulation	10 subjects	cutaneous tolerance test; product was applied to the whole face., including the eyes, 2x/day for 5 days	cutaneous irritant potential is “practically nil”	68
PEG-75 shea butter glycerides	1% in formulation; tested neat; 0.2 g	219 subjects	semi-occlusive HRIPT; nine 24-h patches were applied to a site on the back 3x/wk for 3 wks; the 24-h challenge patch was applied after a 2-wk non-treatment period	not a sensitizer 2 subjects had ± reactions during induction (one on D1; one on D7) 5 subjects had ±1 reactions at challenge	67

Abbreviations: ET₅₀ – time at which percent viability would be 50%; FCA – Freund’s complete adjuvant; HRIPT – human repeated insult patch test; ME – microemulsion; MTT – 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazolium bromide; NHEK – normal, human-derived epidermal keratinocytes; NZW – New Zealand White; PII – primary irritation index; SLS – sodium lauryl sulfate

Table 8. Ocular irritation studies

Test Article	Concentration/Dose	Test System	Procedure	Results	Reference
ALTERNATIVE STUDIES					
PEG-7 glyceryl cocoate	11% in formulation; tested at 50%; 0.3 ml	white Leghorn chicken eggs	HET-CAM	practically no irritation potential mean score 1.25/32	⁷³
PEG-7 glyceryl cocoate	10 and 50% in water and undiluted, 0.3 ml	chicken eggs	HET-CAM	slight irritant	⁷⁴
PEG-7 glyceryl cocoate	100-1468 µg/ml	human keratinocytes (HaCaT)	NRU	cytotoxic to the HaCaT cells, resulting in a median MRU-50 value of 427 µg/ml; considered not classified (≥750 is classified as non-irritant, and <110 is classified as a severe irritant)	⁷⁵
PEG-10 sunflower glyceride	neat; 40 µl	DeKalb chicken eggs	CAMVA	non-irritant RC ₅₀ was 80% (>5.0% is a non-irritant)	⁷⁶
NON-HUMAN STUDIES					
PEG-6 caprylic/capric glycerides	4% in formulation – tested neat (2 groups) and at 10% aq.; 0.1 ml	3 NZW rabbits/group	undiluted test article was instilled into the conjunctival sac of the eye and the eye was rinsed in one group and not rinsed in the other; the eyes of the animals dosed with 10% were not rinsed	not an ocular irritant	⁷⁸
PEG-6 caprylic/capric glycerides	1 and 5% aq.; 0.1 ml	4 rabbits/grp	test substance was applied, and the eyes were evaluated for up to 72 h	no irritant effects 1%: no redness, swelling, secretion, or corneal changes 5%: slight conjunctival redness, slight swelling, and secretion in 1 animal; these signs were reversed by 48 h	⁴²
PEG-6 caprylic/capric glycerides	undiluted; 0.1 ml	2 rabbit	Draize test; the eyes were observed for up to 8 days	strong irritation, including redness, swelling, and secretion, was observed no corneal changes were reported swelling decreased over the 8-day period, but redness was not completely reversible	⁴²
PEG-8 caprylic/capric glycerides	0.5-5.0% in phosphate buffer (pH 7.4); 0.1 ml/dose	6 NZW rabbits	instilled into the conjunctival sac of the left eye 4x/day for 7 days	0.5-3% was not irritating 5% was a slight irritant	⁷⁷
PEG-7 glyceryl cocoate	10% aq.	2 rabbits	instilled into the conjunctival sac	not irritating	²
HUMAN STUDIES					
PEG-75 shea butter glycerides	1% in formulation	10 subjects	ocular tolerance test; product was applied to the whole face., including the eyes, 2x/day for 5 days	no ocular irritation; ocular irritation rate of 0.15%	⁶⁸

Abbreviations: CAMVA – chorioallantoic membrane vascular assay (CAMVA); HET-CAM – hen's egg test utilizing the chorioallantoic membrane; NRU – neutral red uptake; NZW – New Zealand White; OECD – Organisation for Economic Co-operation and Development; RC₅₀ – theoretical concentration producing a positive reaction in 50% of treated eggs

REFERENCES

1. Nikitakis J and Breslawec HP. International Cosmetic Ingredient Dictionary and Handbook. 15 ed. Washington, DC: Personal Care Products Council, 2014.
2. Andersen FA (ed). Final report on the safety assessment of PEG-7, -30, -40, -78, and -80 glyceryl cocoate. *Int J Toxicol*. 1999;18(Suppl 1):33-42.
3. Burnett CL, Heldreth, BA, Bergfeld WF, Belsito DV, Hill RA, Klaassen CD, Liebler DC, Marks JG, Shank RC, Slaga TJ, Snyder PW, and Andersen FA. Amended safety assessment of PEGylated oils as used in cosmetics. 2013. Available at <http://www.cir-safety.org/ingredients>.
4. Bergfeld WF, Belsito DV, Hill RA, Klaassen CD, Liebler DC, Marks Jr JG, Shank RC, Slaga TJ, Snyder PW, and Andersen FA. Amended Safety Assessment of Triethylene Glycol and Polyethylene Glycols (PEGs)-4, -6, -7, -8, -9, -10, -12, -14, -16, -18, -20, -32, -33, -40, -45, -55, -60, -75, -80, -90, -100, -135, -150, -180, -200, -220, -240, -350, -400, -450, -500, -800, -2M, -5M, -7M, -9M, -14M, -20M, -23M, -25M, -45M, -65M, -90M, -115M, -160M and -180M and any PEGs \geq 4 as used in Cosmetics. Washington, DC, Cosmetic Ingredient Review. 2010.
5. Burnett CL, Fiume MM, Bergfeld WF, Belsito DV, Hill RA, Klaassen CD, Liebler DC, Marks JG, Shank RC, Slaga TJ, Snyder PW, and Andersen FA. Final report of Plant-Derived Fatty Acid Oils as Used in Cosmetics. 2011. Available at <http://www.cir-safety.org/ingredients>.
6. Elder RL (ed). Final report on the safety assessment for caprylic/capric triglyceride. *Journal of Environmental Pathology and Toxicology*. 1980;4(4):105-120.
7. Burnett CL, Bergfeld WF, Belsito DV, Klaassen CD, Marks JG Jr, Shank RC, Slaga TJ, Snyder PW, and Andersen FA. Final report on the safety assesment of *Cocos nucifera* (coconut) oil and related ingredients. *Int J Toxicol*. 2011;30(Suppl 1):5S-16S.
8. Andersen FA, Bergfeld WF, Belsito DV, Klaassen CD, Marks JG Jr, Shank RC, Slaga TJ, and Snyder PW. Final report on the safety assessment of cosmetic ingredients derived from *Zea mays* (corn). *Int J Toxicol*. 2011;30(Suppl 1):17S-39S.
9. Andersen FA (ed). Final report on the safety assessment of *Elaeis guineensis* (palm) oil, *Elaeis guineensis* (palm) kernel oil, hydrogenated palm oil and hydrogenated palm kernel oil. *Int J Toxicol*. 2000;19(Suppl 2):7-28.
10. Andersen FA (ed). Final Amended Report on the Safety Assessment of Mink Oil. *Int J Toxicol*. 2005;24(Suppl 3):57-64.
11. Elder RL (ed). Final report on the safety assessment of safflower oil. *J Am Coll Toxicol*. 1985;4(5):171-197.
12. O'Lenick AJ Jr and Parkinson JK. Group selectivity of ethoxylation of hydroxy acids. *J Soc Cosmet Chem*. 1993;44(6):319-328.
13. Nasioudis A, van Velde JW, Heeren RMA, and van den Brink OF. Detailed molecular characterization of castor oil ethoxylates by liquiid chromatography multistage mass spectrometry. *J Chromatogr A*. 2011;1218(40):7166-7172.
14. Zschimmer & Schwarz GmbH & Co KG. 2012. Raw material information PEG-11 Avocado Glycerides. Unpublished data submitted by Personal Care Products Council.
15. Zschimmer & Schwarz GmbH & Co KG. 2013. Raw material information PEG-9 Cocoglycerides. Unpublished data submitted by Personal Care Products Council.
16. Zschimmer & Schwarz GmbH & Co KG. 2013. Raw material information PEG-10 Olive Glycerides. Unpublished data submitted by Personal Care Products Council.
17. Evonik Industries. 2013. Tegosoftware® GMC 6 (PEG-6 Caprylic/Capric Glycerides): Product data record. Unpublished data submitted by Personal Care Products Council.
18. Cremer Oleo GmbH and Co KG. 2014. Softigen 767 (PEG-6 Caprylic/Capric Glycerides) statement about method of manufacture. Unpublished data submitted by Personal Care Products Council.
19. Council of Experts. Caprylocaproyl polyoxyglycerides. http://www.pharmacopeia.cn/v29240/usp29nf24s0_m12363.html. U.S.Pharmacopeia. Date Accessed 5-27-2014.

20. Council of Europe. Caprylocaproyl macrogolglycerides. http://lib.njutc.edu.cn/yaodian/ep/EP5.0/16_monographs/monographs_a-c/Caprylocaproyl%20macrogolglycerides.pdf. European Pharmacopoeia. Date Accessed 5-27-2014.
21. Kociba RJ, McCollister SB, Park C, Torkelson TR, and Gehring PJ. 1,4-Dioxane. I. Results of a 2-year ingestion study in rats. *Toxicol Appl Pharmacol.* 1974;30:275-286.
22. Elder RL (ed). Final Report on the Safety Assessment of PEG-2, -6, -8, -12, -20, -32, -40, -50, -100, and -150 Stearates. *JACT.* 1983;2(7):17-34.
23. Food and Drug Administration (FDA). 1,4-Dioxane. <http://www.fda.gov/cosmetics/productandingredientsafety/potentialcontaminants/ucm101566.htm>. 10903 New Hampshire Ave, Silver Spring, MD 20993. Date Accessed 9-17-2012.
24. Evonik Industries. 2011. Tegosoft® GMC 6 (PEG-6 Caprylic/Capric Glycerides): Product specification. Unpublished data submitted by Personal Care Products Council.
25. Kreilgaard M, Pdersen EJ, and Jaroszewski JW. NMR characterisation and transdermal drug delivery potential of microemulsion systems. *J Control Release.* 2000;69(3):421-433.
26. Gattefossé. Technical data sheet: Labrasol. http://www.gattefosse.com/media/document/tds_labrasol.PDF. Date Accessed 5-14-2014.
27. Abitec. Technical data sheet: Acconon® MC8-2, EP/NF (PEG-8 caprylic/capric glycerides). http://www.abiteccorp.com/wp-content/files_mf/1385392969AccononMC82EP.NFTDSI4.pdf. Date Accessed 5-15-2014.
28. Food and Drug Administration (FDA). Frequency of use of cosmetic ingredients. *FDA Database.* 2014.
29. Personal Care Products Council. 8-21-2014. Updated Concentration of Use by FDA Product Category: PEG Glyceryl Cocoate Ingredients. Unpublished data submitted by Personal Care Products Council.
30. Personal Care Products Council. 7-15-2014. Concentration of use by FDA product category: PEG-6 Hydrogenated Palm/Palm Kernel Glyceride. Unpublished data submitted by Personal Care Products Council.
31. Personal Care Products Council. 5-12-2014. Updated Concentration of Use Information: PEGylated Alkyl Glyceride Ingredients. Unpublished data submitted by Personal Care Products Council.
32. Johnsen MA. The influence of particle size. *Spray Technol Marketing.* 2004;14(11):24-27.
33. Rothe H. Special Aspects of Cosmetic Spray Evaluation. 9-26-2011. Unpublished data presented at the 26 September CIR Expert Panel meeting. Washington, D.C.
34. Rothe H, Fautz R, Gerber E, Neumann L, Rettinger K, Schuh W, and Gronewold C. Special aspects of cosmetic spray safety evaluations: Principles on inhalation risk assessment. *Toxicol Lett.* 2011;205(2):97-104.
35. Bremmer HJ, Prud'homme de Lodder LCH, and Engelen JGM. Cosmetics Fact Sheet: To assess the risks for the consumer; Updated version for ConsExpo 4. 2006. Report No. RIVM 320104001/2006. pp. 1-77.
36. European Commission. CosIng database; following Cosmetic Regulation No. 1223/2009. <http://ec.europa.eu/consumers/cosmetics/cosing/>. Date Accessed 4-28-2014.
37. Djekic, Ljiljana and Primorac, Marija. The influence of cosurfactants and oils on the formation of pharmaceutical microemulsions based on PEG-8 caprylic/capric glycerides. *International Journal of Pharmaceutics.* 2008;352(1-2):231-239.
38. Food and Drug Administration (FDA). Inactive Ingredient Search for Approved Drug Products. <http://www.accessdata.fda.gov/scripts/cder/iig/getiigWEB.cfm>. Date Accessed 10-21-2014.
39. Gannu R, Vishnu YV, Kishan V, and Rao YM. In vitro permeation of carvedilol through porcine skin: effect of vehicles and penetration enhancers. *PDA Journal of Pharmaceutical Science and Technology.* 2008;62(4):256-263.
40. Liu H, Li S, Wang Y, Yao H, and Zhang Y. Effect of vehicles and enhancers on the topical delivery of cyclosporin A. *Int J Pharm.* 3-27-2006;311(1-2):182-186.

41. Shah PP, Desai PR, Patlolla R, Klevans L, and Singh M. Effect of combination of hydrophilic and lipophilic permeation enhancers on the skin permeation of kahalalide F. *J Pharm Pharmacol.* 2-12-2014.
42. Cremer Oleo GmbH and Co KG. 2014. Toxicological and ecotoxicological profile of Softigen 767 (PEG-6 Caprylic/Capric Glycerides). Unpublished data submitted by Personal Care Products Council.
43. Prasad YVR, Eaimtrakarn S, Ishida M, Kusawake Y, Tawa R, Yoshikawa Y, Shibata N, and Takada K. Evaluation of oral formulations of gentamicin containing labrasol in beagle dogs. *International Journal of Pharmaceutics.* 2003;268(1-2):13-21.
44. Delongea JL, de Conchard GV, Beamonte A, Bertheux H, Spire C, Maisonneuve C, Becourt-Lhote N, Goldfain-Blanc F, and Claude N. Assessment of Labrasol/Labrafil/Transcutol (4/4/2, v/v/v) as a non-clinical vehicle for poorly water-soluble compounds after 4-week oral toxicity study in Wistar rats. *Regul Toxicol Pharmacol.* 2010;57(2-3):284-290.
45. Hazleton Laboratories America Inc. 1989. Summary of 28-day subchronic oral toxicity study in rats with a material containing 0.8% PEG-7 Glyceryl Cocoate. Unpublished data submitted by Personal Care Products Council.
46. Personal Care Products Council. 9-19-2014. Summary of a Developmental Toxicity Study on PEG-8 Caprylic/Capric Glycerides. Unpublished data submitted by Personal Care Products Council. 1 pages.
47. NAmSA. 1997. Genotoxicity: *Salmonella typhimurium* reverse mutation assay of PEG-6 caprylic/capric glycerides. Unpublished data submitted by the Personal Care Products Council.
48. NAmSA. 1996. Ames *Salmonella*/mammalian microsome mutagenicity assay of PEG-7 glyceryl cocoate. Unpublished data submitted by the Personal Care Products Council.
49. Consumer Product Testing Co. 1995. The Advanced TissueTM Sciences skin^{2TM} Model ZK1300 *in vitro* toxicity testing system with PEG-6 caprylic/capric glycerides. (Experiment Reference No. V95-0033-2). Unpublished data submitted by Personal Care Products Council.
50. Consumer Product Testing Co. 1995. The Advanced TissueTM Sciences skin^{2TM} Model ZK1300 *in vitro* toxicity testing system with PEG-7 glyceryl cocoate. (Experiment reference no. V95-0033-3.). Unpublished data submitted by the Personal Care Products Council.
51. Consumer Product Testing Co. 1999. The MatTek Corporation EpiDermTM skin model *in vitro* toxicity testing system with PEG-60 almond glycerides. (Experiment Reference No. V99-0122-13). Unpublished data submitted by Personal Care Products Council.
52. Consumer Product Testing Co. 1999. The MatTek Corporation EpiDermTM skin model *in vitro* toxicity testing system PEG-60 almond glycerides. (Experiment Reference No. V99-0122-10). Unpublished data submitted by Personal Care Products Council.
53. Consumer Product Testing Co. 1999. The MakTek Corporation EpiDermTM skin model *in vitro* toxicity testing system with PEG-7 glyceryl cocoate. (Experiment reference no. V99-0122-8.). Unpublished data submitted by the Personal Care Products Council.
54. Consumer Product Testing Co. 2004. The MatTek Corporation EpiDermTM skin model *in vitro* toxicity testing system with PEG-13 sunflower glycerides. (Experiment Reference No. V04-0101-3). Unpublished data submitted by Personal Care Products Council.
55. Thomas J. Stephens & Associates Inc. 1994. Advanced Tissue Sciences' Skin^{2TM} Model 1350 MTT assay for corrosivity testing (protocol #085) with PEG-13 sunflower glycerides (test material 0728-94L). Unpublished data submitted by Personal Care Products Council; File provided has legibility issues - not scanned again.
56. Raltech Scientific Services. 1981. Skin corrosivity (DOT procedure) of a perfume containing 4% PEG-6 caprylic/capric glycerides (P0956). Unpublished data submitted by Personal Care Products Council.
57. Research Toxicology Centre S.p.A. 1995. Delayed dermal sensitization study in the guinea pig (PEG-7 Glyceryl Cocoate). Unpublished data submitted by Personal Care Products Council.
58. Toxicol Laboratories Limited. 1987. Primary skin irritation study PEG-45 Palm Kernel Glycerides (Study Ref. No. 232/8612). Unpublished data submitted by Personal Care Products Council.
59. Toxicol Laboratories Limited. 1987. Primary skin irritation study PEG-12 Palm Kernel Glycerides (Study Ref. No. 229/8612). Unpublished data submitted by Personal Care Products Council.

60. Toxicol Laboratories Limited. 1989. Primary skin irritation study PEG-60 Almond Glycerides (Study Ref. No. A/S/18780). Unpublished data submitted by Personal Care Products Council.
61. Toxicol Laboratories Limited. 1989. Primary skin irritation study PEG-20 Almond Glycerides (Study Ref. No. A/S/18783). Unpublished data submitted by Personal Care Products Council.
62. Fouad SA, Basalious EB, El-Nabarawi MA, and Tayel SA. Microemulsion and poloxamer microemulsion-based gel for sustained transdermal delivery of diclofenac epolamine using in-skin drug depot: In vitro/in vivo evaluation. *International Journal of Pharmaceutics (Amsterdam, Netherlands)*. 2013;453(2):569-578.
63. Anonymous. 1981. Human sensitization test of a perfume containing 4% PEG-6 caprylic/capric glycerides. Unpublished data submitted by Personal Care Products Council.
64. Consumer Product Testing Co. 1996. Repeated insult patch test of PEG-60 almond glycerides. (Experiment Reference No. C95-0520). Unpublished data submitted by Personal Care Products Council.
65. Consumer Product Testing Co. 1996. Repeated insult patch test of PEG-7 of glyceryl cocoate. (Experiment reference no. C96-0088.). Unpublished data submitted by the Personal Care Products Council.
66. Consumer Product Testing Co. 1997. Repeated insult patch test of PEG-6 caprylic/capric glycerides. (Experiment Reference Number C97-0170). Unpublished data submitted by Personal Care Products Council.
67. Harrison Research Laboratories Inc. 2008. Repeated insult patch test of a hair setting lotion containing 1% PEG-75 Shea Butter Glycerides. Unpublished data submitted by Personal Care Products Council.
68. Peritesco. 2004. Ocular and cutaneous tolerance study of self foaming cleansers containing 1% PEG-75 Shea Butter Glycerides. Unpublished data submitted by Personal Care Products Council.
69. Pharmaco UK Ltd. 1994. Human repeat insult patch test (RE-0647.01) of PEG-20 Almond Glycerides, 100% neat tested at 5%. Unpublished data submitted by Personal Care Products Council.
70. TKL Research Inc. 2004. Repeated insult patch test of a styling pomade (BCS578-094) containing 8% PEG-7 Glyceryl Cocoate. Unpublished data submitted by Personal Care Products Council.
71. TKL Research Inc. 2004. Repeated insult patch test of a styling pomade (BCS578-098) containing 8% PEG-7 Glyceryl Cocoate. Unpublished data submitted by Personal Care Products Council.
72. TKL Research Inc. 2013. Repeated insult patch test of a hair styling product containing 11% PEG-7 Glyceryl Cocoate. Unpublished data submitted by Personal Care Products Council.
73. Consumer Product Testing Co. 2013. The hen's egg test- utilizing the chorioallantoic membrane (HET-CAM) of a hair styling product containing 11% PEG-7 glyceryl cocoate. Unpublished data submitted by Personal Care Products Council.
74. Anonymous. 2005. Assessment of the eye irritation potential of PEG-7 Glyceryl Cocoate in the hen's egg test on the chorioallantoic membrane (HET-CAM). Unpublished data submitted by Personal Care Products Council.
75. Anonymous. 2005. Assessment of the eye irritation potential of PEG-7 Glyceryl Cocoate by cytotoxicity measurement in the neutral red uptake assay (NRU) on human keratinocytes (HaCaT). Unpublished data submitted by Personal Care Products Council.
76. Leberco-Celsis Testing. 1997. Ocular irritation screen of PEG-10 Sunflower Glycerides using the chorioallantoic membrane vascular assay (CAMVA). Unpublished data submitted by Personal Care Products Council.
77. Liu Z, Zhang X, Li J, Liu R, Shu L, and Jin J. Effects of Labrasol on the corneal drug delivery of baicalin. *Drug Delivery*. 2009;16(7):399-404.
78. Springborn Institute for Bioresearch, Inc. 1981. Rabbit eye irritation (perfum containing 4% PEG-6 caprylic/capric glycerides).
79. Consumer Product Testing Co. 2005. Safety in use of a female hygiene product containing 0.1% PEG-75 shea butter glycerides. Unpublished data submitted by Personal Care Products Council.
80. Abitec. Acconon® CC-6. http://www.abiteccorp.com/wp-content/files_mf/1310570451AccononCC6TDSI7.pdf. Date Accessed 5-15-2014.

81. SpecialChem. Glycerox PK-70. <http://www.specialchem4cosmetics.com/tds/glycerox-pk-70/croda/15629/index.aspx>. Date Accessed 5-28-2014.
82. SpecialChem. Acconon[®] S-35. <http://www.specialchem4cosmetics.com/tds/acconon-s-35/abitec-corporation/10822/index.aspx>. Date Accessed 5-28-2014.
83. SpecialChem. Oxypon 365. <http://www.specialchem4cosmetics.com/tds/oxypon-365/zschimmer-schwarz-italiana/12620/index.aspx>. Date Accessed 5-28-2014.
84. Zschimmer & Schwarz GmbH & Co KG. Oxypon 365. http://www.zschimmer-schwarz.com/OXYPON_365/simon/zschimmer-schwarz/media/site/downloads/merkblatt/1_S_S_ENG_1108_10_2_140.pdf. Date Accessed 5-28-2014.
85. Anonymous. 2013. Specification/certificate of analysis PEG-11 Avocado Glycerides. Unpublished data submitted by Personal Care Products Council.
86. SpecialChem. Crovol A-40. <http://www.specialchem4cosmetics.com/tds/crovol-ba70g/croda/12994/index.aspx>. Date Accessed 5-28-2014.
87. SpecialChem. Crovol BA70G. <http://www.specialchem4cosmetics.com/tds/crovol-ba70g/croda/12994/index.aspx>. Date Accessed 5-28-2014.
88. Lubrizol Advanced Materials, Inc. Product specifications: Chemonic[™] CCG-6 Surfactant (PEG-6 caprylic/capric glycerides). <http://www.lubrizol.com/personal-care/documents/specifications/chemonic%E2%84%A2-ccg-6-surfactant.pdf>. Date Accessed 5-15-2014.
89. Evonik Industries. Tegoso[®] GC; Tegoso[®] GMC 6. http://www.finecon.sk/admin/pdf/DS_TEGOSOFT_GC_6_e.pdf. Date Accessed 5-28-2014.
90. Zschimmer & Schwarz GmbH & Co KG. Oxypon 401. http://www.zschimmer-schwarz.com/OXYPON_401/simon/zschimmer-schwarz/media/site/downloads/merkblatt/1_S_S_ENG_2815_20_2_140.pdf. Date Accessed 5-28-2014.
91. Croda. Product summary, Crovol[™] M70. <http://www.crodacropcare.com/home.aspx?view=dtl&d=content&s=143&r=256&p=1932&productName=&productfunction=&subfunction=&productgroup=&prodID=182>. Date Accessed 5-28-2014.
92. SpecialChem. Crovol EP70. <http://www.specialchem4cosmetics.com/tds/crovol-ep70/croda/12995/index.aspx>. Date Accessed 5-28-2014.
93. Floratech. Product specifications: Florasolvs[®] PEG-16 macadamia. <file:///C:/Users/mmf/Downloads/D--Hosting-8529091-html-fileMgr-upload-files-SPEC%20SHEETS-Numbers-046.pdf>. Date Accessed 5-28-2014.
94. Floratech. Product information: Florasolvs PEG-16 macadamia. <http://www.floratech.com/mpl/viewdoc.asp?aID=94&searchQ=>. Date Accessed 5-28-2014.
95. AarhusKarlshamn (AAK). Lipex[®] 102 E75; Lipex[®] 203 E70. http://www.aak.se/Global/Products/Beauty%20and%20personal%20care/Surfactants/aak-lfc_lipex_102-203_E75-E70_0808.pdf. Date Accessed 5-28-2014.
96. UL. Lipex[®] 203 E-70. <http://www.innovadex.com/PersonalCare/Detail/1524/130067/Lipex-203-E-70>. Date Accessed 5-28-2014.
97. Zschimmer & Schwarz GmbH & Co KG. Oxypon 288. http://www.zschimmer-schwarz.com/en/OXYPON_288/simon/zschimmer-schwarz/media/site/downloads/merkblatt/1_S_S_ENG_4632_20_2_140.pdf. Date Accessed 5-28-2014.
98. Anonymous. 2013. Specification/certificate of analysis PEG-10 Olive Glycerides. Unpublished data submitted by Personal Care Products Council.
99. SpecialChem. Lexol[®] EO. <http://www.specialchem4cosmetics.com/tds/lexol-eo/inolex/3623/index.aspx>. Date Accessed 5-28-2014.
100. UL. Crovol[™] PK-70. <http://www.innovadex.com/PersonalCare/Detail/134/30933/Crovol-PK-70>. Date Accessed 5-28-2014.
101. UL. Lipex[®] 102 E-75. <http://www.innovadex.com/PersonalCare/Detail/1524/130062/Lipex-102-E-75>. Date Accessed 5-28-2014.

102. Floratech. Product specifications: Florasolvs® PEG-10 sunflower. <file:///C:/Users/mmf/Downloads/D--Hosting-8529091-html-fileMgr-upload-files-SPEC%20SHEETS-Numbers-045.pdf>. Date Accessed 5-28-2014.
103. Floratech. Product information: Florasolvs PEG-16 sunflower. <http://www.floratech.com/mpl/viewdoc.asp?aID=94&searchQ=>. Date Accessed 5-28-2014.
104. Floratech. Florasolvs® PEG-10 Sunflower. https://www.floratech.com/fileMgr/upload/files/PIBS/All/PIB_FlorasolvsPEG-10Sunflower.pdf. Date Accessed 5-28-2014.
105. Natural Sourcing, LLC. Passionfruit Seed Oil. https://www.naturalsourcing.com/media/pr_passionfloweroil.asp. Date Accessed 5-5-2014.
106. Food and Agriculture Organization of the United Nations (FAO). FAO Food and Nutrition Paper, 47/5: Utilization of tropical foods: tropical oil-seeds. http://books.google.com/books?id=OtF6R8vWghEC&pg=PA69&lpg=PA69&dq=shorea+Stenoptera+seed+butter++oil+fatty+acid+composition&source=bl&ots=0eH0KIBzGs&sig=fnTYBqDYE9a_vAWT9oVs_BwjFVg&hl=en&sa=X&ei=uB9nU5HdNqWIyAH9pIGYAg&ved=0CCYQ6AEwAA#v=onepage&q=shorea%20Stenoptera%20seed%20butter%20%20oil%20fatty%20acid%20composition&f=false.

PEG-20 ALMOND GLYCERIDES	05A - Hair Conditioner	4
PEG-20 ALMOND GLYCERIDES	05F - Shampoos (non-coloring)	1
PEG-20 ALMOND GLYCERIDES	05G - Tonics, Dressings, and Other Hair Grooming Aids	2
PEG-20 ALMOND GLYCERIDES	10A - Bath Soaps and Detergents	2
PEG-20 ALMOND GLYCERIDES	12A - Cleansing	1
PEG-20 ALMOND GLYCERIDES	12J - Other Skin Care Preps	1
PEG-60 ALMOND GLYCERIDES	01A - Baby Shampoos	2
PEG-60 ALMOND GLYCERIDES	02D - Other Bath Preparations	1
PEG-60 ALMOND GLYCERIDES	03D - Eye Lotion	4
PEG-60 ALMOND GLYCERIDES	03E - Eye Makeup Remover	1
PEG-60 ALMOND GLYCERIDES	03F - Mascara	1
PEG-60 ALMOND GLYCERIDES	04A - Cologne and Toilet waters	1
PEG-60 ALMOND GLYCERIDES	04B - Perfumes	1
PEG-60 ALMOND GLYCERIDES	05A - Hair Conditioner	1
PEG-60 ALMOND GLYCERIDES	05F - Shampoos (non-coloring)	38
PEG-60 ALMOND GLYCERIDES	05G - Tonics, Dressings, and Other Hair Grooming Aids	15
PEG-60 ALMOND GLYCERIDES	05H - Wave Sets	2
PEG-60 ALMOND GLYCERIDES	05I - Other Hair Preparations	7
PEG-60 ALMOND GLYCERIDES	06D - Hair Shampoos (coloring)	1
PEG-60 ALMOND GLYCERIDES	06F - Hair Lighteners with Color	1
PEG-60 ALMOND GLYCERIDES	06G - Hair Bleaches	4
PEG-60 ALMOND GLYCERIDES	10A - Bath Soaps and Detergents	11
PEG-60 ALMOND GLYCERIDES	10E - Other Personal Cleanliness Products	9
PEG-60 ALMOND GLYCERIDES	11A - Aftershave Lotion	1
PEG-60 ALMOND GLYCERIDES	11G - Other Shaving Preparation Products	1
PEG-60 ALMOND GLYCERIDES	12A - Cleansing	14
PEG-60 ALMOND GLYCERIDES	12C - Face and Neck (exc shave)	22
PEG-60 ALMOND GLYCERIDES	12D - Body and Hand (exc shave)	2
PEG-60 ALMOND GLYCERIDES	12F - Moisturizing	11
PEG-60 ALMOND GLYCERIDES	12G - Night	2
PEG-60 ALMOND GLYCERIDES	12H - Paste Masks (mud packs)	2
PEG-60 ALMOND GLYCERIDES	12I - Skin Fresheners	2
PEG-60 ALMOND GLYCERIDES	12J - Other Skin Care Preps	6
PEG-192 APRICOT KERNEL GLYCERIDES	05B - Hair Spray (aerosol fixatives)	4
PEG-192 APRICOT KERNEL GLYCERIDES	05I - Other Hair Preparations	20
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	01A - Baby Shampoos	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	01C - Other Baby Products	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	02A - Bath Oils, Tablets, and Salts	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	02B - Bubble Baths	6
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	02D - Other Bath Preparations	11
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	03B - Eyeliner	11
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	03E - Eye Makeup Remover	5
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	03F - Mascara	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	04A - Cologne and Toilet waters	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	04B - Perfumes	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	04E - Other Fragrance Preparation	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	05E - Rinses (non-coloring)	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	05F - Shampoos (non-coloring)	64
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	05G - Tonics, Dressings, and Other Hair Grooming Aids	4
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	05I - Other Hair Preparations	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	07I - Other Makeup Preparations	2
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	10A - Bath Soaps and Detergents	191
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	10C - Douches	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	10E - Other Personal Cleanliness Products	128
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	11A - Aftershave Lotion	47
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	11G - Other Shaving Preparation Products	4
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	12A - Cleansing	36

PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	12C - Face and Neck (exc shave)	5
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	12D - Body and Hand (exc shave)	5
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	12F - Moisturizing	8
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	12G - Night	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	12H - Paste Masks (mud packs)	1
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	12I - Skin Fresheners	2
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	12J - Other Skin Care Preps	3
PEG-6 CAPRYLIC/CAPRIC GLYCERIDES	13A - Suntan Gels, Creams, and Liquids	2
PEG-6 CAPRYLIC/CAPRIC TRIGLYCERIDE	10A - Bath Soaps and Detergents	2
PEG-7 CAPRYLIC/CAPRIC GLYCERIDES	12A - Cleansing	2
PEG-8 CAPRYLIC/CAPRIC GLYCERIDES	02A - Bath Oils, Tablets, and Salts	1
PEG-8 CAPRYLIC/CAPRIC GLYCERIDES	02D - Other Bath Preparations	1
PEG-8 CAPRYLIC/CAPRIC GLYCERIDES	05I - Other Hair Preparations	1
PEG-8 CAPRYLIC/CAPRIC GLYCERIDES	10E - Other Personal Cleanliness Products	2
PEG-8 CAPRYLIC/CAPRIC GLYCERIDES	12A - Cleansing	1
PEG-8 CAPRYLIC/CAPRIC GLYCERIDES	12C - Face and Neck (exc shave)	9
PEG-8 CAPRYLIC/CAPRIC GLYCERIDES	12D - Body and Hand (exc shave)	2
PEG-8 CAPRYLIC/CAPRIC GLYCERIDES	12F - Moisturizing	3
PEG-8 CAPRYLIC/CAPRIC GLYCERIDES	12J - Other Skin Care Preps	2
PEG-75 COCOA BUTTER	04B - Perfumes	1
PEG-75 COCOA BUTTER	05A - Hair Conditioner	2
PEG-75 COCOA BUTTER	05F - Shampoos (non-coloring)	3
PEG-75 COCOA BUTTER	07E - Lipstick	2
PEG-75 COCOA BUTTER	07I - Other Makeup Preparations	1
PEG-75 COCOA BUTTER	10E - Other Personal Cleanliness Products	1
PEG-75 COCOA BUTTER	12F - Moisturizing	10
PEG-3 GLYCERYL COCOATE	12D - Body and Hand (exc shave)	2
PEG-7 GLYCERYL COCOATE	01A - Baby Shampoos	5
PEG-7 GLYCERYL COCOATE	01B - Baby Lotions, Oils, Powders, and Creams	1
PEG-7 GLYCERYL COCOATE	01C - Other Baby Products	4
PEG-7 GLYCERYL COCOATE	02A - Bath Oils, Tablets, and Salts	4
PEG-7 GLYCERYL COCOATE	02B - Bubble Baths	9
PEG-7 GLYCERYL COCOATE	02D - Other Bath Preparations	8
PEG-7 GLYCERYL COCOATE	03D - Eye Lotion	1
PEG-7 GLYCERYL COCOATE	03E - Eye Makeup Remover	10
PEG-7 GLYCERYL COCOATE	03G - Other Eye Makeup Preparations	1
PEG-7 GLYCERYL COCOATE	04E - Other Fragrance Preparation	1
PEG-7 GLYCERYL COCOATE	05A - Hair Conditioner	6
PEG-7 GLYCERYL COCOATE	05F - Shampoos (non-coloring)	122
PEG-7 GLYCERYL COCOATE	05G - Tonics, Dressings, and Other Hair Grooming Aids	11
PEG-7 GLYCERYL COCOATE	05I - Other Hair Preparations	2
PEG-7 GLYCERYL COCOATE	06A - Hair Dyes and Colors (all types requiring caution statement	1
PEG-7 GLYCERYL COCOATE	06D - Hair Shampoos (coloring)	9
PEG-7 GLYCERYL COCOATE	06G - Hair Bleaches	1
PEG-7 GLYCERYL COCOATE	07I - Other Makeup Preparations	2
PEG-7 GLYCERYL COCOATE	08C - Nail Creams and Lotions	1
PEG-7 GLYCERYL COCOATE	10A - Bath Soaps and Detergents	342
PEG-7 GLYCERYL COCOATE	10B - Deodorants (underarm)	14
PEG-7 GLYCERYL COCOATE	10C - Douches	1
PEG-7 GLYCERYL COCOATE	10E - Other Personal Cleanliness Products	89
PEG-7 GLYCERYL COCOATE	11A - Aftershave Lotion	9


PEG-7 GLYCERYL COCOATE	11E - Shaving Cream	1
PEG-7 GLYCERYL COCOATE	11G - Other Shaving Preparation Products	6
PEG-7 GLYCERYL COCOATE	12A - Cleansing	114
PEG-7 GLYCERYL COCOATE	12C - Face and Neck (exc shave)	32
PEG-7 GLYCERYL COCOATE	12D - Body and Hand (exc shave)	18
PEG-7 GLYCERYL COCOATE	12F - Moisturizing	11
PEG-7 GLYCERYL COCOATE	12G - Night	2
PEG-7 GLYCERYL COCOATE	12H - Paste Masks (mud packs)	6
PEG-7 GLYCERYL COCOATE	12I - Skin Fresheners	1
PEG-7 GLYCERYL COCOATE	12J - Other Skin Care Preps	13
PEG-30 GLYCERYL COCOATE	01C - Other Baby Products	1
PEG-30 GLYCERYL COCOATE	05F - Shampoos (non-coloring)	22
PEG-30 GLYCERYL COCOATE	07C - Foundations	1
PEG-30 GLYCERYL COCOATE	10A - Bath Soaps and Detergents	6
PEG-30 GLYCERYL COCOATE	10E - Other Personal Cleanliness Products	5
PEG-30 GLYCERYL COCOATE	12A - Cleansing	9
PEG-40 GLYCERYL COCOATE	02D - Other Bath Preparations	1
PEG-40 GLYCERYL COCOATE	03E - Eye Makeup Remover	2
PEG-40 GLYCERYL COCOATE	04E - Other Fragrance Preparation	1
PEG-40 GLYCERYL COCOATE	05F - Shampoos (non-coloring)	3
PEG-40 GLYCERYL COCOATE	10A - Bath Soaps and Detergents	6
PEG-40 GLYCERYL COCOATE	12A - Cleansing	10
PEG-80 GLYCERYL COCOATE	05F - Shampoos (non-coloring)	1
PEG-80 GLYCERYL COCOATE	10A - Bath Soaps and Detergents	2
PEG-80 GLYCERYL COCOATE	10E - Other Personal Cleanliness Products	8
PEG-16 MACADAMIA GLYCERIDES	10A - Bath Soaps and Detergents	3
PEG-16 MACADAMIA GLYCERIDES	10E - Other Personal Cleanliness Products	1
PEG-16 MACADAMIA GLYCERIDES	12A - Cleansing	1
PEG-16 MACADAMIA GLYCERIDES	12F - Moisturizing	1
PEG-16 MACADAMIA GLYCERIDES	12J - Other Skin Care Preps	2
PEG-70 MANGO GLYCERIDES	05B - Hair Spray (aerosol fixatives)	4
PEG-70 MANGO GLYCERIDES	05G - Tonics, Dressings, and Other Hair Grooming Aids	4
PEG-70 MANGO GLYCERIDES	05I - Other Hair Preparations	22
PEG-70 MANGO GLYCERIDES	10E - Other Personal Cleanliness Products	1
PEG-10 OLIVE GLYCERIDES	05G - Tonics, Dressings, and Other Hair Grooming Aids	
PEG-45 PALM KERNEL GLYCERIDES	02A - Bath Oils, Tablets, and Salts	1
PEG-45 PALM KERNEL GLYCERIDES	02D - Other Bath Preparations	8
PEG-45 PALM KERNEL GLYCERIDES	05B - Hair Spray (aerosol fixatives)	4
PEG-45 PALM KERNEL GLYCERIDES	05F - Shampoos (non-coloring)	1
PEG-45 PALM KERNEL GLYCERIDES	05H - Wave Sets	1
PEG-45 PALM KERNEL GLYCERIDES	05I - Other Hair Preparations	1
PEG-45 PALM KERNEL GLYCERIDES	07I - Other Makeup Preparations	1
PEG-45 PALM KERNEL GLYCERIDES	10A - Bath Soaps and Detergents	16
PEG-45 PALM KERNEL GLYCERIDES	12A - Cleansing	1
PEG-45 PALM KERNEL GLYCERIDES	12C - Face and Neck (exc shave)	5
PEG-45 PALM KERNEL GLYCERIDES	12D - Body and Hand (exc shave)	2
PEG-45 PALM KERNEL GLYCERIDES	12J - Other Skin Care Preps	2

PEG-75 SHEA BUTTER GLYCERIDES	02A - Bath Oils, Tablets, and Salts	1
PEG-75 SHEA BUTTER GLYCERIDES	05A - Hair Conditioner	1
PEG-75 SHEA BUTTER GLYCERIDES	05B - Hair Spray (aerosol fixatives)	1
PEG-75 SHEA BUTTER GLYCERIDES	05F - Shampoos (non-coloring)	1
PEG-75 SHEA BUTTER GLYCERIDES	05I - Other Hair Preparations	2
PEG-75 SHEA BUTTER GLYCERIDES	10A - Bath Soaps and Detergents	8
PEG-75 SHEA BUTTER GLYCERIDES	10E - Other Personal Cleanliness Products	6
PEG-75 SHEA BUTTER GLYCERIDES	12A - Cleansing	2
PEG-75 SHEA BUTTER GLYCERIDES	12D - Body and Hand (exc shave)	1
PEG-75 SHEA BUTTER GLYCERIDES	12F - Moisturizing	1
PEG-50 SHEA BUTTER	01C - Other Baby Products	1
PEG-50 SHEA BUTTER	08B - Cuticle Softeners	1
PEG-50 SHEA BUTTER	10E - Other Personal Cleanliness Products	1
PEG-50 SHEA BUTTER	12D - Body and Hand (exc shave)	1
PEG-50 SHEA BUTTER	12F - Moisturizing	1
PEG-35 SOY GLYCERIDES	12A - Cleansing	1
PEG-10 SUNFLOWER GLYCERIDES	02B - Bubble Baths	1
PEG-10 SUNFLOWER GLYCERIDES	05F - Shampoos (non-coloring)	5
PEG-10 SUNFLOWER GLYCERIDES	05G - Tonics, Dressings, and Other Hair Grooming Aids	2
PEG-10 SUNFLOWER GLYCERIDES	10A - Bath Soaps and Detergents	1
PEG-10 SUNFLOWER GLYCERIDES	11G - Other Shaving Preparation Products	1
PEG-10 SUNFLOWER GLYCERIDES	12A - Cleansing	1
PEG-13 SUNFLOWER GLYCERIDES	05G - Tonics, Dressings, and Other Hair Grooming Aids	1



Memorandum

TO: Lillian Gill, D.P.A.
Director - COSMETIC INGREDIENT REVIEW (CIR)


FROM: Beth A. Lange, Ph.D.
Industry Liaison to the CIR Expert Panel 

DATE: September 19, 2014

SUBJECT: Summary of a Developmental Toxicity Study on PEG-8 Caprylic/Capric Glycerides

A supplier has provided the following summary of a developmental toxicity study on PEG-8 Caprylic/Capric Glycerides.

Test material: PEG-8 Caprylic/Capric Glycerides
Study Type: Segment II developmental toxicity (treatment from gestation days 6 through 17)
Route: Oral (gavage)
Species: rat
Doses: vehicle/control: purified water; 0, 1000 , 2000 and 3000 mg/kg/day
Results/Conclusion: Maternal NOAEL 2000 mg/kg/day (effects on body weight were observed at 3000 mg/kg/day)
Embryo/Fetal NOAEL 3000 mg/kg/day; no signs of embryotoxicity, fetotoxicity or teratogenicity were noted at any dose-level.



Memorandum

TO: Lillian Gill, D.P.A.
Director - COSMETIC INGREDIENT REVIEW (CIR)

FROM: Beth A. Lange, Ph.D.
Industry Liaison to the CIR Expert Panel

DATE: September 3, 2014

SUBJECT: Comments on the Draft Report on PEGylated Alkyl Glycerides Prepared for the September 8-9, 2014 CIR Expert Panel Meeting

Key Issue

Composition and Impurities - The statement that industry “removes” 1,4-dioxane from ingredients before blending into cosmetic formulation is misleading and may lead to the expectation of “zero” 1,4-dioxane in these ingredients. As analytical methods improve, lower and lower concentrations of 1,4-dioxane can be measured. Please change the word “remove” to “limit”.

Additional Comments

Cosmetic Use - In the paragraph describing use of the PEG Glyceryl Cocoate ingredients in 1996, “PEG-6 glyceryl cocoate” needs to be corrected to “PEG-7 glyceryl cocoate” and “PEG-7 cocoate” needs to be corrected to “PEG-7 glyceryl cocoate”.

Irritation and Sensitization, Dermal - What concentration of PEG-7 Glyceryl Cocoate was mildly irritating to rabbits?

To which compound does “8 or 11% glyceryl cocoate” refer?

Irritation and Sensitization, Dermal and Summary - As there was a limit on the use of PEG Glyceryl Cocoate ingredients of 10%, the negative HRIPT (n=205) on the hair product containing 11% PEG-7 Glyceryl Cocoate should be mentioned in both the Irritation and Sensitization section and the Summary.

Table 4, Glycine Soja (Soybean) Oil - The linoleic row says “0000”

Table 7, Alternative Studies - Please correct: “polyclyceryl”



Memorandum

TO: Lillian Gill, D.P.A.
Director - COSMETIC INGREDIENT REVIEW (CIR)

FROM: Beth A. Lange, Ph.D.
Industry Liaison to the CIR Expert Panel

DATE: October 1, 2014

SUBJECT: Comments on the Tentative Report: Safety Assessment of PEGylated Alkyl Glycerides as Used in Cosmetics

Key Issues

The Dictionary definitions of the PEGylated alkyl glyceride ingredients all indicate that the number in the name represents the average number of moles of ethylene oxide. In contrast, the Abstract, Introduction, Chemistry and Summary sections all state that these ingredients “are mono-, di-, and/or tri-glycerides that have been modified with a specific number of ethylene glycol repeat units”. The word specific is not correct, as the number represents the average number of moles of ethylene oxide. In addition, in Table 1, the added text for each ingredient says “and the sum of all cases of n is [x]”. Because x represents the average number of moles of ethylene oxide, please add the word “average” before sum.

Irritation and Sensitization, Dermal, Summary - In addition to discussing studies that reported effects, the text should discuss the highest concentrations of these ingredients that were tested in HRIPTs and not found to be sensitizing. This is especially important for this report because the previous limit (10% in leave-on products) for the PEG Glyceryl Cocoate ingredients was based in part on the highest concentration found not to be sensitizing.

Additional Comments

Introduction, last paragraph - Please correct “*incarnate*” to “*incarnata*”

Table 1 - Since Figure 2 is of PEG Caprylic/Capric Glycerides, why is it necessary to repeat that figure 3 times in this table? Instead of repeating the figure, please refer to Figure 2 in a manner similar to the references to Figure 1 in the definitions of the other PEG alkyl glyceride ingredients.

Table 2 - Caprylic/Capric Triglyceride - Please revise: "A 4% solution in ethanol were not sensitizers in guinea pigs."

Corn Oil - In the Conclusion row, correct "glyceride" to "glycerides"

Safflower Oil - Please correct the last sentence: "The oil increased the incidence of 12-dimethylbenz(a)anthracene in rats." The oil likely had an effect on tumors induced by 12-dimethylbenz(a)anthracene.

PEGylated Oils - Did these compounds really have an effect on "skeletal function"? Or was the effect on "skeletal muscle function"?

Table 5 - PEG-7 Glyceryl Cocoate - As it is not known for sure if the "other fragrance preparation" containing 6% PEG-7 Glyceryl Cocoate is a spray, it should also have a footnote a.

PEG-10 Olive Glycerides - The deodorant product should be noted as "not spray"

PEG-45 Palm Kernel Glycerides - There is a body and hand product (not spray) containing 0.92% PEG-45 Palm Kernel Glycerides. CIR protocol indicates that this product should be in the "powder" row with a footnote that it may or may not be a powder product.

PEG-75 Shea Butter Glycerides - As it is not known for sure if the "perfume" containing 0.005% PEG-75 Shea Butter Glycerides is a spray, it should also have a footnote a.

Table 7 - Rather than stating that some *in vitro* assays use "MTT exposure", please state: "MTT was used to measure cell viability".

Last *in vitro* row - Please correct: "tyo be"

First Non-Human row - Please correct: ""well-define erythema"

Table 8 - Unless "Leghorn" and "DeKalb" represent species other than chicken, please state "chicken" with "Leghorn" and "DeKalb" (as was done for NZW rabbits).

Last row - It is not clear what "ocular irritation rate of 3/2000" represents. Is this an ocular irritation "score"? A rate generally represents the number of subjects of the total that reacted - but only 10 subjects were tested, so a "rate of 3/2000" does not make sense.